



Ethnomedicinal plants used to treat skin diseases by Tharu community of district Udham Singh Nagar, Uttarakhand, India



Jyotsana Sharma^{a,*}, Sumeet Gairola^{b,*}, Yash Pal Sharma^a, R.D. Gaur^c

^a Department of Botany, University of Jammu, Jammu-Tawi 180006, Jammu and Kashmir, India

^b Herbarium and Plant Systematics Section, Biodiversity and Applied Botany Division, CSIR-Indian Institute of Integrative Medicine, Jammu-Tawi 180001, Jammu and Kashmir, India

^c Department of Botany, HNB Garhwal University, Srinagar Garhwal 246174, Uttarakhand, India

ARTICLE INFO

Article history:

Received 12 March 2014

Received in revised form

29 September 2014

Accepted 4 October 2014

Available online 14 October 2014

Keywords:

Antioxidant

Antimicrobial

Anti-inflammatory

Toxicological

Wound healing

Tyrosinase inhibition

ABSTRACT

Ethnopharmacological relevance: Tharu community is the largest primitive indigenous community of the Uttarakhand, India. In this article we have scientifically enumerated medicinal plants and herbal preparations used by the Tharu community to treat various skin diseases, and discussed dermatological properties of these plants in the light of previous ethnomedicinal, microbiological, pharmacological, toxicological, phytochemical and clinical studies.

Materials and methods: Ethnomedicinal survey was conducted in different villages of Tharu community located in district Udham Singh Nagar, Uttarakhand, India. Ethnomedicinal information on plants used to treat various skin diseases was collected from 122 individuals (93 males and 29 females), including 35 experienced herbal practitioners and 87 local villagers. For each of the recorded plant species the use value (UV) and fidelity level (FL) was calculated. The informant consensus factor (F_{ic}) was also calculated to find out the homogeneity in the information given by the informants.

Results: A total of 90 plant species belonging to 86 genera and 48 families were used by the Tharu community to treat various skin diseases viz., wounds (38 spp.), boils (32 spp.), cuts (18 spp.), leprosy (11 spp.), eczema (10 spp.), itching (7 spp.), ringworm (5 spp.), burns (4 spp.), leucoderma (4 spp.), cracked heels (2 spp.), dandruff (3 spp.), body infection (2 spp.), chilblains (2 spp.), hair fall (2 spp.) and toes infection (2 spp.). Information on botanical name, family, vernacular name, ailments treated, mode and dose of herbal preparations, UV and FL values are provided for each of the recorded species. According to UV value most preferred plant species used to treat skin diseases by Tharu community was *Ricinus communis* L. followed by *Tridax procumbens* (L.) L., *Azadirachta indica* A. Juss., *Ageratum conyzoides* and *Allium cepa* L.

Conclusions: The present study has revealed significant information on various medicinal plants used to treat skin diseases by Tharu community. Literature review has confirmed most of the claims made by the Tharu community regarding treatment of various skin diseases by the reported plants. The literature review has also revealed that products from very few of the reported plants are available in market, while most of the reported plants are still under preclinical or clinical trials. There are various known phytochemicals, and antibiotic, antibacterial, antiviral and antifungal agents present in these plants

Abbreviations: AAPE, arachidonic acid (AA) induced paw edema model; ABTS, 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) assay; AIA, anti-inflammatory activity; AOTS, acute oral toxicity study; APE, adjuvant induced paw edema model; BKPE, Bradykinin induced paw edema model; BWI, burn wound model; C, cultivated; CAP, carrageenan air pouch model; CFAA, complete Freund's adjuvant (CFA) induced arthritis model; CHG, chlorhexidine gluconate; CIP, carrageenan induced peritonitis model; CODE, croton oil induced dermatitis of the mouse ear model; CPE, carrageenan induced paw edema model; CPG, cotton pellet granuloma model; CTS, chronic toxicity study; DPE, dextran induced paw edema model; DPPH, 2,2-diphenyl-1-picrylhydrazyl assay; DSW, dead space wound model; DXM, dexamethasone; EWM, excision wound model; F_{ic} , consensus factor; FL, Fidelity level; FoPE, formaldehyde induced paw edema model; FPE, formaline induced paw edema model; FRA, framycetin; FRAP, ferric reducing antioxidant powder assay; FSC, framycetine sulfate cream; H, herb; HPE, histamine induced paw edema model; HRBC, human blood cell membrane stabilization method; HTPE, 5-hydroxytryptamine induced paw edema model; IPE, immunologically induced paw edema model; IWM, incision wound model; KPE, kaolin induced paw edema model; LPE, leukotriene induced paw edema model; NA, not available; NEO, neosporin; NFZ, nitrofurazone; NPE, nystatin induced paw edema model; NT/NM, non-toxic/no mortality; PAPE, phlogistic agents induced paw edema model; PGNIN, peptidoglycan (PGN) induced inflammatory reaction; PGPE, prostaglandin E2 induced paw edema model; PIW, punch incision wound model; PPCIN, polyinosinic: polycytidylic acid (polyI:C) induced inflammatory reaction; PVI, povidone iodine ointment; RP, reducing power; S, shrub; SATS, sub acute toxicity study; SOF, soframycin; SPE, serotonin induced paw edema model; SSD, silver sulphadiazine; ST, sulphathiazole; TAC, total antioxidant capacity; TC, number of times cited in the reviewed literature to treat any skin disease in India; TPAP, tetradecanoylphorbol acetate (TPA) induced paw edema model; Tr, tree; UV, use value; W & C, both wild as well as cultivated; W, wild; XEE, xylene induced ear edema model; ZNO, zinc oxide

* Corresponding authors.

E-mail addresses: sharmajyotsana76@gmail.com (J. Sharma), sumeetgairola@iiim.ac.in (S. Gairola).

which may be synthesized or transformed to make pharmaceuticals. Some of the reported plants have shown promising results in preclinical trials and there is a need of clinical trials to see their safety and efficacy in treating various skin diseases. These plants may be targeted for development of new medicines, ointments or drugs for the treatment of skin diseases. However further toxicological, preclinical and clinical studies are needed to validate claims about little worked out plant species reported in the present study viz., *Sida cordata* (Burm. F.) Borss. Waalk., *Milletia extensa* (Benth.) Baker, *Caesulia axillaris* Roxb., *Ehretia laevis* Roxb., *Vanda tessellate* (Roxb.) Hook. Ex G.Don. and *Eualaliopsis binata* (Retz.) C.E. Hubb. Further studies on these plants are recommended to assess their potential in development of new skin care products.

© 2014 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

The human skin is the outer covering of the body that provides contact with the environment and protects human body from unfavorable external factors. According to Grice et al. (2009), “Human skin is a large, heterogenous organ that protects the body from pathogens while sustaining microorganisms that influence human health”. Skin ailments affect people of all sex and ages, right from the neonate to the elderly stage, causing harm to human health in number of ways. It has been estimated that skin diseases amount to as high as 34% of all occupational diseases (Spiewak, 2000; Abbasi et al., 2010a). It is supposed to be the most common ailment amongst rural population (Policepatel and Manikrao, 2013). Skin diseases or infectious dermatological conditions are common in tropical countries where most of the population live in underdeveloped areas with lack of sanitation and inattentiveness to hygienic food habits. Skin diseases such as ringworm, itching, wound, skin disorders, leprosy, dermatitis, skin allergy, eczema, psoriasis, scabies and swelling are caused by a variety of micro-organisms and uncomfortable environment (Suresh et al., 2012). Traditional herbal medicines have played an important role in the management of dermatological conditions (Saikia et al., 2006). Hundreds of medicinal plants worldwide are used in the traditional medicine for treatment of skin diseases caused by bacteria, fungi and viruses (Kumar and Vidyasagar, 2008).

Uttarakhand Himalaya is a rich reservoir of biodiversity including enormous plants species and wildlife. It is a house to many indigenous communities like Bhotias, Boxas, Gujjars, Tharus, Rajis and Jaunsaries, who have excellent traditional knowledge on medicinal plants used to treat various human health problems. Apart from these, many other forest dwelling communities and local people also possess unique knowledge on uses of medicinal plants available in the region. Several researchers have documented ethnomedicinal plants used for treating various human health related ailments by different indigenous communities in sub-Himalayan region of Uttarakhand (Gaur, 1999; Gaur et al., 2010; Gaur and Sharma, 2011; Sharma and Painuli, 2011; J. Sharma et al., 2012; Sharma et al., 2013a, 2013b; Gairola et al., 2013). Although, Tharu community is the largest primitive indigenous community of the Uttarakhand, but very little information is available about ethnomedicinal plants used by this community. Recently J. Sharma et al. (2011, 2012) and Sharma et al. (2013b) have recorded some ethnomedicinal information on plants used by Tharu community to treat some general diseases, jaundice and epilepsy, respectively. But information about plants used to treat various skin diseases by Tharu community of Udham Singh Nagar is almost lacking except some preliminary information recorded by J. Sharma et al. (2011) about plants used to treat wounds (10 plants) and cuts (2 plants). Therefore, it was realised that the documentation of ethnomedicinal knowledge of this community is required. Keeping the aforesaid facts in view, the present study was undertaken to scientifically enumerate medicinal plants and herbal preparations used by Tharu community to treat various skin diseases, and discuss dermatological properties of these plants in the light of

previous ethnomedicinal, microbiological, pharmacological, toxicological, phytochemical and clinical studies.

2. Research strategy and methods

2.1. Study area

The Uttarakhand state is situated in the northern part of India between 28°43' and 31°28'N latitude and 77°34' and 81°03'E longitude, and shares an international boundary with Tibet Autonomous Region in the north and Nepal in the east (Fig. 1). It embodies mountainous terrain ranging from 300 to 7800 m asl and has a geographic area of 53,483 km², which is even bigger in size than nearby Himalayan country of Bhutan. The state has very rich biological wealth and cultural heritage, and because of its unique geography and diverse climatic conditions it possesses the large number of medicinal plant species. Majority of the land area in Uttarakhand is mountainous (93%) and 64.79% area is covered by the forests (FSI, 2011). The present study was conducted in different villages inhabited by Tharu community in district Udham Singh Nagar of Uttarakhand, popularly known as the “Gateway to Kumaon hills”. There are three main sub-divisions of Udham Singh Nagar – Rudrapur, Kashipur and Khatima. The district consists of seven tehsils viz., Bajpur, Gadarpur, Jaspur, Kashipur, Kichha, Khatima and Sitarganj. District is located in the Terai region and is endowed with a very rich fertile land. Agriculture is the mainstay and there are several agriculture related activities and industries located here. It is surrounded by Nainital district in the north, Bijnour, Moradabad and Rampur in the west, Bareilly and Pilibhit in the south and district Champawat in the east. It also shares south east boundary with the international border of the Nepal. Reserved forest area lies at the borders of district Nainital and Champawat. The villages visited in district Udham Singh Nagar to collect ethnomedicinal information were Bawanpuri, Beeriya, Bhagora, Bhagori, Bharuni, Bidora–Majola, Chamarpur, Chaumela, Dhohra–Pajaniya, Enjania, Gaganpur, Gaurikhedra, Ghosari, Harriaya, Karkata, Kharoona, Khujri, Kicha, Lamkhera, Lauka, Magarsanda, Malara, Nadai, Nakulia–Bithora, Paseni, Sadhunagar, Sehjani, Sisalkhera, Sisoona, Sitargang, Taumala, Tisoor and Tukari–Bichua.

2.2. Studied community and ethnomedicinal survey

Frequent field trips were undertaken in different seasons viz., winter (November–March), summer (April–June) and rainy (July–October) in order to collect information on ethnomedicinal plants used to treat various skin diseases by Tharu community. Tharu community is the largest primitive indigenous community of the Uttarakhand, who lives in interior forests sustaining a close association with their ambient environment. They are shy and relatively timid people due to lack of exposure outside their community. Tharus have Mongoloid facial features and a unique




Fig. 1. Map of the study area.

dialect. According to some people, the word “Tharu” is derived from “Taru” which means trees or forest, as these people generally reside in forests. Whereas, according to others the word “Tharu” is derived from Sanskrit word “Thal” (ground), as they are settled in the forest ground in the terai region. According to Crooke (1896), one of the Kshatriya kings of plain named these people “Tharu” (wine bibber) because of their drunken habit. The main occupation of Tharu people is agriculture. Extraction of fibers and threshing of crops is common among these people. The social life style patterns of Tharus are similar in many ways with nearby Hindu societies. They follow Hinduism religion and worship all the Hindu God and Goddess. Besides that, they also worship the tribal God or indigenous deities such as “Bhumisen Devta” or “Bhumisen”,

“Bhuniya” and others. The Tharu community has a strong belief in witchcraft (Jhar-phoonk and Tantra-mantra). A local doctor or Vaidhya, locally known as “Bharara” or “Bharra” is found in each village. These medicinal practitioners have very good knowledge of traditional herbal therapies and village folk rely on them for recovering from diseases.

Ethnomedicinal information on plants used to treat various skin diseases was collected from 122 individuals (93 males and 29 females), including 35 experienced herbal practitioners and 87 local villagers. Verbal consent from the participants was taken and they were clearly informed about the objective of the study and their right to withdraw from the study at any time. Participants were specifically asked about plants and plant parts used for the

treatment of various skin diseases along with method of preparation and mode of administration of the prescribed medicine. Interviews were conducted in local language and according to symptoms described by participants, the skin diseases were classified in to 15 categories viz., boils, cuts, leprosy, eczema, itching, ringworm, burns, leucoderma, cracked heels, dandruff, body infection, chilblains, hair fall, toes infection and wounds.

The plant specimens were identified with the help of Flora of the District Garhwal North West Himalaya (Gaur, 1999). For the confirmation of identification the specimens were compared with the authentic specimens lodged at Herbaria of HNB Garhwal University Srinagar (GUH), Botanical Survey of India, Dehradun (BSD) and Forest Research Institute, Dehradun (DD). Identified specimens were properly labeled with detailed botanical name, family, locality, field number, date of collection and other necessary remarks and deposited in the internationally indexed Herbarium of Department of Botany, HNB Garhwal University Srinagar Garhwal (GUH). Valid botanical names along with the author citations of all the recorded plant species were verified from www.theplantlist.org version 1.1 (TPL, 2013).

2.3. Statistical analysis and literature review

For each of the recorded plant species the use value (UV), as adapted by Ferreira et al. (2009) from the proposal of Phillips et al. (2002) was calculated. This quantitative method evaluates the relative importance of each medicinal plant species based on its relative use among informants. UV was calculated using the following formula:

$$UV = \sum \frac{U}{n}$$

where U is the number of times a species is cited and n is the number of informants. The use value of each species is therefore based objectively on the importance attributed by the informants and does not depend on the opinion of the researcher (Ferreira et al., 2009). Fidelity level (FL) was calculated using the following formula:

$$FL(\%) = \frac{Ip}{Iu} \times 100$$

where Ip is the number of informants who independently indicated the use of a species for the same major ailment and Iu the total number of informants who mentioned the plant for any major ailment (Friedman et al., 1986). To test homogeneity of knowledge about the medicinal plants, the factor informant consensus (F_{ic}) was used (Heinrich et al., 1998; Gazzaneo et al., 2005). The F_{ic} was

calculated using the following formula:

$$F_{ic} = \frac{n_{ur} - n_t}{n_{ur} - 1}$$

where n_{ur} is the number of use reports for particular ailment category, and n_t is the number of species used for a particular ailment category by all the informants.

Detailed literature survey was carried out to gather and discuss information about the earlier ethnomedicinal studies recording uses of the listed plants in treatment of any skin disease, other studies showing their antimicrobial, anti-inflammatory, wound healing, toxicological, antioxidant and tyrosinase inhibitory properties and compound or molecules present in these plants which are known antibiotic, antibacterial, antiviral and antifungal agents. Literature was searched from various scientific databases viz., Web of Science, CAB international, Google Scholar, Science direct, SciFinder, PubMed, Scopus and DNP CHEMnetBASE.

3. Results and discussion

3.1. Ethnomedicinal plants used by Tharu community

The present exploration has brought in to light the antidermatophytic medicinal plants and herbal medicines used against various skin diseases by Tharu community living in Udham Singh Nagar district of Uttarakhand. A total of 90 plants belonging to 86 genera and 48 families were used by Tharu community to treat 15 categories of skin diseases (Fig. 2). Recently Gairola et al. (2014) have reported list of 321 medicinal plant taxa used to treat dermatological or skin related diseases by various indigenous communities of Jammu & Kashmir. Whereas, Sharma et al. (2013a) have also reported 109 medicinal plants used to treat various skin diseases by Gujjar community of sub-Himalayan tract of Uttarakhand, India. Fifty plant species were used to cure only one skin disease, whereas 40 plant species were used to cure multiple skin diseases. Results of the present investigation are given in Table 1, where information about reported plants species are given along with their botanical names, families, vernacular names, skin disease treated, mode and dose of herbal preparations, UV values, distribution, habitat, earlier ethnomedicinal studies in India citing use of plant for treatment of any skin disease and number of times species cited in the reviewed literature to treat any skin disease in India. Frequently used family in skin care by Tharu community was Leguminosae (11 taxa) followed by Lamiales (7 taxa), Apocynaceae (6 taxa), Euphorbiaceae (5 taxa), Poaceae (5 taxa), Asteraceae (4 taxa), Anacardiaceae (3 taxa),




Fig. 2. Number of plant species used to treat different categories of skin diseases.

Table 1

Plants and herbal preparations used as remedy of skin diseases by the Tharu community of district Udham Singh Nagar, Uttarakhand, India.

Plant species (family) voucher specimen number	Vernacular name (Habit)	Mode of preparation and dose	UV	Earlier ethnomedicinal studies in India citing use of plant for treatment of any skin disease	Distribution ^a ; Habitat (TC)	Uses by Gujjar Community of sub-Himalayan tract (Sharma et al., 2013a) ^b
Acanthaceae						
<i>Dicliptera paniculata</i> (Forssk.) L. Darbysh. [<i>*Peristrophe paniculata</i> (Forssk.) Brummitt] (GUH-JS 20154)	Harjodi (H)	The paste of leaves is applied topically on wounds.	0.016	Adhikari et al. (2010), Gaur (1999)	Throughout India, common along waste places, road sides and crop fields to 1400 m asl; W (2)	–
Acoraceae						
<i>Acorus calamus</i> L. (GUH-JS 20355)	Papri (H)	The leaf paste is applied topically on wounds. Paste of rhizome is mixed with <i>Curcuma domestica</i> rhizome and applied topically on eczema.	0.074	Chopda and Mahajan (2009), Kingston et al. (2009), Pradhan and Badola (2008), Shiddamallayya et al. (2010), Sivaranjani and Ramakrishnan (2012), M. Yadav et al. (2012)	Throughout India in damp marshy places, ascending to 1800 m asl; W & C (6)	–
Amaranthaceae						
<i>Achyranthes aspera</i> L. (GUH-JS 18893)	Chattisa/ Chircita (H)	The leaves are crushed and juice is applied topically on boils.	0.336	Bhandary and Chandrashekhar (2002), Bhat et al. (2012), Bhat et al. (2014), Chopda and Mahajan (2009), Gaur and Sharma (2011), Johnsy et al. (2012), Kumar and Vidyasagar (2008), Muthu et al. (2006), Nandagopalan et al. (2011), Patil et al. (2009), Ray et al. (2011), Revathi and Parimelazhagan (2010), Seetharam et al. (1999), J. Sharma et al. (2011), Sharma et al. (2013a), A.K. Sharma et al. (2013), Shiddamallayya et al. (2010), Singh and Maheswari. (1994), Sinha et al. (2013), Sivaranjani and Ramakrishnan (2012), Subramanian et al. (2011), Venkataswamy et al. (2010), Wadankar et al. (2011)	Throughout tropical and subtropical region of India, up to 2100 m asl; W (23)	The paste of leaves is applied externally on skin allergy.
<i>Amaranthus spinosus</i> L. (GUH-JS 18832)	Katmarsa (H)	The leaves are pounded and are applied externally on wounds between toes in rainy season. The paste of leaves is mixed with turmeric powder and applied topically on boils. Juice of leaves is applied topically on eczema.	0.311	Adhikari et al. (2010), Choudhary et al. (2011), Sharma et al. (2013a), A.K. Sharma et al. (2013)	Native to tropical America. Common weed in cultivated fields, waste places and along roadsides; W (4)	The paste of roots is applied on boils and carbuncle.
Amaryllidaceae						
<i>Allium cepa</i> L. (GUH-JS 20356)	Palandu (H)	The whole plant is crushed and its paste is applied on ringworm.	0.500	Abbasi et al. (2010a), Bhat et al. (2012), Gaur (1999), Sharma et al. (2013a)	Cultivated throughout India, up to 1800 m asl in Himalaya; C (4)	The bulbs are pounded and paste is applied externally on skin allergy.
Anacardiaceae						
<i>Buchanania cochinchinensis</i> (Lour.) M.R. Almeida [<i>*Buchanania</i>	Chironji (Tr)	The bark paste is applied topically on cuts and burns. Decoction of bark is taken internally for the same, one teaspoonful, twice a day until the disease cures.	0.090	Gaur (1999), Kala (2009), Kumar and Vidyasagar (2008), Sharma et al. (2013a), Singh et al. (2002)	Drier parts of India; W (5)	Seed powder is mixed with milk and applied on pimples. The paste of leaves is applied over wounds.

<i>lanzan</i> Spreng.] (GUH-JS 20322)						
<i>Lannea coromandelica</i> (Houtt.) Merr. (GUH-JS 18876)	<i>Jingan</i> (Tr)	The juice of leaves is applied topically on cuts and wounds.	0.197	Adhikari et al. (2010), Dahare and Jain (2010), Sharma et al. (2013a), Singh et al. (2002), Upadhyay et al. (2010)	Throughout India, ascending to 1500 m asl in the Himalayas; W (5)	The paste of bark is applied externally on wounds.
<i>Semecarpus anacardium</i> L.f. (GUH-JS 20261)	<i>Bhilwa</i> (Tr)	The gum is applied topically on leprosy.	0.025	Bhat et al. (2012), Kumar et al. (2012), Pradhan and Badola (2008), Singh et al. (2002), Wadankar et al. (2011)	Sub Himalayan tracts from Himachal Pradesh to Sikkim, Punjab, Assam, Khasi Hills, Madhya Pradesh and Peninsular India; W (5)	–
Annonaceae						
<i>Annona squamosa</i> L. (GUH-JS 20297)	<i>Sitaphal</i> (Tr)	The leaves are ground and its paste is applied topically on boils. The powder of seeds is applied on scalp to treat dandruff.	0.131	Bhat et al. (2012), Chopda and Mahajan (2009), Choudhary et al. (2011), Dash and Misra (1999), Gaur (1999), Gaur and Sharma (2011), Nayak et al. (2004), Raja et al. (2009), J. Sharma et al. (2011), A.K. Sharma et al. (2013), Subramanian et al. (2011), Wadankar et al. (2011)	Native to South America and the West Indies, now cultivated throughout India; C (12)	–
Apocynaceae						
<i>Calotropis procera</i> (Aiton) Dryand. (GUH-JS 20351)	<i>Madar</i> (S)	The leaves and flowers are crushed together, made into paste and applied topically on boils.	0.279	Choudhary et al. (2011), Gangwar et al. (2010), Kumari and Singh (2009), J. Sharma et al. (2011), Sharma et al. (2013a), Upadhyay et al. (2010)	An evergreen shrub distributed throughout India; W (6)	The roots are ground and applied externally on leucoderma.
<i>Cryptolepis dubia</i> (Burm.f.) M.R. Almeida [* <i>Cryptolepis buchananii</i> Roem. & Schult.] (GUH-JS 19730)	<i>Dudhi bel</i> (H)	The latex of the plant is applied topically on wounds, boils or sores	0.066	Adhikari et al. (2010), Bhat et al. (2012), Bhat et al. (2014), Mairh et al. (2010), Subramanian et al. (2011)	Throughout India, ascending to 1200 m asl in Himalaya; W (5)	–
<i>Holarrhena pubescens</i> Wall. ex G.Don (GUH-JS 18830)	<i>Kurchi</i> (Tr)	The decoction of seeds is recommended internally to cure leprosy. The paste of leaves is applied topically on boils.	0.279	Bhat et al. (2012), Bhat et al. (2014), Gangwar et al. (2010), Girach et al. (1999), J. Sharma et al. (2011)	Throughout India, tropical Himalayas going up to an altitude of 1100 m asl; W (5)	–
<i>Rauvolfia serpentina</i> (L.) Benth. ex Kurz (GUH-JS 18866)	<i>Jhaberbarua</i> (H)	Root paste is topically applied for boils.	0.025	Bhat et al. (2012), Bhat et al. (2014), Gaur et al. (2011), Sharma et al. (2013a)	The sub-Himalayan tract from Punjab to Nepal, Sikkim, Assam, Western Ghats and Andamans; W (4)	The root paste is mixed with oil of <i>Cinamomum tamala</i> (tejpaat) and externally applied on leucoderma in night.
<i>Vallisneria spiralis</i> (L.) Kuntze (GUH-JS 19763)	<i>Bakerghaneli</i> (S)	The latex is topically applied on wounds.	0.049	Adhikari et al. (2010), Gaur (1999), Sharma et al. (2013a)	Throughout India, Cultivated in gardens; W & C (3)	The plant paste is applied on eczema. The latex is applied on wounds.
<i>Wrightia arborea</i> (Dennst.) Mabb. (GUH-JS 20372)	<i>Dhudli</i> (Tr)	The latex is applied topically to heal on cuts and wounds.	0.148	Gaur (1999), Reddy et al. (2007), Sharma et al. (2013a)	Sub Himalayan tracts, Punjab, Rajasthan, Bihar, Assam and Western Peninsula; W (3)	The latex is applied on cuts, wounds and on skin allergy.
Asteraceae						
<i>Ageratum conyzoides</i> (L.) L. (GUH-JS 18868)	<i>Gindhoni</i> (H)	The leaf juice is applied topically on cuts to stop bleeding. The whole plant decoction is taken orally to cure leprosy, 2 ml thrice a day.	0.541	Gaur (1999), Sharma et al. (2010), Sharma et al. (2013a), M. Yadav et al. (2012)	Throughout India, as a weed up to an altitude of 1800 m asl; W (4)	The leaf juice is applied externally on cuts to stop bleeding.
<i>Caesulia axillaris</i> Roxb. (GUH-JS 20210)	<i>Gorghanta</i> (H)	The paste of flowers is applied topically on cuts and wounds.	0.180	J. Sharma et al. (2011)	Throughout the warmer parts of India, ascending to 1000 m asl; W (1)	–
<i>Eclipta prostrata</i> (L.) L. (GUH-JS 18829)	<i>Dudhli</i> (H)	The juice of whole plant is applied topically on cuts, wounds and boils	0.344	Adhikari et al. (2010), Bhat et al. (2014), Chopda and Mahajan (2009), Kamble et al. (2010)	Throughout India, up to 2000 m asl on the hills; W (4)	–
	<i>Mundi</i> (H)		0.623			

Table 1 (continued)

Plant species (family) voucher specimen number	Vernacular name (Habit)	Mode of preparation and dose	UV	Earlier ethnomedicinal studies in India citing use of plant for treatment of any skin disease	Distribution ^a ; Habitat (TC)	Uses by Gujjar Community of sub-Himalayan tract (Sharma et al., 2013a) ^b
<i>Tridax procumbens</i> (L.) L. (GUH-JS 18817)		The paste of leaves is topically applied on cuts and wound to stop bleeding. The juice of leaves is applied on boils.		Gaur (1999), Gaur et al. (2011), Jyothi et al. (2010), Kamble et al. (2010), Kumar and Vidyasagar (2008), Muthu et al. (2006), J. Sharma et al. (2011), Sharma et al. (2013a), Singh et al. (2002), Sinha et al. (2013), Upadhyay et al. (2010)	Waste places, road sides and hedges throughout India; W (11)	The paste of leaves is applied on cuts and wound to stop bleeding.
Bignoniaceae						
<i>Oroxylum indicum</i> (L.) Kurz (GUH-JS 20273)	Phargat (Tr)	The bark powder is used topically on wounds. Bark paste is applied topically on burns.	0.033	Gaur and Sharma (2011)	Throughout the greater part of India to 1500 m asl; W (1)	–
Boraginaceae						
<i>Ehretia laevis</i> Roxb. (GUH-JS 20101)	Chamror/ Chamrod (Tr)	The leaves are ground and its paste is topically applied on wounds	0.041	Gaur and Sharma (2011), Meena and Yadav (2011)	Throughout India, also grown along roadsides; W & C (2)	–
Brassicaceae						
<i>Brassica juncea</i> (L.) Czern. (GUH-JS 20302)	Lahi (H)	The paste of seeds is applied topically used on boils.	0.303	Bhat et al. (2012), Sharma et al. (2013a), A.K. Sharma et al. (2013)	Cultivated in Punjab, West Bengal, Uttar Pradesh and Gujarat; C (3)	The paste of seeds is externally used on boils and skin allergy.
Cannabaceae						
<i>Cannabis sativa</i> L. (GUH-JS 20224)	Bhang (S)	The leaves are fried with mustard oil, made into paste and applied topically on burns.	0.295	Chopda and Mahajan (2009), Negi et al. (2011), Sharma et al. (2004), Sharma et al. (2010)	Cultivated all over the country. Commonly occurs in waste grounds, along road side; W & C (4)	–
Celastraceae						
<i>Celastrus paniculatus</i> Willd. (GUH-JS 19734)	Kakundan (S)	The seed paste is applied topically on leprosy.	0.066	Adhikari et al. (2010), Bhat et al. (2014), Chakraborty and Bhattacharjee (2006), Pradhan and Badola (2008), Sharma et al. (2013a)	Sub-Himalayan tract up to 2000 m asl and South Indian hills; W (5)	The paste of leaves is applied externally on eczema. The ash of roots is mixed with fruit piece of <i>Citrus pseudolimon</i> (galgal) and placed in an earthen pot for a whole night; the whole material is dried in shade in early morning and made into paste, this preparation is given orally (2–3 gm), twice a day for 10–15 days with cold water to treat body allergy, milk is avoided during treatment.
Cleomaceae						
<i>Cleome viscosa</i> L. (GUH-JS 20170)	Hulhul (H)	The whole plant paste is applied topically on wounds. The paste of seeds is applied on scalp during hair fall or to avoid baldness.	0.107	Adhikari et al. (2010), Ayyanar and Ignacimuthu (2009), Ganesan and Kesaven (2003), Gaur (1999), Girach et al. (1999), Kuvar and Bapat (2010), Muthu et al. (2006), Upadhyay et al. (2010)	Throughout India, Tripura, West Bengal and Gangetic valley as a weed; W (8)	–
Combretaceae						
	Kawa (Tr)	The paste of bark is applied topically on leprosy.	0.033			–

<i>Terminalia arjuna</i> (Roxb. ex DC.) Wight & Arn. (GUH-JS 20334)				Kumari and Singh (2009), Muthu et al. (2006), Shiddamallayya et al. (2010)	Throughout the greater part of India, also grown as an avenue tree; W & C (3)	
Commelinaceae						
<i>Commelina benghalensis</i> L. (GUH-JS 20160)	Kana (H)	The paste of whole plant is applied topically on wounds	0.025	Adhikari et al. (2010), Ayyanar and Ignacimuthu (2009), Begum and Nath (2000), Gaur (1999), Lal et al. (1996), Sharma et al. (2013a)	Throughout India in moist places; W (6)	The paste of plant is applied externally on wounds. The juice of leaves (one teaspoonful) is taken internally twice a day for 1–2 weeks, in the treatment of leprosy.
Convolvulaceae						
<i>Cuscuta reflexa</i> Roxb. (GUH-JS 19749)	Sarai-babiya (H)	The paste of whole plant is applied topically to treat eczema. The whole plant paste is applied on scalp to treat dandruff.	0.189	Gaur (1999), Jyothi et al. (2010), Sharma et al. (2013a), Sinha et al. (2013), Upadhyay et al. (2010)	A parasitic climber common throughout India up to 3000 m asl; W (5)	The paste of plant is applied on leprosy.
<i>Ipomoea carnea</i> Jacq. (GUH-JS 20138)	Beshram (H)	The leaves are fried in <i>Brassica campestris</i> oil (sarsoon) and tied topically on cuts and wounds.	0.377	Adhikari et al. (2010), Sharma et al. (2013a), Singh and Maheswari (1994)	Native to tropical America, widely naturalized in India, common on waste places; W (3)	The leaves are fried in <i>Brassica campestris</i> oil (sarsoon) and tied externally on cuts and wounds to get quick relief.
Cucurbitaceae						
<i>Momordica dioica</i> Roxb. ex Willd. (GUH-JS 20237)	Karauna (H)	The paste of roots is applied topically on boils. Paste of fruit pulp is also used for the same.	0.107	Bhat et al. (2014), Sharma et al. (2013a)	Throughout India, up to 1500 m asl in the Himalaya; C (2)	The paste of roots is applied on boils.
Dipterocarpaceae						
<i>Shorea robusta</i> Gaertn. (GUH-JS 20369)	Kurela (Tr)	The gum is applied topically on cracks of feet's.	0.041	Gaur and Sharma (2011), Singh et al. (2002)	North, east and central India; W (2)	–
Euphorbiaceae						
<i>Acalypha indica</i> L. (GUH-JS 18853)	Kuphi (H)	The paste of leaves is mixed with little quantity of salt and applied topically on eczema. The paste of leaves is mixed with turmeric and topically on affected places to heal wounds.	0.230	Ayyanar and Ignacimuthu (2009), Chopda and Mahajan (2009), Johnsy et al. (2012)	Occurs throughout the plains of India, ascending the hills in Orissa up to 210 m asl; W (3)	–
<i>Euphorbia hirta</i> L. (GUH-JS 18829)	Dudhli (H)	The latex of the plant is applied topically on wounds. Latex is also applied on eczema. The paste of whole plant is applied topically on ringworm.	0.156	Ayyanar and Ignacimuthu (2009), Gaur (1999), Gaur et al. (2011), Girach et al. (1999), Jain et al. (2005), Katewa and Galav (2005), Kingston et al. (2009), Kumar et al. (2012), Kumari and Singh (2009), Muthu et al. (2006), Sharma et al. (2013a), A.K. Sharma et al. (2013), Sinha et al. (2013), Sivaranjani and Ramakrishnan (2012), Upadhyaya et al. (1998), M. Yadav et al. (2012)	Throughout warmer regions of India; W (16)	The plant paste is applied externally on cuts and wounds.
<i>Euphorbia thymifolia</i> L. (GUH-JS 20374)	Dhudiya (H)	The whole plant paste is used topically for the treatment of itches.	0.033	Shiddamallayya et al. (2010)	Found in tropical plains and low hills of India; W (1)	–
<i>Mallotus philippensis</i> (Lam.) Mull.Arg. (GUH-JS 20148)	Rohini (Tr)	The juice of leaves is applied topically on cuts and wounds.	0.066	Gaur et al. (2011), Kamble et al. (2010), Sharma et al. (2010), Sharma et al. (2013a), Singh and Singh (2009), Verma and Chauhan (2007), M. Yadav et al. (2012)	Throughout tropical regions of India; W (7)	The powder of fruits is mixed with ghee, made into paste and externally applied on boils.
<i>Ricinus communis</i> L. (GUH-JS 18877)	Arandi (S)	The leaves are fried in oil of mustard, made into paste and topically applied on wounds. The seed paste is applied topically on heel cracks. Seed oil is used to cure eczema.	0.631	Gangwar et al. (2010), Gaur et al. (2011), Johnsy et al. (2012), Kala (2009), Kumar et al. (2012), Kumar and Vidyasagar (2008), Sharma et al. (2013a), Uniyal and Shiva (2005), Upadhyay et al. (2010)	Native of tropical Africa, Found throughout India. Cultivated chiefly in Andhra Pradesh, Maharashtra, Karnataka and Orissa; W & C (9)	The leaves are fried in oil of mustard, made into paste and externally applied on wounds.

Table 1 (continued)

Plant species (family) voucher specimen number	Vernacular name (Habit)	Mode of preparation and dose	UV	Earlier ethnomedicinal studies in India citing use of plant for treatment of any skin disease	Distribution ^a ; Habitat (TC)	Uses by Gujjar Community of sub-Himalayan tract (Sharma et al., 2013a) ^b
Hypoxidaceae						
<i>Curculigo orchoides</i> Gaertn. (GUH-JS 18861)	Musli (H)	The juice of leaves is squeezed on cuts. Juice of leaves is also used for treating itching.	0.066	Ray et al. (2011), Sharma et al. (2013a), Singh et al. (2002)	Sub-tropical Himalayas from Kumaon eastward, Western Ghats from Konkan Southwards; W (3)	The paste of leaves is applied externally on skin allergy.
Lamiaceae						
<i>Anisomeles indica</i> (L.) Kuntze (GUH-JS 18814)	Basinga (S)	The paste of stem is mixed with coconut oil (<i>Cocos nucifera</i>) and applied topically on wounds.	0.262	Adhikari et al. (2010), Sharma et al. (2010)	Throughout India, ascending to 1500 m asl in Himalayas; W (2)	–
<i>Callicarpa macrophylla</i> Vahl (GUH-JS 19733)	Tilkhi/Dehgala (S)	The juice of leaves applied topically on chilblains.	0.041	Arya and Agarwal (2008), Sharma et al. (2013a)	Sub-Himalayan tracts, from Hazara eastwards to Assam, up to 1500 m asl; W (2)	The paste of leaves applied externally on chilbalins.
<i>Clerodendrum infortunatum</i> L. [<i>*Clerodendrum viscosum</i> Vent.] (GUH-JS 18892)	Bhatar (S)	The leaf juice is applied topically on cuts.	0.074	Gaur (1999), Jain et al. (2005), Johnsy et al. (2012), J. Sharma et al. (2011), Vijayan et al. (2007), Bhat et al. (2014), Bhat et al. (2012)	Throughout India; W (7)	–
<i>Hyptis suaveolens</i> (L.) Poit. (GUH-JS 20323)	Vantulsi (H)	The paste of leaves is applied topically on cuts and wounds. Juice of leaves is also used for treating itching.	0.098	Kumar and Vidyasagar (2008)	Native to tropical America. Distributed throughout India; W (1)	–
<i>Leonotis nepetifolia</i> (L.) R.Br. (GUH-JS 18887)	Hijurchi (H)	The paste of leaves is applied topically on ringworm.	0.066	Dash and Misra (1999), Kumar and Vidyasagar (2008), Kuvar and Bapat (2010), Sharma et al. (2013a), Upadhyaya et al. (1998)	Throughout the warmer parts of India; W (5)	The inflorescence is pounded and its paste is applied on skin allergy.
<i>Premna mollissima</i> Roth [<i>*Premna mucronata</i> Roxb.] (GUH-JS 20368)	Baka (S)	The decoction of bark (2 teaspoonfuls) is taken orally twice a day as a remedy for leprosy and its paste is also applied externally for the same. The latex is applied topically on boils.	0.156	Adhikari et al. (2010), Gaur (1999)	Submontane Himalaya in Garhwal to Sikkim, Peninsular India, Bihar, West Bengal and North-eastern India; W (2)	The paste of bark is applied externally on ringworm.
<i>Vitex negundo</i> L. (GUH-JS 18836)	Nirgndi (S)	The paste of leaves is used for healing of wounds. The paste of flowers is topically applied on eczema.	0.475	Bhat et al. (2012), Gaur et al. (2011), Ghorband and Biradar (2011), Prakasha and Krishnappa (2006), J. Sharma et al. (2011), Sharma et al. (2013a), Subramanian et al. (2011), Verma and Chauhan (2007)	Throughout India in the warmer zones; ascending to 900 m asl in the North-western Himalaya; W (8)	The leaves are fried in mustard oil, made in to paste and applied on boils.
Lauraceae						
<i>Litsea glutinosa</i> (Lour.) C.B.Rob. (GUH-JS 18823)	Maida lakri (Tr)	The bark paste is also applied topically on boil.	0.025	Kumar et al. (2012), Parhar and Biswas (1998), Ramadas et al. (2000), Sharma et al. (2013a)	Throughout the warmer parts of India, sub-montane Himalaya and South India; W (4)	The bark paste is applied on skin allergy.
Leguminosae						
<i>Albizia lebbek</i> (L.) Benth. (GUH-JS 19740)	Siris (Tr)	The juice of crushed leaves is applied topically on boils.	0.262	Chopda and Mahajan (2009), Joshi and Tyagi (2011), Kumari and Singh (2009), Sharma et al. (2010), Shiddamallayya et al. (2010)	All over India, from the plains up to 900 m asl in the Himalayas, also in the Andamans; W (5)	–
<i>Bauhinia variegata</i> L. (GUH-JS 19780)	Kachnar (Tr)	The root bark decoction is orally taken thrice a day to treat leprosy.	0.057	Negi et al. (2011), Singh et al. (2002), Upadhyay et al. (2010)	Throughout the warmer parts of India up to 1200 m asl, Punjab, Western Peninsula and Assam. Also cultivated in gardens; W & C (3)	–
<i>Butea monosperma</i> (Lam.) Taub. (GUH-JS 18802)	Palash (Tr)	The fresh gum is used as antiseptic and applied on cuts and wounds. Juice of flowers is used to cure itching and leprosy.	0.246	Bhat et al. (2014), Gaur (1999), Kala (2009), Katewa et al. (2004), Kumar et al. (2012), Kumari and Singh (2009), Singh et al. (2002)	Throughout India, up to 1200 m asl except in very arid regions; W (7)	–

<i>Cassia fistula</i> L. (GUH-JS 18842)	<i>Amaltas</i> (T)	The root paste is applied topically on leprosy. Powder of roots (2–3 gm) is taken internally two or three times a day to cure allergy or itching.	0.279	Bhat et al. (2012), Bhat et al. (2014), Gaur (1999), Johnsy et al. (2012), Jyothi et al. (2010), Kumar et al. (2012), Kumar and Vidyasagar (2008), Kumari and Singh (2009), Mamatha et al. (2006), Rajakumar and Shivanna (2009), Sharma et al. (2013a), A.K. Sharma et al. (2013), Shiddamallayya et al. (2010), Shivanna and Rajakumar (2010), Silja et al. (2008), Sinha et al. (2013), Verma and Chauhan (2007)	Cultivated as an ornamental throughout India; C (17)	Leaf paste is applied externally in ringworm.
<i>Crotalaria juncea</i> L. (GUH-JS 20362)	<i>Sanai</i> (S)	The seed paste is mixed with mustard oil and applied topically on wounds	0.033	Gaur and Sharma (2011), Shiddamallayya et al. (2010)	Throughout the warmer parts of India, especially in South India; W (2)	–
<i>Dalbergia sissoo</i> DC. (GUH-JS 20373)	<i>Siswa</i> (Tr)	Stem bark is soaked in water for 4–5 hours, made into paste and applied topically on leprosy.	0.148	Johnsy et al. (2012), Joshi and Tyagi (2011), Gaur and Sharma (2011), Sharma et al. (2013a), Gaur (1999)	The sub-Himalayan tract, up to 1200 m asl from Indus to Assam and in plains throughout India; W (5)	Leaves and bark are mixed together, made into paste and applied externally on eczema
<i>Desmodium gangeticum</i> (L.) DC. (GUH-JS 20191)	<i>Salperni</i> (H)	The paste of leaves is applied topically on boils.	0.033	Chopda and Mahajan (2009), Jain et al. (2005)	Common on lower hills and plains throughout India, ascending to 1500 m asl on the Himalaya; W (2)	–
<i>Milletia extensa</i> (Benth.) Baker (GUH-JS 20183)	<i>Bambiri</i> (S)	The root paste is applied topically on boils or sores.	0.033	Gaur (1999), Kamble et al. (2010), Sharma et al. (2013a)	Himachal Pradesh to Sikkim, Outer submontane Himalaya to 1500 m asl; W (3)	The bark paste is applied over wounds.
<i>Mucuna pruriens</i> (L.) DC. (GUH-JS 20180)	<i>Gaunj</i> (S)	The paste of seeds used topically on boils.	0.123	Gaur (1999), Majumdar and Datta (2007), Uniyal and Shiva (2005)	Throughout India, including Andaman and Nicobar Islands; W (3)	–
<i>Pongamia pinnata</i> (L.) Pierre (GUH-JS 18847)	<i>Papri</i> (Tr)	The paste of seeds is applied on wounds. The oil of seeds is applied topically to treat eczema.	0.369	Ayyanar and Ignacimuthu (2009), Bhat et al. (2012), Bhat et al. (2014), Ghorband and Biradar (2011), Kingston et al. (2009), Kumar et al. (2012), Muthu et al. (2006), Patil et al. (2009), Sharma et al. (2013a), Shiddamallayya et al. (2010), Silja et al. (2008), Upadhyay et al. (2010)	Native to the Western Ghats, India but found all over India. Also cultivated in parks; W & C (12)	The oil extracted from seeds is externally applied on itching and skin allergy.
<i>Senna tora</i> (L.) Roxb. [* <i>Cassia tora</i> L.] (GUH-JS 20146)	<i>Kasonji</i> (H)	Leaves are soaked in water, made into paste and applied topically on boils.	0.336	Adhikari et al. (2010), Gaur (1999), Gaur et al. (2011), Jyothi et al. (2010), Kumar et al. (2012), Sharma et al. (2010), Shiddamallayya et al. (2010), Singh et al. (2002), Sinha et al. (2013), Sivaranjani and Ramakrishnan (2012), Bhat et al. (2014)	Throughout India as a weed; W (11)	The seed powder is mixed with flour of <i>Triticum aestivum</i> (wheat), made into paste and applied externally in the treatment of boils.
Linaceae						
<i>Linum usitatissimum</i> L. (GUH-JS 20120)	<i>Alsi</i> (H)	The paste of whole plant is applied topically on wounds.	0.025	Adhikari et al. (2010), Sharma et al. (2013a)	Cultivated throughout India; C (2)	The paste of seeds is applied externally on boils.
Loranthaceae						
<i>Dendrophthoe falcata</i> (L.f.) Ettingsh. (GUH-JS 20294)	<i>Banda</i> (H)	The juice of bark is applied topically on wounds.	0.025	Ayyanar and Ignacimuthu (2009), Mairh et al. (2010), Sharma et al. (2013a), Singh et al. (2002)	Throughout India; W (4)	The roots are crushed and its juice is applied on skin allergy.
Lythraceae						
<i>Lawsonia inermis</i> L. (GUH-JS 18843)	<i>Mehndi</i> (S)	The leaf paste is applied on cracked heels in rainy season due to mud infection	0.311	Gaur (1999), Jain et al. (2005), Kingston et al. (2009), Kumar and Vidyasagar (2008), Kumar et al. (2012), Muthu et al. (2006), A.K. Sharma et al. (2013), Silja et al. (2008), M. Yadav et al. (2012)	Native to Arabia and Persia; Cultivated in northern and central India; C (9)	–

Table 1 (continued)

Plant species (family) voucher specimen number	Vernacular name (Habit)	Mode of preparation and dose	UV	Earlier ethnomedicinal studies in India citing use of plant for treatment of any skin disease	Distribution ^a ; Habitat (TC)	Uses by Gujjar Community of sub-Himalayan tract (Sharma et al., 2013a) ^b
Malvaceae						
<i>Sida cordata</i> (Burm. F.) Borss. Waalk. (GUH-JS 19753)	Bareyara (H)	The paste of leaves is topically applied on boils.	0.238	Adhikari et al. (2010), Katewa et al. (2004), Kumar and Sankar (2003), Purkayastha et al. (2005), Reddy et al. (2010), Senthilkumar et al. (2006a), J. Sharma et al. (2011)	Throughout hotter parts of India; W (7)	–
<i>Sida rhombifolia</i> L. (GUH-JS 20320)	Bariyara (H)	The root paste is topically applied on boils or abscess. The leaves are crushed and applied on wounds.	0.352	Adhikari et al. (2010), Gaur (1999), Singh and Maheswari (1994)	Throughout India, in moist places; W (3)	–
Martyniaceae						
<i>Martynia annua</i> L. (GUH-JS 20105)	Bicchua (S)	The oil of seeds is used topically in treatment of eczema.	0.303	Chakraborty and Bhattacharjee (2006), Adhikari et al. (2010), Jain et al. (2005)	Native of Mexico, found throughout India; W (3)	–
Meliaceae						
<i>Azadirachta indica</i> A. Juss. (GUH-JS 18826)	Nimba (Tr)	The stem bark is burnt and ash is applied topically on boils. Decoction of leaves is used to bath for the treatment of body infection. Its decoction is also taken orally for the treatment of the same. The seed oil is used externally to kill lice and to treat dandruff.	0.615	Bhat et al. (2012), Bhat et al. (2014), Choudhary et al. (2011), Dahare et al. (2010), Gaur (1999), Hiremath et al. (2010), Johnsy et al. (2012), Jyothi et al. (2010), Kingston et al. (2009), Kumar and Vidyasagar (2008), Kumar et al. (2012), A.K. Sharma et al. (2013), Shiddamallayya et al. (2010), Singh and Maheswari (1994), Singh et al. (2002), Sinha et al. (2013), Subramanian et al. (2011), Upadhyay et al. (2010), M. Yadav et al. (2012)	Native to Burma, found all over India; W (19)	–
Menispermaceae						
<i>Cissampelos pareira</i> L. (GUH-JS 18827)	Madrachi (H)	The juice of leaves is applied topically to cure itching. The roots are crushed made into paste and applied topically on leucoderma. The infusion of leaves (2 ml) is also taken internally, twice a day to heal wounds.	0.115	Ayyanar and Ignacimuthu (2009), Kuvar and Bapat (2010), Pradhan and Badola (2008), Sharma et al. (2013a)	The tropical and sub-tropical parts of India; W (4)	The paste of leaves is externally applied on boils.
Moraceae						
<i>Ficus benghalensis</i> L. (GUH-JS 18878)	Bargad (Tr)	The latex is applied topically on boils. The paste of leaves is mixed with coconut oil and applied topically on burns.	0.262	Ayyanar and Ignacimuthu (2009), Gaur and Sharma (2011), Johnsy et al. (2012), Katewa et al. (2004), J. Sharma et al. (2011)	Sub-Himalayan tract and Peninsular India. Planted along roadsides, and in gardens; W & C (5)	–
<i>Ficus racemosa</i> L. (GUH-JS 20127)	Gillor (Tr)	The latex is applied topically on boils	0.172	Ayyanar and Ignacimuthu (2009), Bhat et al. (2012), Negi et al. (2002), Punjani (2002), Sen and Behara (2003), Sharma et al. (2013a), Shiddamallayya et al. (2010), Shiddamallayya et al. (2010), Silja et al. (2008), Upadhyay et al. (2010)	Throughout India. Grows wild in forests and hills; W (10)	The paste of leaves is used in healing of wounds. The decoction (1–2 teaspoonfuls) of fruits is taken internally once a day for 2 weeks, to cure leprosy and its paste is applied externally for the same.
<i>Ficus religiosa</i> L. (GUH-JS 20366)	Peepal (Tr)	Bark paste is applied topically on boils	0.246	Ayyanar and Ignacimuthu (2009), Gaur (1999), Johnsy et al. (2012), Bhat et al. (2014)	Sub-Himalayan tracts, West Bengal, Central and South India; planted throughout India as an avenue tree; W & C (4)	–
Nyctaginaceae						
<i>Boerhavia diffusa</i> L. (GUH-JS 18845)	Punarnava (H)	The whole plant is pounded and its paste is applied topically on boils.	0.213	Chopda and Mahajan (2009)	Throughout India as a weed; W (1)	–

<i>Mirabilis jalapa</i> L. (GUH-JS 20343)	<i>Gulbans</i> (H)	The leaf juice is applied topically on boils or abscess.	0.057	Bhat et al. (2012) , Hiremath et al. (2010) , Kamble et al. (2010) , Sharma et al. (2013a) , Tomar (2009)	North-West Himalayas, Bengal and Manipur; W (5)	The leaf juice is applied externally on boils.
Orchidaceae						
<i>Vanda tessellata</i> (Roxb.) Hook. ex G.Don. (GUH-JS 20349)	<i>Harjori</i> (H)	Root paste is applied topically on boils or carbuncle.	0.016	J. Sharma et al. (2011) , Sharma et al. (2013a)	Sub-Himalayan tracts to 1500 m asl, Assam; W (2)	The root paste is applied externally on boils.
Papaveraceae						
<i>Argemone mexicana</i> L. (GUH-JS 18838)	<i>Pili kantiya</i> (H)	The paste of seeds is applied topically on itching. The seed oil is applied topically to treat ringworm. The yellow latex of the leaves is applied topically to cure chilblains. The yellow latex is applied between toes while working on paddy fields.	0.336	Chopda and Mahajan (2009) , Choudhary et al. (2011) , Dahare et al. (2010) , Johnsy et al. (2012) , Kingston et al. (2009) , Kumar and Vidyasagar (2008) , Kumar et al. (2012) , Kumari and Singh (2009) , Reddy et al. (2010) , Sharma et al. (2013a) , Sivaranjani and Ramakrishnan (2012)	Native to America, naturalized throughout India; W (11)	The latex is applied externally in cracks on feet. The latex along with cow milk is placed in copper pot for 3 days and applied externally in leucoderma. The yellow latex is applied between toes while working on paddy fields.
Pedaliaceae						
<i>Sesamum indicum</i> L. [* <i>Sesamum orientale</i> L.] (GUH-JS 20143)	<i>Til</i> (H)	The oil of seeds is applied topically in leprosy. The paste is applied on boils.	0.090	A.K. Sharma et al. (2013)	Up to 1500 m asl in Himalaya and in Uttar Pradesh, Madhya Pradesh, Rajasthan, Orissa, Gujarat, Andhra Pradesh, Tamil Nadu and Maharashtra; W & C (1)	–
Phyllanthaceae						
<i>Phyllanthus amarus</i> Schumach. and Thonn. (GUH-JS 19748)	<i>Jarmala</i> (H)	The juice of roots is applied topically on cuts and wounds.	0.172	Gaur (1999) , Sharma et al. (2013a) , Silja et al. (2008) , Upadhyay et al. (2010)	Throughout the hotter parts of India, particularly in cultivated land, up to 1000 m asl; W (4)	The crushed leaves are made into paste and applied externally on skin allergy.
Plantaginaceae						
<i>Scoparia dulcis</i> L. (GUH-JS 20229)	<i>Bichu</i> (H)	Paste of leaves is topically applied on wounds.	0.098	Ayyanar and Ignacimuthu (2009)	Native to tropical America. Introduced in India, commonly found in many parts of India; W (1)	–
Poaceae						
<i>Bambusa bambos</i> (L.) Voss [* <i>Bambusa arundinacea</i> Willd.] (GUH-JS 20358)	<i>Bans</i> (S)	The paste of stem is applied topically on wounds.	0.033	Chopda and Mahajan (2009) , Johnsy et al. (2012)	Wild throughout India, especially in the hill forests; W (2)	–
<i>Chrysopogon zizanioides</i> (L.) Roberty [* <i>Vetiveria zizanioides</i> (L.) Nash] (GUH-JS 19729)	<i>Seenk</i> (S)	The paste of roots is topically applied on boils.	0.295	Gaur (1999) , J. Sharma et al. (2011)	A perennial grass cultivated almost throughout India; C (2)	–
<i>Cynodon dactylon</i> (L.) Pers. (GUH-JS 20293)	<i>Dhoob</i> (H)	The whole plant paste is applied topically on cuts and wounds to stop bleeding	0.328	Abbasi et al. (2010) , Bhandary and Chandrashekhara (2002) , Bhat et al. (2012) , Bhat et al. (2014) , Chopda and Mahajan (2009) , Gangwar et al. (2010) , Kingston et al. (2009) ,	Throughout India up to 3000 m asl; W (14)	The plant paste is applied on cuts and wounds to stop bleeding.

Table 1 (continued)

Plant species (family) voucher specimen number	Vernacular name (Habit)	Mode of preparation and dose	UV	Earlier ethnomedicinal studies in India citing use of plant for treatment of any skin disease	Distribution ^a ; Habitat (TC)	Uses by Gujjar Community of sub-Himalayan tract (Sharma et al., 2013a) ^b
<i>Desmostachya bipinnata</i> (L.) Stapf (GUH-JS 20303)	Dubh (S)	The roots are dried and its powder is sprinkled on wounds.	0.033	Negi et al. (2011), Pradhan and Badola (2008), Sharma et al. (2013a), Shiddamallayya et al. (2010), Sivaranjani and Ramakrishnan (2012), Subramanian et al. (2011), Upadhyay et al. (2010)	Throughout the plains of India in dry and hot areas and in sandy deserts; W (1)	–
<i>Eulaliopsis binata</i> (Retz.) C.E.Hubb. (GUH-JS 18808)	Baib (H)	The juice of leaves is applied topically on cuts and wounds.	0.115	Sharma et al. (2013a)	Northern India; W (1)	The plant juice is applied on cuts.
Polygonaceae						
<i>Persicaria barbata</i> (L.) H.Hara (GUH-JS 18806)	Bhidiya (H)	The paste of leaves is applied topically on boils. Juice or paste of leaves is applied on wounds to cure pain.	0.115	Gaur and Sharma (2011), Sharma et al. (2013a)	Throughout the warmer parts of India, ascending to 1400 m asl; W (2)	The juice of leaves is applied on cuts.
Ranunculaceae						
<i>Ranunculus sceleratus</i> L. (GUH-JS 18810)	Jaldhaniya (H)	The paste of leaves is applied topically on boils.	0.205	Gaur (1999), Rani et al. (2013)	The plains of northern India, and the warm valleys of the Himalayas from Kashmir to Assam; W (2)	–
Rhamnaceae						
<i>Ziziphus nummularia</i> (Burm. f.) Wight & Arn. (GUH-JS 20328)	Jharberiya (S)	The bark paste is applied on eczema. The seeds are burnt, ash is mixed with a little amount of honey and applied to avoid baldness (hair fall).	0.123	Katewa et al. (2004), Mairh et al. (2010), Singh et al. (2002)	Throughout warmer parts of India in wastelands; W (3)	–
Rubiaceae						
<i>Mitragyna parvifolia</i> (Roxb.) Korth. (GUH-JS 19793)	Kaim (Tr)	The paste of leaves is applied topically on cuts and wounds. The bark paste is applied topically on leucoderma.	0.107	Nil	All over India, up to 1200 m asl in the outer Himalaya; W (0)	–
Scrophulariaceae						
<i>Verbascum thapsus</i> L. (GUH-JS 18811)	Chudighass (H)	The paste of leaves is mixed with sarsoon oil and topically applied on boils.	0.139	Lal and Singh (2008), Sharma et al. (2013a)	Temperate Himalayas, Western Ghats and the Nilgiris; up to 2500; W (2)	The seed paste is used externally on leucoderma.
Solanaceae						
<i>Solanum incanum</i> L. (GUH-JS 20329)	Kachriya (S)	The roots are pounded and applied topically on leucoderma.	0.049	Gaur (1999)	Sub Himalayan tracts and South India; W (1)	–
<i>Datura stramonium</i> L. (GUH-JS 20364)	Kala daturu (H)	The leaves are grounded with little quantity of black pepper powder and applied topically on wounds.	0.287	Kumar and Vidyasagar (2008), Kumari and Singh (2009), Sivaranjani and Ramakrishnan (2012)	From Kashmir to Sikkim hilly districts of Central and South India, up to 2700 m asl; W (3)	–
<i>Nicotiana plumbaginifolia</i> Viv. (GUH-JS 19713)	Bhangraleeg (H)	The paste of roots is applied topically on leucoderma. The leaf paste is applied topically on wounds. Juice of leaves is applied on itches.	0.148	Dangwal et al. (2010), Sharma et al. (2013a)	Native of Mexico and West Indies, widely distributed in India, common along road sides; W (2)	The leaf paste is applied externally on wounds and itching.

Typhaceae

<i>Typha domingensis</i> Pers. [<i>*Typha angustata</i> Bory & Chaub.] (GUH-JS 20308)	<i>Pater</i> (S)	The fruit paste is topically applied on wounds.	0.008	Katewa et al. (2004)	Throughout India, up to 1730 m – asl; W (1)	
Ulmaceae						
<i>Holoptelea integrifolia</i> Planch. (GUH-JS 20219)	<i>Chilbil</i> (Tr)	The paste of stem bark is applied topically on ringworm.	0.074	Gaur and Sharma (2011) , Meena and Yadav (2011) , Sharma et al. (2013a) , Upadhyay et al. (2010)	Throughout greater parts of India, also grown in gardens; W & C (4)	The tender leaves paste is applied externally on eczema. The paste of inner bark is applied on blisters.
Verbenaceae						
<i>Lantana camara</i> L. (GUH-JS 20223)	<i>Lalten</i> (S)	Leaf juice is applied topically on cuts and wounds.	0.352	Bhat et al. (2012) , Gaur (1999) , Jyothi et al. (2010) , Kumar and Vidyasagar (2008) , Mamatha et al. (2006) , Pradhan and Badola (2008) , Rajakumar and Shivanna (2009) , Sharma et al. (2013a) , Sinha et al. (2013) , Subramanian et al. (2011) , Upadhyay et al. (2010) , Wadankar et al. (2011)	Native to tropical America. Naturalized and occurs throughout India; W (12)	The paste of leaves is applied on ringworm.

Abbreviations: C=cultivated; FL=fidelity level; H=herb; S=shrub; TC=number of times species cited in the reviewed literature to treat any skin disease in India; Tr=Tree; UV=use value; W & C=both wild as well as cultivated; W=Wild.

^a Sources ([Gaur, 1999](#); [Khare, 2007](#)).

^b Uses in bold denotes exactly same use of same plant part as recorded in the present study.

Moraceae (3 taxa), Solanaceae (3 taxa), Amaranthaceae (2 taxa), Convolvulaceae (2 taxa), Malvaceae (2 taxa) and Nyctaginaceae (2 taxa). In some earlier studies it was estimated that approximately one third of all the traditional medicines are for treatment of wounds or skin disorders, compared to only 1–3% of modern drugs (Mantle et al., 2001). Literature review revealed that in most of the earlier studies very few plants have been reported for treatment of hair fall, dandruff and toes infection, whereas large numbers of plants have been reported for treatment of wounds, cuts and boils. In the present study maximum number of plants (38 spp.) were used to treat wounds followed by boils (32 spp.), cuts (18 spp.), leprosy (11 spp.), eczema (10 spp.), itching (7 spp.), ringworm (5 spp.), burns (4 spp.), leucoderma (4 spp.), cracked heels (2 spp.), dandruff (3 spp.), body infection (2 spp.), chilblains (2 spp.), hair fall (2 spp.) and toes infection (2 spp.) (Fig. 2).

As shown in Fig. 3, among different plant parts, leaves (48.28%) were the most commonly used for preparation of herbal ointment followed by seeds (17.24%), roots (14.94%), whole plant (13.79%), stem (5.75%), flower (4.60%) and fruit (2.30%). It has been noticed that leaves were frequently used as a remedy to cure skin ailments as compared to any other plant part due to their easy availability in case of cuts and wounds to stop bleeding and fast relief. Latexes of 8 plant species were directly applied topically on the affected body part viz., latexes of *Cryptolepis dubia* (Burm.f.) M.R.Almeida, *Ficus benghalensis* L., *Ficus racemosa* L. and *Premna mollissima* Roth for boils; latexes of *Cryptolepis dubia* (Burm.f.) M.R.Almeida, *Euphorbia hirta* L., *Vallis solanacea* (Roth) Kuntze and *Wrightia arborea* (Dennst.) Mabb. for wounds; yellow latex of *Argemone mexicana* L. for chilblains; latex of *Cryptolepis dubia* (Burm.f.) M.R.Almeida for sores; latex of *Euphorbia hirta* L. for eczema; latex of *Wrightia arborea* (Dennst.) Mabb. for cuts. Similarly, gums of three plant species were also used topically on the skin i.e. fresh gum of *Butea monosperma* (Lam.) Taub. as antiseptic on cuts and wounds; gum of *Semecarpus anacardium* L.f. to cure leprosy; and gum of *Shorea robusta* Gaertn. on cracked heels.

As shown in Fig. 4, most frequently used form of external administration of herbal medicine was paste (65.05%), which is prepared by pulverizing plant matter with water. Externally, a squeezed juice of the plant part was directly used in 18.45% of preparations, followed by use of latex (8.74%), seed oil (5.83%), powder (3.88%), gum (2.91%) and ash (1.94%). In most of the cases herbal combinations were applied topically on affected parts to heal skin disorders but internally decoction (6 preparations) was most commonly used mode of administration followed by infusion and powder form. Seeds were usually applied in the form of paste over the skin, but in some cases seed oil was used viz. seed oil of *Argemone mexicana* L., *Martynia annua* L., *Pongamia pinnata* (L.) Pierre, *Ricinus communis* L. and *Sesamum indicum* L. Addition of oil of other plant species i.e., mustard (*Brassica rapa* L.) and coconut (*Cocos nucifera* L.) with the recorded plants in some medicinal formulations was also recorded. The oil of mustard seeds was mixed with seed paste of *Crotalaria juncea* L. and applied on the wound. In some cases leaves of *Cannabis sativa* L., *Ricinus communis* L. and *Ipomoea carnea* Jacq. were separately fried in mustard oil, made into paste and applied on affected body part. Leaf pastes of *Anisomeles indica* (L.) Kuntze and *Ficus benghalensis* L. were mixed with coconut (*Cocos nucifera* L.) oil and used topically for healing. Besides that black pepper (*Piper nigrum* L.) and common salt (NaCl or Sodium Chloride) in minute quantities were also added in some formulations, for example the leaf paste of *Acalypha indica* L. was mixed with a little quantity of salt and applied topically on eczema. The possible reason for use of common salt is that it enters the tissue and in effect binds the water, inhibiting the activity of bacteria and fungi, which could in turn help in rapid healing of the skin. Similarly, very little quantity of black pepper powder was grounded with leaves of *Datura stramonium* L. and

applied on wounds. This seems to be very scientific as Piperine a major active component of black and long peppers, has been reported to enhance drug bioavailability by inhibiting certain enzyme metabolism (Atal et al., 1985). Addition of salt and black pepper was believed to have synergistic effect in the treatment of skin ailment.

Available ethnomedicinal literature was reviewed to study whether plants reported in the present study are used by any other indigenous community in India to treat any skin ailment (Table 1). *Mitragyna parvifolia* (Roxb.) Korth. was not reported in any of the earlier studies to treat any skin disease. Some little known ethnomedicinal plant species used to treat skin diseases reported in the present study, which were cited in only one study were *Boerhavia diffusa* L., *Caesulia axillaris* Roxb., *Desmostachya bipinnata* (L.) Stapf, *Eulaliopsis binata* (Retz.) C.E.Hubb., *Euphorbia thymifolia* L., *Hyptis suaveolens* (L.) Poit., *Oroxylum indicum* (L.) Kurz, *Scoparia dulcis* L., *Sesamum indicum* L., *Solanum incanum* L. and *Typha domingensis* Pers. (Table 1). According to literature survey the most commonly used plant species to treat skin diseases in India is *Achyranthes aspera* L. (23), followed by *Azadirachta indica* A. Juss. (19 citations), *Cassia fistula* L. (17 citations), *Euphorbia hirta* L. (16 citations), *Cynodon dactylon* (L.) Pers. (14 citations), *Annona squamosa* L. (12 citations), *Pongamia pinnata* (L.) Pierre (12 citations), *Lantana camara* L. (12 citations), *Tridax procumbens* (L.) L. (11 citations), *Senna tora* (L.) Roxb. (11 citations), *Argemone mexicana* L. (10 citations), *Ficus racemosa* L. (10 citations), *Ricinus communis* L. (9 citations) and *Lawsonia inermis* L. (9 citations).

Out of the recorded 90 plant species, 8 plant species are cultivated (*Allium cepa* L., *Annona squamosa* L., *Brassica juncea* (L.) Czern., *Cassia fistula* L., *Chrysopogon zizanioides* (L.) Roberty, *Lawsonia inermis* L., *Linum usitatissimum* L. and *Momordica dioica* Roxb. ex Willd.), whereas 12 plant species (*Acorus calamus* L., *Bauhinia variegata* L., *Cannabis sativa* L., *Ehretia laevis* Roxb., *Ficus benghalensis* L., *Ficus religiosa* L., *Holoptelea integrifolia* Planch., *Pongamia pinnata* (L.) Pierre, *Ricinus communis* L., *Sesamum indicum* L., *Terminalia arjuna* (Roxb. ex DC.) Wight & Arn. and *Vallis solanacea* (Roth) Kuntze.) are used from both wild as well as cultivated sources. Remaining 70 plant species are available in wild.

All the reported plant species are widely distributed through out India (Table 1). Twelve plant species are non-native or introduced to India, but they are now naturalized throughout the country including the study area. These plant species are *Amaranthus spinosus* L., *Annona squamosa* L., *Argemone mexicana* L., *Azadirachta indica* A. Juss., *Hyptis suaveolens* (L.) Poit., *Ipomoea carnea* Jacq., *Lantana camara* L., *Lawsonia inermis* L., *Martynia annua* L., *Nicotiana glauca* Viv., *Ricinus communis* L. and *Scoparia dulcis* L. According to UV value most preferred plant species to treat skin diseases by Tharu community is *Ricinus communis* L. (0.631) followed by *Tridax procumbens* (L.) L. (0.623), *Azadirachta indica* A. Juss. (0.615), *Ageratum conyzoides* (L.) L. (0.541) and *Allium cepa* L. (0.500). The F_{ic} values for different categories of skin diseases and FL of all the reported plant species are given in Table 2. Values of F_{ic} ranged from 0.880 for leucoderma to 0.976 for cracked heels. High F_{ic} values in the present study suggest that there is ample exchange of information between informants and there is well defined selection criterion in the community for use of these plants (Gazzaneo et al., 2005).

The usage pattern of medicinal plant resources by any particular community is usually unique to that particular community and is part of its cultural traditional knowledge, which is passed on from one generation to another. Most of the studies on medicinal plants focus on the role of these plants within one culture or one ethnic group and little emphasis have been given to the comparison of medicinal plant species in various cultures (Heinrich et al., 1998). In past few decades some researchers have drawn attention of research community to the lack of information on the relative

importance of a medicinal plant within a culture and the need for comparing the use of plants interculturally (Heinrich et al., 1992, 1998; Aguilar et al., 1994; Etkin, 1994; Moreman, 1996). Earlier some workers have studied intercultural importance of medicinal plants between various cultures in different parts of the world (Heinrich et al., 1998; Leonti et al., 2006, 2009, 2010; Lardos and Heinrich, 2013; Gairola et al., 2014). This intercultural comparison approach is practical since both consensus and variations can be addressed by this approach (Heinrich et al., 1998). Therefore to see how widespread presently recorded plants are used to treat skin diseases in the sub-Himalayan region of Uttarakhand, we compared results of the present study with results of one of our earlier published study on Gujjar community inhabiting forests with similar vegetation in nearby districts of Dehradun, Haridwar and Pauri in sub-Himalayan region of Uttarakhand (Sharma et al., 2013a). In that study we had recorded 109 medicinal plants used to treat skin diseases by Gujjar community (for details see Sharma et al. (2013a)). Gujjar community is one of the important migratory communities of the Himalaya. When we compared results of these two studies we observed that although both the communities inhabit forests with similar vegetation but medicinal plant usage pattern of both the communities is very much different from each other and unique to them. A total of 154 plant species were used by both the communities combined to treat skin diseases out of which use of 46 plant species was common between these communities. But out of these 46 plant species only 16 plant species were used in exactly same way to treat same type of skin disease (Table 1 and Fig. 5). Although rest 30 plant species were also used by both the communities, but plant part used, skin disease treated and methods of preparation were different. For comparison, information about uses of these 46 plants by Gujjar community is given in Table 1. Since these communities inhabit nearby forest regions from hundreds of years, some exchange of information regarding usage of medicinal plants between these communities is inevitable, which could be one of the possible reasons for consensus on these 16 plants with exactly same usage.

Availability of herbal resources is sometimes suggested as one of the main criterion for selection of plants by the communities (Moreman, 1996; Heinrich et al., 1998). But in the present case both the communities inhabit forests with similar vegetation and availability of herbal resources are almost identical to them. Therefore, it seems that selection of medicinal plants by these communities is not random or based solely on the availability of the resources and there are some specific criteria, which are used by each community for selecting these plants. These criteria could well be related to cultural, social and religious backgrounds of these communities and are responsible for variation in medicinal plant usage pattern. However, in future studies it would be interesting to assess differences in specific criteria used by these




Fig. 3. Percentage of plant parts used in herbal preparations as remedy of skin diseases.




Fig. 4. Percentage of different modes of preparation of herbal remedies used to treat skin diseases.

communities to select medicinal plants, and exchange of knowledge on uses of medicinal plants between these and other communities inhabiting sub-Himalayan region of Uttarakhand.

3.2. Pharmacological, toxicological and clinical considerations

Some categories of skin diseases are caused by bacteria and fungi, while others are caused by environmental conditions. For example, boils are caused by bacteria or fungi present on the skin surface, leprosy is a chronic infection caused by *Mycobacterium leprae* and *Mycobacterium lepromatosis*, eczema is a bacterial infection caused by the bacteria *Staphylococcus aureus*, ringworm is a fungal infection caused by dermatophytes of genera *Trichophyton* and *Microsporum*. Whereas, leucoderma, also known as vitiligo is a condition in which loss of skin pigments take place. Causes of vitiligo are unknown but according to some studies it is autoimmune condition in which immune system of body destroy its own cells and tissues. Wounds and cuts are of external physical origin, chilblains are tissue injury develop as an abnormal response to cold, Itching also known as pruritus is an irritating sensation on skin, which may be caused by allergy and variety of micro-organisms. Tharu community use medicinal plants to treat almost all the category of skin diseases. Details of earlier studies showing wound healing, toxicological, antioxidant and tyrosinase inhibitory properties of the plants recorded in the present study have been provided in Table 3. Whereas, summary of the earlier studies showing antiinflammatory and antimicrobial activities, and active compounds in the plants reported in the present study have been given in Table 4. Toxicological, pharmacological, clinical and other information on the recorded plants related to the skin diseases have been discussed below.

Acalypha indica L. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in earlier the studies (Tables 3 and 4). Its ethanol extract showed significant antioxidant activity in superoxide and hydroxyl radical scavenging activity and effect on lipid peroxidation tests (Joy and Mathew, 2010), whereas aqueous extract showed DPPH radical scavenging activity with IC₅₀ value of 527.86 µg/ml (Guha et al., 2011). In toxicity test ethanolic extract of whole plant has been found to be non-toxic to male Wistar albino rats even at dose of 2000 mg/kg bw (Sathya et al., 2012). In the present study it is reported to be used topically on wounds, which corroborates with the aforesaid scientific studies done on this plant. However human clinical trial related to any skin disease has not been conducted on this plant.

Achyranthes aspera L. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). The methanol extract, alkaloid, non-alkaloid and saponin fractions of *Achyranthes aspera* L. leaves exhibited

Table 2

Consensus factor values for different illness categories and Fidelity level of the plant species used to treat various skin diseases by Tharu community.

Illness categories (F_{ic})	Plant species (FL)
Body infection (0.933)	<i>Acalypha indica</i> (14.29), <i>Azadirachta indica</i> (16.00)
Boils (0.944)	<i>Achyranthes aspera</i> (48.78), <i>Albizia lebeck</i> (100.00), <i>Amaranthus spinosus</i> (63.16), <i>Annona squamosa</i> (31.25), <i>Azadirachta indica</i> (41.33), <i>Boerhavia diffusa</i> (100.00), <i>Brassica juncea</i> (100.00), <i>Calotropis procera</i> (100.00), <i>Chrysopogon zizanioides</i> (100.00), <i>Cryptolepis dubia</i> (37.50), <i>Desmodium gangeticum</i> (100.00), <i>Eclipta prostrata</i> (21.43), <i>Ficus benghalensis</i> (40.63), <i>Ficus racemosa</i> (100.00), <i>Ficus religiosa</i> (100.00), <i>Holarrhena pubescens</i> (20.59), <i>Litsea glutinosa</i> (100.00), <i>Milletia extensa</i> (100.00), <i>Mirabilis jalapa</i> (100.00), <i>Momordica dioica</i> (100.00), <i>Mucuna pruriens</i> (100.00), <i>Persicaria barbata</i> (28.57), <i>Premna mollissima</i> (42.11), <i>Ranunculus sceleratus</i> (100.00), <i>Rauvolfia serpentina</i> (100.00), <i>Senna tora</i> (100.00), <i>Sesamum indicum</i> (45.45), <i>Sida cordata</i> (100.00), <i>Sida rhombifolia</i> (32.56), <i>Tridax procumbens</i> (36.84), <i>Vanda tessellata</i> (100.00), <i>Verbascum thapsus</i> (100.00)
Burn (0.953)	<i>Buchanania cochinchinensis</i> (63.64), <i>Cannabis sativa</i> (100.00), <i>Ficus benghalensis</i> (59.38), <i>Oroxylum indicum</i> (75.00)
Chilblain (0.900)	<i>Argemone mexicana</i> (14.63), <i>Callicarpa macrophylla</i> (100.00)
Cracked heels (0.976)	<i>Lawsonia inermis</i> (100.00), <i>Shorea robusta</i> (100.00)
Cuts (0.949)	<i>Ageratum conyzoides</i> (84.85), <i>Buchanania cochinchinensis</i> (36.36), <i>Butea monosperma</i> (46.67), <i>Caesulia axillaris</i> (77.27), <i>Clerodendrum infortunatum</i> (100.00), <i>Curculigo orchioideis</i> (50.00), <i>Cynodon dactylon</i> (65.00), <i>Eclipta prostrata</i> (47.62), <i>Eulaliopsis binata</i> (78.57), <i>Hyptis suaveolens</i> (66.67), <i>Ipomoea carnea</i> (73.91), <i>Lannea coromandelica</i> (79.17), <i>Lantana camara</i> (60.47), <i>Mallotus philippensis</i> (50.00), <i>Mitragyna parvifolia</i> (23.08), <i>Phyllanthus amarus</i> (71.43), <i>Tridax procumbens</i> (63.16), <i>Wrightia arborea</i> (77.78)
Dandruff (0.966)	<i>Annona squamosa</i> (68.75), <i>Azadirachta indica</i> (42.67), <i>Cuscuta reflexa</i> (69.57)
Eczema (0.959)	<i>Acalypha indica</i> (64.29), <i>Acorus calamus</i> (55.56), <i>Amaranthus spinosus</i> (36.84), <i>Cuscuta reflexa</i> (30.43), <i>Euphorbia hirta</i> (26.32), <i>Martynia annua</i> (100.00), <i>Pongamia pinnata</i> (100.00), <i>Ricinus communis</i> (62.34), <i>Vitex negundo</i> (60.34), <i>Ziziphus nummularia</i> (46.67)
Hair fall (0.933)	<i>Cleome viscosa</i> (61.54), <i>Ziziphus nummularia</i> (53.33)
Itching (0.885)	<i>Argemone mexicana</i> (24.39), <i>Cassia fistula</i> (58.82), <i>Cissampelos pareira</i> (50.00), <i>Curculigo orchioideis</i> (50.00), <i>Euphorbia thymifolia</i> (100.00), <i>Hyptis suaveolens</i> (16.67), <i>Nicotiana plumbaginifolia</i> (33.33)
Leprosy (0.916)	<i>Ageratum conyzoides</i> (15.15), <i>Bauhinia variegata</i> (100.00), <i>Butea monosperma</i> (40.00), <i>Cassia fistula</i> (41.18), <i>Celastrus paniculatus</i> (100.00), <i>Dalbergia sissoo</i> (100.00), <i>Holarrhena pubescens</i> (79.41), <i>Premna mollissima</i> (57.89), <i>Semecarpus anacardium</i> (100.00), <i>Sesamum indicum</i> (54.55), <i>Terminalia arjuna</i> (100.00)
Leucoderma (0.880)	<i>Cissampelos pareira</i> (35.71), <i>Mitragyna parvifolia</i> (53.85), <i>Nicotiana plumbaginifolia</i> (44.44), <i>Solanum incanum</i> (100.00)
Ringworm (0.961)	<i>Allium cepa</i> (100.00), <i>Argemone mexicana</i> (46.34), <i>Euphorbia hirta</i> (31.58), <i>Holoptelea integrifolia</i> (100.00), <i>Leonotis neopetifolia</i> (100.00)
Toes infection (0.962)	<i>Achyranthes aspera</i> (51.22), <i>Argemone mexicana</i> (14.63)
Wound (0.888)	<i>Acalypha indica</i> (21.43), <i>Acorus calamus</i> (44.44), <i>Anisomeles indica</i> (100.00), <i>Bambusa bambos</i> (100.00), <i>Butea monosperma</i> (13.33), <i>Caesulia axillaris</i> (22.73), <i>Cissampelos pareira</i> (14.29), <i>Cleome viscosa</i> (38.46), <i>Commelina benghalensis</i> (100.00), <i>Crotalaria juncea</i> (100.00), <i>Cryptolepis dubia</i> (62.50), <i>Cynodon dactylon</i> (35.00), <i>Datura stramonium</i> (100.00), <i>Dendrophthoe falcata</i> (100.00), <i>Desmostachya bipinnata</i> (100.00), <i>Dicliptera paniculata</i> (100.00), <i>Eclipta prostrata</i> (30.95), <i>Ehretia laevis</i> (100.00), <i>Eulaliopsis binata</i> (21.43), <i>Euphorbia hirta</i> (42.11), <i>Hyptis suaveolens</i> (16.67), <i>Ipomoea carnea</i> (26.09), <i>Lannea coromandelica</i> (20.83), <i>Lantana camara</i> (39.53), <i>Linum usitatissimum</i> (100.00), <i>Mallotus philippensis</i> (50.00), <i>Mitragyna parvifolia</i> (23.08), <i>Nicotiana plumbaginifolia</i> (22.22), <i>Oroxylum indicum</i> (25.00), <i>Persicaria barbata</i> (71.43), <i>Phyllanthus amarus</i> (28.57), <i>Ricinus communis</i> (37.66), <i>Scoparia dulcis</i> (100.00), <i>Sida rhombifolia</i> (67.44), <i>Typha domingensis</i> (100.00), <i>Vallisneria spiralis</i> (100.00), <i>Vitex negundo</i> (39.66), <i>Wrightia arborea</i> (22.22)

Abbreviations: F_{ic} =Consensus factor, FL=Fidelity level.

significant inhibitory effects (concentration 100 µg) on the Epstein–Barr virus early antigen activation induced by the tumor promoter 12-O-tetradecanoylphorbol-13-acetate in Raji cells (Chakraborty et al., 2002). The leaf extract and the non-alkaloid fraction of this plant are valuable antitumor promoters in carcinogenesis (Chakraborty et al., 2002). *Achyranthes aspera* L. also has trypsin inhibitory properties (Geetha and Davuluri, 2013). The ethanol and aqueous extracts of leaves of *Achyranthes aspera* L. have shown wound healing activity in excision and incision wound models (Edwin et al., 2008). The ethanol and aqueous extracts of leaves of *Achyranthes aspera* L. showed antioxidant properties in DPPH radical scavenging and superoxide scavenging activities (Edwin et al., 2008). Antioxidant activities were also reported in leaves (Tahiliani and Kar, 2000) and seeds (Malarvili and Gomathi, 2009). The alcoholic extracts of leaves and seeds of *Achyranthes aspera* L. showed anti-inflammatory activity at the oral dose of 250 mg/kg body weight in carrageenan-induced paw edema model and formalin model in Wistar albino rats of either sex (weighing 200–250 g) (Mehta et al., 2009). Ethanolic extracts of *Achyranthes aspera* L. significantly inhibited paw edema induced by carrageenan and Freund's complete adjuvant and prevent accumulation of inflammatory cells in carrageenan-induced peritonitis at doses of 100–200 mg/kg bw (Gokhale et al., 2002). In acute and sub acute oral toxicity tests, this plant was found to be non-toxic to swiss albino mice (Table 3). Clinical trial of the plant on patients with shoth (general anasarca) (Shankar et al., 1980) and Obesity (Mangal and Sharma, 2009) showed promising results. However human clinical trial related to any skin disease has not been conducted on this plant.

Acorus calamus L. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Several studies have recognized anti-inflammatory potential of *Acorus calamus* L. Rhizome extract showed anti-inflammatory property in acute, chronic and immunologic models of inflammation (Varde et al., 1988; Vohra et al., 1989), whereas leaf extract showed anti-inflammatory activity in study on human keratinocyte HaCaT cells (Kim et al., 2009). Antiinflammatory effects of *Acorus calamus* L. could be mediated by suppression of NF-κB and interferon regulatory factor 3 (IRF3) (Kim et al., 2009). Mild but acceptable toxicity of rhizome extract of this plant was observed at high doses by Muthuraman and Singh (2012). β-asarone, a compound from this plant is potentially toxic and carcinogenic (Taylor et al., 1967; Keller and Stahl, 1983), whereas α-asarone has hepatocarcinogenic and mutagenic activity in mice (Chamarro et al., 1998). β-asarone has an oral LD₅₀ of 1010 mg/kg bw in rats and i.p. LD₅₀ of 184 mg/kg bw in mice (JECFA, 1981). The free radical scavenging activity of *Acorus calamus* L. has been found to be useful to overcome excess production of reactive oxygen species generated due to continuous exposure to loud noise and β-Asarone is believed to be involved in reducing the stress (Manikandan and Devi, 2005). Clinical studies on extracts of *Acorus calamus* L. on the management of depression (Tripathi and Singh, 1995), generalized anxiety disorder (Bhattacharyya et al., 2011), ischemic heart disease (Mamgain and Singh, 1994) and bronchial asthma (Rajasekharan and Srivastava, 1977) in humans have shown promising results but clinical trials related to the skin diseases are still lacking. However, *Acorus calamus* L. along with 21 other plants is part of a polyherbal Ayurvedic formulation of The Himalaya Drug Company known as “Muscle & Joint Rub”, which is recommended for the topical management of acute and chronic musculoskeletal disorders. Phase III clinical trial of Muscle & Joint Rub showed that it is effective and safe in the management of muscle sprains, contusions and inflammatory musculoskeletal disorders (Rajanna and Kolhapure, 2005). Since the product studied was a combination of different plants, it is not clear what exact effect *Acorus calamus* L.




Fig. 5. Comparison between plants used to treat skin diseases by two important indigenous communities of sub-Himalayan region of Uttarakhand viz., Gujar and Tharu.

had on the studied disease. Additional clinical studies are required before a firm conclusion can be made.

Ageratum conyzoides (L.) L. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Leaf extract of this plant was not toxic even at high doses of 5000 mg/kg bw in acute oral toxicity test on Wistar rats (Diallo et al., 2010). Clinical trials of aqueous extract of the whole plant was done with patients with arthrosis and results showed analgesic effect in 66% patients and improvement in articulation mobility in 24% patients without side effects (Marques-Neto et al., 1988). *Albizia lebbbeck* (L.) Benth. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). This plant also has trypsin inhibitory properties (P. Sharma et al., 2012). Histamine plays major role in allergic diseases and its action is mediated mainly by H₁ receptor (H1R) (Nurul et al., 2011). The effect of the extract from the bark of *Albizia lebbbeck* (L.) Benth. (AL) on H1R and HDC gene expression using toluene-2,4-diisocyanate (TDI) sensitized allergy model rats and HeLa cells expressing endogenous H1R was studied by Nurul et al. (2011). Their study suggested that AL alleviated nasal symptoms by inhibiting histamine signaling in TDI-sensitized rats through suppression of H1R and HDC gene transcriptions; moreover suppression of Th2-cytokine signaling by AL also suggests that it could affect the histamine–cytokine network (Nurul et al., 2011). Methanolic leaf extract of this plant was non-toxic even at a dose of 2000 mg/kg bw in acute oral toxicity study on Wistar rats (Sivakumar et al., 2013). However human clinical trial related to any skin disease has not been conducted on *Ageratum conyzoides* (L.) L. and *Albizia lebbbeck* (L.) Benth.

Allium cepa L. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). This plant also has trypsin inhibitory properties (Deshimaru et al., 2003) as well as tyrosinase inhibitory properties (Arung et al., 2011). *Allium cepa* L. has many biologically active antifungal, antibacterial and antimicrobial compounds (Table 4) and is non-toxic orally even at high doses (Bhanot and Shri, 2010) (Table 3). Some skin care products made from extract of this plant are already available in the market. *In vivo* clinical study on thirty people with hypertrophic scars or keloids showed that topical applications of a gel containing *Allium cepa* L. extract, pentaglycan, and allantoin twice a day for 24 weeks is useful in reducing neoangiogenesis in hypertrophic scars and keloids, resulting in clinical improvement of skin lesions (Campanati et al., 2010).

Table 3

Earlier studies showing wound healing, toxicological, antioxidant and tyrosinase inhibitory properties of the plants recorded in the present study.

Plant species	Wound healing studies		Toxicological studies		Antioxidant activities	Tyrosinase inhibitory activities
	Plant part; Application; Extract; Animal (Weight)	Animal Model; Reference control; Effective treatment [Source for columns 2 & 3]	Plant part; Extract; Animal (Weight)	Animal model (Dose); Toxicity status [Source for columns 4 & 5]		
<i>Acalypha indica</i> L.	Whole plant; Topical application; 10% (w/v) alcoholic extract; Lewis Wistar albino rats of either sex (100–200 g)	EWM, IWM; NA; 10% (w/v) alcoholic extract (Reddy et al., 2002)	Whole plant; Ethanolic; Male Wistar albino rat (150 to 200 g)	AOTS (5, 50, 300, 500 and 2000 mg/kg bw), SATS (100,200,300,400, and 500 mg/kg/day bw); NT/NM (Sathya et al., 2012)	Aqueous (IC ₅₀ =527.86 µg/ml), methanol (IC ₅₀ =483.87 µg/ml), hexane (IC ₅₀ =805.27 µg/ml) and chloroform (IC ₅₀ =1688.00 µg/ml) extracts of whole plant showed DPPH radical scavenging activity (Guha et al., 2011).	NE
<i>Achyranthes aspera</i> L.	Leaves; Topical application; 2.5%, 5% and 10% (w/w) methanolic extract; Albino rats of either sex (200–250 g)	EWM, IWM; 1% SSD; 5% and 10% (w/w) methanolic extract (Fikru et al., 2012)	Leaves; Methanolic; Swiss mice (35 to 45 g)	AOTS (2, 4, 6 and 8 gm/kg bw); NT/NM (Sadashiv and Krishna, 2011)	Aqueous (IC ₅₀ =179.35 µg/ml), methanol (IC ₅₀ =291.92 µg/ml), hexane (IC ₅₀ =559.32 µg/ml) and chloroform (IC ₅₀ =652.60 µg/ml) extracts of whole plant showed DPPH radical scavenging activity (Guha et al., 2011).	NE
<i>Acorus calamus</i> L.	Leaves; Topical application; 40% and 20% (w/w) ethanolic extract; Rats of either sex (NA)	EWM, IWM; PIO; 20% (w/w) ethanolic extract (Jain et al., 2010)	Whole Plant; Methanolic; Swiss albino mice (20–25 g) Rhizome; Hydroalcoholic; Mice (either sex, 20–30 g) and Wistar rats (either sex, 200–230 g) Rhizome; Ethanolic; Female Wistar rats (150–200 g) and Wistar albino rats of either sex (100–190 g)	AOTS (150, 200, 250 mg/kg bw), SATS (25 and 50 mg/kg/day bw); NT/NM (Reddy and Kamble, 2014) AOTS (2500, 5000, 7500 and 10000 mg/kg bw), SATS (200 to 1000 mg/kg bw); NT/NM but mild and acceptable toxicity potential at high dose (Muthuraman and Singh, 2012) AOTS (175 to 5000 mg/kg), CTS (200 to 600 mg/kg bw for 60 days); NT/NM (P.D. Shah et al., 2012)	Rhizome showed antioxidant activities in DPPH (IC ₅₀ =79.9 µM trolox/100 g dw), ABTS (IC ₅₀ =8.66 µM trolox/100 g dw) and FRAP (IC ₅₀ =78.9 µM trolox/100 g dw) assays (Wojdylo et al., 2007). Methanol extract of leaf (IC ₅₀ =20 µg/ml) and rhizome (IC ₅₀ =20.8 µg/ml) showed activity in DPPH assay, methanol extract of leaf (IC ₅₀ =100.0 µg/ml) and rhizome (IC ₅₀ =30.5 µg/ml) showed Superoxide anion-scavenging activity, methanol extract of leaf (IC ₅₀ =18.8 µg/ml) and rhizome (IC ₅₀ =33.3 µg/ml) showed ability of chelating ferrous ion (Devi and Ganjewala, 2011).	NE
<i>Ageratum conyzoides</i> (L.) L.	Leaves; Topical application; 10% (w/w) petroleum ether, chloroform, methanolic and aqueous extracts; Wistar albino rats of either sex (150–250 g)	EWM (normal and infected), IWM, DSW; 0.2% (w/w) NFZ; Methanolic and aqueous extracts (Dash and Murthy, 2011a)	Leaf; Hydroalcoholic; Wistar rats of either sex (150–200 g) Whole plants; Alcoholic; Swiss albino mice (25–30 g) and Wistar albino rats of the both sexes (150–200 g)	AOTS (5000 mg/kg bw), SATS (500 and 1000 mg/kg bw); NT/NM (Diallo et al., 2010) AOTS (0.5, 1.0, 2.0, 3.0 and 4.0 g/kg bw); NT/NM (M.A. Rahman et al., 2012)	Whole plant methanol extract (IC ₅₀ =22.50 µg/ml) and essential oil (IC ₅₀ =570 µg/ml) showed activity in DPPH assay, whole plant methanol extract (IC ₅₀ =1290.00 µg/ml) and essential oil (IC ₅₀ =15.50 µg/ml) showed Lipid peroxidases activity, whole plant methanol extract (IC ₅₀ =4.48 mg/ml) and essential oil (IC ₅₀ =570 mg/ml) showed activity in FRAP assay (Patil et al., 2010). Crude ethanolic extract of leaves at the concentration of 100 µg/ml showed 91.72% inhibition in DPPH free radical scavenging assay (IC ₅₀ =18.91 µg/ml), ethanolic leaf extracts exhibited 2.011 average absorbance at 700 nm at 100 µg/ml concentrations in Reducing power assay, Ethanolic leaf extracts showed 76.0393% chelating ability at 100 µg/ml	NE

concentration (IC_{50} = 16.28 μ g/ml) in Ferrous ion chelating ability (Dewan et al., 2013).

Activities of antioxidant enzymes, such as superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPX) and glutathione S transferase (GST) were assessed in Male albino rats (Sprague–Dawley strain weighing 120–150 g) using aqueous extract of leaves. The activities of these enzymes returned to normalcy in all administered rats (Resmi et al., 2006).

Ethyl acetate fraction of Red onion peel was studied. Antioxidant activity (AOA) was 97.4% as performed by autoxidation of β -carotene and linoleic acid coupled reaction method, free radical scavenging activity (FRSA) was 75.3% as measured by using DPPH radical solution, Reducing power was 1.6 ascorbic acid equivalents per milligram as determined by ferric reducing antioxidant power assay (B.N. Singh et al., 2009).

Free radical scavenging activities (FRSA) in different layers ranged with IC_{50} values from 0.1 to 15.2 mg/ml (Prakash et al., 2007).

DPPH (IC_{50} = 87.50 μ g/ml), superoxide (IC_{50} = 98.80 μ g/ml), hydroxyl (IC_{50} = 106.25 μ g/ml), nitric oxide (IC_{50} = 88.70 μ g/ml) and ABTS (IC_{50} = 147.50 μ g/ml) radical scavenging activities were reported in methanol extract of leaves (B.S.A. Kumar et al., 2010).

DPPH (IC_{50} = 29 μ g/ml), scavenges superoxide (IC_{50} = 66–70 μ g/ml), hydrogen peroxide (IC_{50} = 120–125 μ g/ml), hydroxyl radicals (IC_{50} = 140–145 μ g/ml) and nitric oxide (IC_{50} = 135–140 μ g/ml) activities were reported in 50% ethanolic extract of whole plant (Zeashan et al., 2009).

Methanol extract of whole plant showed dose dependant free radical scavenging activity at concentrations of 0.05 to 1.0 mg/ml in DPPH free radical assay (Huang et al., 2012).

Purified Ovatodiolide from whole plant inhibited mushroom tyrosinase activity (IC_{50} = 0.253 mM), intracellular tyrosinase activity (IC_{50} = 0.469 mM) and decreased the amount of melanin (IC_{50} = 0.435 mM) in B16F10 cells. Methanol extract of whole plant showed minor inhibitory effects on melanogenesis in B16F10 melanoma on intracellular tyrosinase activity and melanin content in B16F10 cells (Huang et al., 2012).

NE

Ethanolic extract of leaves showed 99.07%, 89.77% and 73.64% values in ABTS, DPPH and nitric oxide radical

<i>Albizia lebbek</i> (L.) Benth.	Bark; Topical application; 5% and 10% (w/v) aqueous and ethanolic extracts and Oral application: 250 mg/kg and 500 mg/kg bw; Wistar rats (150–200 g)	EWM, IWM, DSW; 5% (w/w) PVI; 10% (w/v) ethanolic extract (Upadhyay et al., 2013)	Leaves; Methanolic; Wistar rats (150–200 g)	AOTS (2000 mg/kg bw); NT/NM up to 2000 mg/kg bw (Sivakumar et al., 2013)	Activities of antioxidant enzymes, such as superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPX) and glutathione S transferase (GST) were assessed in Male albino rats (Sprague–Dawley strain weighing 120–150 g) using aqueous extract of leaves. The activities of these enzymes returned to normalcy in all administered rats (Resmi et al., 2006).	NE
<i>Allium cepa</i> L.	Bulbs; Oral application; 300 mg/kg bw ether, chloroform, alcoholic and chloroform–water extracts; Wistar albino rats of either sex (150–200 g)	EWM, IWM, DSW; NA; Alcoholic extract (Shenoy et al., 2009a)	Dry skin and edible portion of bulbs; Methanolic; Swiss albino mice (25–35 g) of either sex	AOTS; NT/NM (Bhanot and Shri, 2010)	Ethyl acetate fraction of Red onion peel was studied. Antioxidant activity (AOA) was 97.4% as performed by autoxidation of β -carotene and linoleic acid coupled reaction method, free radical scavenging activity (FRSA) was 75.3% as measured by using DPPH radical solution, Reducing power was 1.6 ascorbic acid equivalents per milligram as determined by ferric reducing antioxidant power assay (B.N. Singh et al., 2009).	Quercetin 4'-O- β -D-glucopyranoside from dried skin showed IC_{50} value of 4.3 and 52.7 μ M using L-tyrosine or L-DOPA as a substrate (Arung et al., 2011).
<i>Amaranthus spinosus</i> L.	Leaves; Topical application; Benzene and alcoholic extracts; Rats	EWM, DSW, IWM; Benzene and alcoholic extract (S.G. Kumar et al., 2009)	Leaves; Ethanolic; Wistar albino rats of both sex (200–250 g)	AOTS (5, 50, 300 and 2000 mg/kg bw); NT/NM up to 2000 mg/kg bw (S.B. Mishra et al., 2012)	DPPH (IC_{50} = 87.50 μ g/ml), superoxide (IC_{50} = 98.80 μ g/ml), hydroxyl (IC_{50} = 106.25 μ g/ml), nitric oxide (IC_{50} = 88.70 μ g/ml) and ABTS (IC_{50} = 147.50 μ g/ml) radical scavenging activities were reported in methanol extract of leaves (B.S.A. Kumar et al., 2010).	NE
<i>Anisomeles indica</i> (L.) Kuntze	NE	NE	Aerial parts; Ethyl acetate and chloroform; Male and female Wistar rats (200–250 g)	AOTS (2000 mg/kg bw); NT/NM (Sundriyal et al., 2013)	DPPH (IC_{50} = 29 μ g/ml), scavenges superoxide (IC_{50} = 66–70 μ g/ml), hydrogen peroxide (IC_{50} = 120–125 μ g/ml), hydroxyl radicals (IC_{50} = 140–145 μ g/ml) and nitric oxide (IC_{50} = 135–140 μ g/ml) activities were reported in 50% ethanolic extract of whole plant (Zeashan et al., 2009).	Purified Ovatodiolide from whole plant inhibited mushroom tyrosinase activity (IC_{50} = 0.253 mM), intracellular tyrosinase activity (IC_{50} = 0.469 mM) and decreased the amount of melanin (IC_{50} = 0.435 mM) in B16F10 cells. Methanol extract of whole plant showed minor inhibitory effects on melanogenesis in B16F10 melanoma on intracellular tyrosinase activity and melanin content in B16F10 cells (Huang et al., 2012).
<i>Annona squamosa</i> L.	Leaves; Topical application; Ethanol extract (100 mg/kg bw); Streptozotocin-nicotinamide-induced diabetic rats	EWM (open); No reference control; Ethanolic extract	Seed; Crude extracts; Albino female Wistar rats (100–120 g)	AOTS (2000 mg/kg bw); 300 mg/kg bw is NT/NM, whereas 2000 mg/kg bw was toxic (Aneela et al., 2011)	Ethanolic extract of leaves showed 99.07%, 89.77% and 73.64% values in ABTS, DPPH and nitric oxide radical	NE

Table 3 (continued)

Plant species	Wound healing studies		Toxicological studies		Antioxidant activities	Tyrosinase inhibitory activities
	Plant part; Application; Extract; Animal (Weight)	Animal Model; Reference control; Effective treatment [Source for columns 2 & 3]	Plant part; Extract; Animal (Weight)	Animal model (Dose); Toxicity status [Source for columns 4 & 5]		
<i>Argemone mexicana</i> L.	Leaves; Topical application; Petroleum ether, chloroform, methanolic and aqueous extracts; Wistar albino rats of either sex (150–250 g)	(Ponrasu and Suguna, 2012) EWM (normal and infected), IWM, DSF; 0.2% (w/w) NFZ; Methanolic and aqueous extracts (Dash and Murthy, 2011b)	Root bark; Aqueous; Swiss mice of both sexes	AOTS (3, 5, 7 gm/kg bw); NT/NM even at high dose (Pingale, 2013)	scavenging activity at 1000 µg/ml concentration (Shirwaikar et al., 2004). Ethanol extract of roots possesses antioxidant activity at a dose of 100 µg/ml concentration, the extract showed high scavenging activity against DPPH (85.17%), ABTS (75.27%) and H ₂ O ₂ (84.25%) radicals (Perumal et al., 2010).	NE
			Whole plant; Crude extract; Mice (18–25 g)	AOTS; Toxic with LD ₅₀ value of 400 mg/kg bw (Ibrahim and Ibrahim, 2009)	Leaves extracts showed superoxide anion scavenging activity in Nitro blue tetrazolium assay with maximum free radical scavenging at a dosage of 200 µg/ml (Bhardwaj et al., 2011).	
<i>Azadirachta indica</i> A. Juss.	Leaves; Topical application; 5% (w/w) ethanolic extracts; Wistar albino rats of either sex (180–200 g)	EWM; 5% (w/w) PVI; Ethanolic extract (Purohit et al., 2013)	Stem bark; Ethanolic; Male albino rats of Wistar strains (174.40 ± 12.30)	AOTS (50,100, 200 and 300 mg/kg bw); only 50 mg/kg bw relatively safe other doses toxic (Ashafa et al., 2012)	Reducing power of bark extracts at 10 mg/ml concentration: 1.710 (80% ethanol extract), 1.460 (80% Methanol extract), 1.460 (80% Acetone extract) (Sultana et al., 2007). DPPH radical scavenging activity was recorded as 82.45%, 8.89%, 83.73%, 91.03%, 93.11% for methanol crude, Hexane, Ethyl acetate, n-Butanol and Water extracts of bark respectively, whereas DPPH radical scavenging activity was recorded as 12.56%, 9.00%, 15.00%, 33.25%, 5.83% for methanol crude, Hexane, Ethyl acetate, n-Butanol and Water extracts of leaf respectively (Chimeray et al., 2009). Hydroxyl (OH) radical scavenging activity was recorded as 87.84%, 84.89%, 86.38%, 87.87% and 92.12% for methanol crude, Hexane, Ethyl acetate, n-Butanol and Water extracts of bark respectively, whereas Hydroxyl (OH) radical scavenging activity was recorded as 36.47% (Methanol crude), 36.17% (Hexane), 38.72% (Ethyl acetate), 35.72% (n-Butanol), 43.86% for methanol crude, Hexane, Ethyl acetate, n-Butanol and Water extracts of leaf respectively (Chimeray et al., 2009).	Methanolic extract of leaf showed 38.3% Tyrosinase inhibition in mushroom tyrosinase inhibitory activity (Adhikari et al., 2008).
			Seed oil; Three generation study on rats of either sex.	Feed on diet with 10% oil; NT/NM (Chinnasamy et al., 1993)		
<i>Bambusa bambos</i> (L.) Voss	Leaves; Topical application; 0.5% and 1% (w/w) petroleum ether, chloroform and methanolic extracts; Albino rats of either sex (230 g)	EWM, IWM; 5% (w/w) FRA; Methanolic extract (Muniappan et al., 2014)	Leaves; Aqueous; Swiss albino mice of either sex (20 to 25 g)	AOTS (400, 800, 1000, 1200 mg/kg bw); NT/NM up to 1000 mg/kg bw (Kundu et al., 2011)	Aqueous, methanolic and butanolic extract of leaves showed DPPH radical scavenging activity with IC ₅₀ values of 964, 273 and 1103 µgm/ml (Macwan et al., 2010).	NE
<i>Bauhinia variegata</i> L.	Root; Topical application; Ethanolic and aqueous extracts	EWM, IWM; FRA; Methanolic and	Root; Aqueous and ethanolic; Wistar		Root water (IC ₅₀ =37.73 µ/ml), Root alcohol (IC ₅₀ =36.01 µ/ml), stem water	NE

	(200 and 400 mg/kg bw); Albino Wistar rats	ethanolic extract (Sharma, 2010)	albino rats of either sex (150–200 g)	AOTS (200, 400, 2000 mg/kg bw); NT/NM up to 2000 mg/kg bw (R.K. Sharma et al., 2011)	IC ₅₀ =45.85 µl/ml, stem alcohol (IC ₅₀ =30.50 µl/ml) extracts showed DPPH free radical scavenging activity, Root water (IC ₅₀ =478.80 µl/ml), Root alcohol (IC ₅₀ =415.20 µl/ml), stem water IC ₅₀ =405.80 µl/ml, stem alcohol (IC ₅₀ =362.90 µl/ml) showed Nitric oxide free radical scavenging activity, Root water (IC ₅₀ =502.10 µl/ml), Root alcohol (IC ₅₀ =445.30 µl/ml), stem water IC ₅₀ =481.30 µl/ml, stem alcohol (IC ₅₀ =414.60 µl/ml) also showed super oxide free radical scavenging activity, Root water (IC ₅₀ =11.74 µl/ml), Root alcohol (IC ₅₀ =10.78 µl/ml), stem water IC ₅₀ =10.23 µl/ml, stem alcohol (IC ₅₀ =9.85 µl/ml) showed Hydrogen peroxide free radical scavenging activity (Rajani and Ashok, 2009).	
<i>Boerhavia diffusa</i> L.	NE	NE	Leaves; Methanolic; Female Swiss albino mice and Wistar rats of either sex Leaves; Aqueous; Male Wistar albino rats (104–214 g) and albino mice (18.2–24.5 g)	AOTS (300 and 2000 mg/kg bw), Repeated dose toxicity (1000 mg/kg/day bw for 28 days); NT/NM in both tests (Kulkarni and Warriar, 2012) AOTS (2000 mg/kg bw), SATS (500, 1000 and 2000 mg/kg bw); NT/NM in both tests (Orisakwe et al., 2003)	Aqueous (IC ₅₀ =200.82 µg/ml), Methanolic (IC ₅₀ =327.40 µg/ml), Chloroform (IC ₅₀ =409.81 µg/ml), hexane (IC ₅₀ =2351.60 µg/ml) extracts of whole plant showed DPPH radical scavenging activity (Guha et al., 2011). BuOH fraction of leaf controls glucose metabolism and reduces lipid peroxidation as well as the level of oxygen radicals in rats with streptozotocin-induced diabetes (Kim et al., 2003).	NE
<i>Brassica juncea</i> (L.) Czern.	Leaves; Topical application; 200 mg/kg petroleum ether, chloroform, ethanolic and aqueous extracts; Male Wistar albino rats (180–200 g)	EWM; 5% (w/w) PVI; aqueous extract (Malan et al., 2011)	Leaves; Petroleum ether, chloroform, ethanol and water; Sprague Dawley rats	Acute dermal toxicity-fixed dose method (1000–2000 mg/kg bw topically); NT/NM (Malan et al., 2011)		NE
<i>Buchanania cochinchinensis</i> (Lour.) M. R.Almeida	Fruits; Oral administration; 300 mg/kg po ethanolic extracts; Albino rats of either sex (150–200 g)	EWM, IWM, DSW; 0.17 mg/kg DXM; Ethanolic extract (Chitra et al., 2009)	Seed; Methanolic; Female Swiss albino mice (25–30 g)	AOTS (50, 300 and 2000 mg/kg bw); NT/NM (Singh and Bothara, 2012)	Ferric reducing antioxidant power (FRAP) at 500 µg/ml of methanol extracts of bark is 49.05 mg GAE/g of extracts (Venkata et al., 2012).	NE
<i>Butea monosperma</i> (Lam.) Taub.	Bark; Topical and oral applications; 25 mg/kg bw and 50 mg/kg bw ethanolic extract; Wistar albino rats (150 to 200 g) Bark; Topical application; 200 µL of alcoholic extract; Male albino Wistar rats (150–200 g)	EWM, IWM, DSW; SOF; Flavonoid fraction from the ethanolic extract (Muralidhar et al., 2013) EWM, IWM; NA; NA (Sumitra et al., 2005)	Kernels; Methanolic; Wistar rats of either sex (150–200 g) Seed; Powder suspension; Charles Foster strain albino rats of either sex (180–220 g)	AOTS (2000 mg/kg bw); NT/NM (Warokar et al., 2010) CTS (800 mg/kg/day bw powder suspension given orally for 90 days); Toxic effect when administered in powder form (Donga et al., 2011)	Ethyl acetate (EC ₅₀ =15.44 µg/ml) and butanol (EC ₅₀ =41.53 µg/ml) fractions showed DPPH free radical scavenging activity, whereas aqueous fractions showed no inhibition (Lavhale and Mishra, 2007). DPPH free radical scavenging activity of Hydromethanol extract of leaves showed IC ₅₀ value of 25.96 µg/ml (Hasan et al., 2009).	NE
<i>Caesulia axillaris</i> Roxb.	NE	NE	Essential oil; Mice (30 g)	Essential oil from 0.05 to 0.5 ml were orally given; LD ₅₀ value was calculated as 9166.6 µl/kg bw (S.B. Mishra et al., 2012)	Essential oil showed DPPH radical scavenging activity with IC ₅₀ value of 18 µl/ml (S.B. Mishra et al., 2012).	NE
<i>Callicarpa macrophylla</i> Vahl	NE	NE	NE	NE	NE	NE
<i>Calotropis procera</i>		EWM; NA; Latex (Rasik et al., 1999)	Root barks; Aqueous and hydroalcoholic;	AOTS (2000 mg/kg bw both extract), SATS (20 mg/kg/day bw aqueous extract	Methanol extract of leaves showed DPPH free radical scavenging activity with EC ₅₀	NE

Table 3 (continued)

Plant species	Wound healing studies		Toxicological studies		Antioxidant activities	Tyrosinase inhibitory activities
	Plant part; Application; Extract; Animal (Weight)	Animal Model; Reference control; Effective treatment [Source for columns 2 & 3]	Plant part; Extract; Animal (Weight)	Animal model (Dose); Toxicity status [Source for columns 4 & 5]		
(Aiton) Dryand.	Latex; Topical application; 1.0% sterile solution; Male Guinea Pigs of Swiss strain (NA)		Male and female NMRI mice (27 ± 4 g) and Wistar rats of either sex (160 ± 42 g) Latex; Fed through diet; Black rat, <i>Rattus rattus</i> .	for 3 and 6 weeks); Aqueous extract relatively safe as compared to hydroalcoholic extract (Ouedraogo et al., 2013). Toxicity study (concentrations of 5, 7.5 and 10% w/w); Toxic (Pahwa and Chatterjee, 1988)	value of 110.25 $\mu\text{g/ml}$ (Yesmin et al., 2008). DPPH free radical scavenging activities of methanol extract of stem was 93% at 1.3 ml concentration and leaf was 88% at 1.5 ml concentration (Patel et al., 2010). Showed 5% inhibition of DOPA auto-oxidation at concentration of 500 $\mu\text{g/ml}$ (Lee et al., 1997).	
<i>Cannabis sativa</i> L.	NE	NE	Leaves; Aqueous, alcoholic and chloroform; Female Wistar albino rats (120 to 170 g) Seed oil; direct; Wistar albino rats of different sex (100–150 g)	AOTS; NT/NM up to 4000 mg/kg bw (Zade et al., 2013) Toxicity study (Orally at daily doses of 0.01, 0.1 and 1 ml/kg/day bw); Toxic (Dahab et al., 2013)		Whole plant extract showed 3% inhibition of tyrosinase at concentration of 333 $\mu\text{g/ml}$ (Lee et al., 1997).
<i>Cassia fistula</i> L.	Leaves/Topical application: alcoholic extract	Albino rat (NA): NA; NA; NA (Senthilkumar et al., 2006b)	Seed; Methanolic; Mice Pod pulp; direct; Rats and mice	AOTS (5000 mg/kg bw); NT/NM (Jothy et al., 2011) AOTS; NT/NM (Sakulpanich et al., 2012)	Ethanol extracts of leaves and methanol extracts of stem bark, pulp and flowers showed moderate antioxidant activity in linoleic acid peroxidation system between 37 and 62% at concentration of 0.2 mg/ml (Siddhuraju et al., 2002). Stem bark (93%), leaves (74.9%), BHT (37.8%), flowers (33.2%) and pulp (15.7%) extracts showed DPPH radical scavenging activity (Siddhuraju et al., 2002).	Methanol extract of pods showed tyrosinase inhibitory activity with IC_{50} value of 39.2 μg (Khan et al., 2013).
<i>Celastrus paniculatus</i> Willd.	Leaves; Topical application; Petroleum ether extract (8 mg/ml of 0.2% sodium alginate gel); Swiss albino rats (175–225 g)	EWM, IWM, DSW; NFZ; Triterpene compound lupeol from petroleum ether extract (Harish et al., 2008)	Stem bark; Alcoholic; Rats Whole Plant; Methanolic; Swiss albino mice (18–22 g) and Wistar albino rats (150–180 g) of either sex Seeds; Ethanolic and methanolic; Albino mice (25–32 g)	AOTS (2000 mg/kg bw); NT/NM (Agnihotri and Singh, 2013) AOTS (2000 mg/kg bw); NT/NM (Atigari et al., 2012) AOTS (1000 and 5000 mg/kg bw); NT/NM (Parimala et al., 2009)	Aqueous, chloroform, and methanolic extracts of the seeds showed 99%, 95.2%, 92% inhibition respectively in DPPH radical scavenging assay, whereas chloroform, methanolic and aqueous extracts seeds demonstrated 806.5 $\mu\text{mol Trolox/g}$ extract, 705.25 $\mu\text{mol Trolox/g}$ extract, 609.5 $\mu\text{mol Trolox/g}$ extract ABTS scavenging activity respectively in Trolox equivalent antioxidant capacity (TEAC) assay (Arora and Pandey-Rai, 2014). Chloroform, methanolic and aqueous extracts of seed showed FRAP value of 245 $\mu\text{mol Fe(II)/g}$ extract, 211.11 $\mu\text{mol Fe(II)/g}$ and 210 $\mu\text{mol Fe(II)/g}$ extract respectively in Ferric reducing antioxidant power assay (FRAP),	NE

<i>Chrysopogon zizanioides</i> (L.) Roberty	NE	NE	Root; Methanolic; Wistar albino rats (200–220 g) of either sex	AOTS (5000 mg/kg bw); NT/NM (Parmar et al., 2008)	whereas aqueous, methanolic and chloroform extract showed 52.8%, 43.7 ± 2.85% and 42.98% inhibition in Lipoxigenase inhibition assay (Arora and Pandey-Rai, 2014). DPPH activity, Total antioxidant capacity (TAC), Reducing Power (RP), Ferric reducing antioxidant powder assay (FRAP) power increased with the increase in concentration of the extracts (Luqman et al., 2009). Vetiver oil (10 µL/ml) dissolved in methanol exhibited ~93% free radical scavenging activity in the DPPH assay and ~34% Fe ²⁺ chelating activity in the metal chelating assay (Kim et al., 2005). DPPH free radical scavenging activity of the alkaloidal fraction of roots showed IC ₅₀ value of 63.44 µg/ml (Bafna and Mishra, 2010). 50% ethanol extract of roots showed significant antioxidant activity in the DPPH assay and was found to significantly scavenge superoxide, hydrogen peroxide, hydroxyl radicals, and nitric oxide at a dose regimen of 50 to 400 µg/kg <i>in vitro</i> (Amresh et al., 2007a).	Essential oil markedly decreased tyrosinase activity in α-melanin-stimulating-hormone stimulated B16 cells (Peng et al., 2014).
<i>Cissampelos pareira</i> L.	NE	NE	Oil; direct; Albino mice of either sex (20–25 g) and Albino rats of either sex (150–225 g) Whole plant; Hydroethanolic; Rats	AOTS (884, 1250, 1768, 2500, 3536 and 5000 mg/kg bw), SATS (100, 250, 500 and 1500 mg/kg/day bw); NT/NM in Both tests, LD ₅₀ value 2985.38 mg/kg bw (Tripathi et al., 2006) AOTS (2 g/kg bw), SATS (1 and 2 g/kg/day bw); NT/NM in Both tests (Amresh et al., 2008)	70% methanolic extract of leaves showed high free radical scavenging activity in DPPH (IC ₅₀ =373.18 µg/ml) and hydroxyl radical (IC ₅₀ =573.55 µg/ml) assays (Gupta et al., 2011). Ethanol extract of leaves showed DPPH scavenging activity of 92.6% at 250 µg/ml concentration, whereas chloroform and petroleum ether extracts showed 52.2% and 16.7% inhibition at the same concentration. Ethanol extract of leaves exhibited 68.58%, 62.06% and 52.65% hydroxyl, superoxide anion and nitric oxide radical activity at 250 µg/ml respectively (Gouthamchandra et al., 2010).	NE
<i>Cleome viscosa</i> L.	Leaves and whole plant; Topical application; 5% and 10% (w/w) methanolic extract; Wistar albino rats of either sex (130–180 g)	EWM, IWM; 0.2% (w/w) NEO; 5% (w/w) methanolic extract (Mohammad and Mazumder, 2013)	Leaves and whole plant; Ethanolic; Wistar albino rats of either sex	AOTS (2 g/kg bw); NT/NM (Panduraju et al., 2011)	70% methanolic extract of leaves showed high free radical scavenging activity in DPPH (IC ₅₀ =373.18 µg/ml) and hydroxyl radical (IC ₅₀ =573.55 µg/ml) assays (Gupta et al., 2011). Ethanol extract of leaves showed DPPH scavenging activity of 92.6% at 250 µg/ml concentration, whereas chloroform and petroleum ether extracts showed 52.2% and 16.7% inhibition at the same concentration. Ethanol extract of leaves exhibited 68.58%, 62.06% and 52.65% hydroxyl, superoxide anion and nitric oxide radical activity at 250 µg/ml respectively (Gouthamchandra et al., 2010).	Ethanolic extract of leaf at 200 µg/ml concentration showed 31.47% inhibition in tyrosinase inhibitory activity using DOPA chrome method (Meechai et al., 2010).
<i>Clerodendrum infortunatum</i> L.	NE	NE	Leaves; Methanolic; Adult male Swiss albino mice (18–25 g)	AOTS (2000 mg/kg bw), SATS (500 mg/kg bw); NT/NM (Das et al., 2011)	Ethanol extract of leaves showed DPPH scavenging activity of 92.6% at 250 µg/ml concentration, whereas chloroform and petroleum ether extracts showed 52.2% and 16.7% inhibition at the same concentration. Ethanol extract of leaves exhibited 68.58%, 62.06% and 52.65% hydroxyl, superoxide anion and nitric oxide radical activity at 250 µg/ml respectively (Gouthamchandra et al., 2010).	NE
<i>Commelina benghalensis</i> L.	Root; Topical application; alcoholic and aqueous extracts; Albino rats (NA)	EWM, IWM, DSW; NA; NA (Sambrekar et al., 2011)	Leaves; Hydroethanolic; Wistar albino female rats (130–150 g)	AOTS (2000 mg/kg bw), SATS (200 and 400 mg/kg/day bw for 14 days); NT/NM in both tests (Tiwari et al., 2013)	Methanol extract of aerial part exhibited strong DPPH radical scavenging activities (IC ₅₀ =138.7 µg/ml) (Anusuya et al., 2012).	NE
<i>Crotalaria juncea</i> L.	NE	NE	Leaves; Ethanolic; Male albino Wistar rats (150–250 g)	AOTS (2000 mg/kg bw); NT/NM (Ashok et al., 2006)	Seed oil showed IC ₅₀ value of 122.52 µg/ml for the DPPH radical scavenging activity and IC ₅₀ value of 286.409 µg/ml for hydroxyl radical scavenging activity (Chouhan et al., 2011).	NE
<i>Cryptolepis dubia</i> (Burm. f.) M.R. Almeida	NE	NE	Leaves and whole plant; Mice, rat and rabbit	AOTS (0.324, 0.31, 0.356 g/kg bw); NT/NM (Minh and Tuan, 2013)	NE	NE
	Tubers; Topical application; 200 and 400 mg/kg bw petroleum	EWM; 0.2% (w/w) NFZ; 200 mg/kg			Methanolic extract enhanced the antioxidant defense against reactive	NE

Table 3 (continued)

Plant species	Wound healing studies		Toxicological studies		Antioxidant activities	Tyrosinase inhibitory activities
	Plant part; Application; Extract; Animal (Weight)	Animal Model; Reference control; Effective treatment [Source for columns 2 & 3]	Plant part; Extract; Animal (Weight)	Animal model (Dose); Toxicity status [Source for columns 4 & 5]		
<i>Curculigo orchioides</i> Gaertn.	ether, chloroform, ethyl acetate and methanolic extracts; Wistar albino rats (150–200 g)	methanolic extract (Agrahari et al., 2010a)	Rhizome; Hydroalcoholic; Wistar rats	AOTS (50, 100, 300, 1000, and 2000 mg/kg bw); NT/NM up to 2000 mg/kg bw (Asif and Kumar, 2010)	oxygen species produced under hyperglycemic conditions (Anandakirouchenane et al., 2013). Hydroxyl radical scavenging activities were reported in eight compounds isolated from rhizome viz., Orcinol glucoside (IC ₅₀ =1.39 mM), orcinol-1-O-β-D-glucopyranosyl-(1→6)-β-D-glucapyranoside (IC ₅₀ =0.87 mM), orcinol-1-O-β-D-apiofuranosyl-(1→6)-β-D-glucopyranoside (IC ₅₀ =1.17 mM), curculigoside (IC ₅₀ =0.54 mM), curculigoside B (IC ₅₀ =1.11 mM), curculigoside C (IC ₅₀ =0.25 mM), 2,6-dimethoxyl benzoic acid (IC ₅₀ =1.51 mM) and syringic acid (IC ₅₀ =2.61 mM) (Wu et al., 2005). Superoxide anion radical scavenging activities were reported in eight compounds isolated from rhizome viz., Orcinol glucoside (IC ₅₀ =2.49 mM), orcinol-1-O-β-D-glucopyranosyl-(1→6)-β-D-glucapyranoside (IC ₅₀ =1.56 mM), orcinol-1-O-β-D-apiofuranosyl-(1→6)-β-D-glucopyranoside (IC ₅₀ =1.84 mM), curculigoside (IC ₅₀ =1.35 mM), curculigoside B (IC ₅₀ =1.48 mM), curculigoside C (IC ₅₀ =0.88 mM), 2,6-dimethoxyl benzoic acid (IC ₅₀ =3.21 mM) and syringic acid (IC ₅₀ =3.46 mM) (Wu et al., 2005).	
<i>Cuscuta reflexa</i> Roxb.	NE	NE	Stem; Aqueous and alcoholic; Albino mice of either sex (16–20 g)	AOTS (2000 mg/kg); NT/NM (Katiyar et al., 2012)	Methanolic extract showed DPPH radical scavenging activity with IC ₅₀ value of 359.48 µg/ml (Vijikumar, 2011). Alcoholic extract showed DPPH radical scavenging activity with IC ₅₀ value of 6.89 mg/ml (Sakshy et al., 2012). Methanolic leaf extracts showed total antioxidant capacity of 10.48 µg/ml AAE in phosphomolybdenum method, with reducing power of 0.726 (Krishanti et al., 2010). Water (IC ₅₀ =152.96 mg/L), methanol (IC ₅₀ =203.29 mg/L) and acetone (IC ₅₀ =206.69 mg/L) extracts of whole plants showed activity in DPPH free radical assay, whereas water (IC ₅₀ =57.21 mg/L) and acetone	Methanolic extract of whole plant showed 28.9% Tyrosinase inhibition in mushroom tyrosinase inhibitory assay (Adhikari et al., 2008).
<i>Cynodon dactylon</i> (L.) Pers.	Whole plant; Topical application; 11.5% (w/w) hydroalcoholic extract; Male Wistar albino rats (180–200 g)	EWM; PVI; Hydroalcoholic extract (Kumar and Kashyap, 2013)	Whole plant; Chloroform, ethanolic extract; Mice	AOTS (100–2000 mg/kg bw); NT/NM (Yogesh et al., 2013)		NE

<i>Dalbergia sissoo</i> DC.	NE	NE	Bark; Alcoholic extract; Wistar rats	AOTS (50, 100, 300, 1000, and 3000 mg/kg bw); NT/NM up to dose of 3000 mg/kg bw (Mojahid-ul-Islam and Elhddad, 2012)	(IC ₅₀ =179.77 mg/L) showed activity in ABTS assay (Khlifi et al., 2013). Aqueous (IC ₅₀ =12.23 µg/ml) and methanol (IC ₅₀ =23.63 µg/ml) extract of the stem bark showed DPPH radical scavenging activity (Roy et al., 2011).	NE
<i>Datura stramonium</i> L.	NE	NE	Seeds; Alkaloids; Male and female Albino Wistar rats	AOTS; NT/NM (Abdelouahab et al., 2011)	Leaf extract showed free lipid peroxidation activity with IC ₅₀ value of 1.45 mg/ml (Kumar et al., 2008).	NE
<i>Dendrophthoe falcata</i> (L.f.) Ettingsh.	NE	NE	Leaves; Ethanolic; Swiss albino mice of either sex (20–25 g)	Brine shrimp lethality assay for general toxicity; Low level of general toxicity (LC ₅₀ 100 µg/ml) (Hasan et al., 2006)	Ethanolic extract of aerial parts showed potent antioxidant activity by lipid peroxidation inhibition, reduced glutathione, superoxide dismutase levels and increased the catalase activity (Pattanayak and Sunita, 2008).	NE
<i>Desmodium gangeticum</i> (L.) DC.	NE	NE	Aerial parts; Petroleum ether, benzene, chloroform, acetone and alcohol; Healthy albino Wistar rats (200–250 g)	AOTS (2000 mg/kg bw); NT/NM (Bisht and Bhattacharya, 2013)	Ethanolic extract of leaves showed potent antioxidant activity in DPPH, superoxide scavenging, reducing power, nitric oxide scavenging, 2,2 azino-bis(3-ethylbenzothiazoline-6-sulfonate), hydroxyl radical, ferric reducing antioxidant power and chelating ability assays (Venkatachalam and Muthukrishnan, 2012). Hydroalcoholic extract of the aerial parts strongly scavenged DPPH radical with the IC ₅₀ being 2.01 mg/ml, total antioxidant capacity of the extract was 149.91 nmol ascorbic acid/g, extract inhibited the ferrylbipyridyl (chromogen) formation in a dose dependent fashion with the IC ₅₀ value of 0.115 mg/ml (Govindarajan et al., 2003).	NE
<i>Desmostachya bipinnata</i> (L.) Stapf	NE	NE	Roots; Alcoholic and aqueous; Female albino mice (20–30 g)	AOTS (2000 mg/kg bw); NT/NM (Hegde et al., 2010)	Hydroalcoholic extract showed significant antioxidant activity in a dose-dependent manner with an IC ₅₀ value of 264.18 µg/ml in H ₂ O ₂ scavenging assay (Golla and Bhimathati, 2014).	NE
<i>Dicliptera paniculata</i> (Forssk.) L. Darbysh.	NE	NE	Leaves and stem; Aqueous, methanol and butanolic; Wistar rats (150–250 g)	AOTS (10 to 5000 mg/kg bw); NT/NM up to 5000 mg/kg bw (Abdulazeez et al., 2010)	NE	NE
<i>Eclipta prostrata</i> (L.) L.	NE	NE	Whole plant; Ethanolic; Swiss albino mice and Wistar albino rats	AOTS (0.5, 1.0, 2.0, 3.0 and 4.0 g/kg bw); NT/NM up to 4 g/kg bw (Md.A. Rahman et al., 2012)	Trolox equivalent antioxidant capacity TEAC of methanolic and aqueous extracts of aerial parts was 54.3 µmol trolox/100 g DW and 148.1 µmol trolox/100 g DW (Cai et al., 2004).	NE
<i>Ehretia laevis</i> Roxb.	NE	NE	NE	NE	Ethanol extract of leaves and stem showed DPPH radical scavenging activity with IC ₅₀ values of 2.44 and 29.88 µg/ml respectively (Torane et al., 2011).	NE
<i>Eulaliopsis binata</i> (Retz.) C.E.Hubb.	NE	NE	NE	NE	NE	NE
<i>Euphorbia hirta</i> L.	Whole plant; Topical application; 2% (w/w) ethanolic extract; Male Wistar albino rats (150–200 g);	BWM; 0.2% (w/w) NFZ; 2% w/w ethanolic extract (Chandramohan and Reddy, 2006)	Fruits; Ethanolic and aqueous suspended in 0.5% w/v Sodium-CMC; Adult Swiss albino mice (20–25 g)	AOTS (1000, 2000, 3000, 4000, 5000 mg/kg bw); Toxic, morality was observed at a dose of 2500 and 3000 mg/kg (P. Das et al., 2010)	Methanolic extract of leaves (72.96%), flowers (52.45%), roots (48.59%) and stems (44.42%) exhibited potent DPPH scavenging activity, where IC ₅₀ values of methanol extract of leaves, flowers, roots and stems were 0.803, 0.972, 0.989 and	NE

Table 3 (continued)

Plant species	Wound healing studies		Toxicological studies		Antioxidant activities	Tyrosinase inhibitory activities
	Plant part; Application; Extract; Animal (Weight)	Animal Model; Reference control; Effective treatment [Source for columns 2 & 3]	Plant part; Extract; Animal (Weight)	Animal model (Dose); Toxicity status [Source for columns 4 & 5]		
<i>Euphorbia thymifolia</i> L.	Whole plant; Topical application; 200, 400 and 600 mg/kg, bw total flavonoid fraction from methanolic extract; Male Wistar rats (175 g) NE	EWM, IWM, DSW; DXM (0.34, i.m), PVI; 400 and 600 mg/kg methanolic extract (Bigoniya et al., 2013) NE	Whole Plant; Alcohol, chloroform, and aqueous; Albino mice (25–30 g)	AOTS (50 to 2000 mg/kg bw); NT/NM up to 2000 mg/kg bw (Mamatha et al., 2014)	1.358 mg/ml, respectively (Basma et al., 2011). Methanol, chloroform, ethylacetate, n-butanol and water fractions and pure compounds (3-O-galloyl-4,6-(5)-HHDP-D-glucose, rugosin B and 1,3,4,6-tetra-O-galloyl-K-beta-D-glucose) possessed antioxidant activities (Lin et al., 2002).	NE
<i>Ficus benghalensis</i> L.	Bark; Topical application; 10% and oral application: 200 mg/kg/day ethanolic and aqueous extracts; Female Wistar albino rats (200–220 g)	EWM, IWM; NA; Both Ethanolic and aqueous extracts (Garg and Paliwal, 2011)	Bark; Ethanol and aqueous; Female Albino Wistar rats (200–220 g)	AOTS (5, 50, 300 and 2000 mg/kg bw); NT/NM up to 2000 mg/kg bw (Garg and Paliwal, 2011)	Aqueous extract of stem bark showed dose dependent antioxidant activity in microsomal lipid peroxidation inhibition assay with IC ₅₀ value of 80.24 µg/ml (Satish et al., 2013).	NE
<i>Ficus racemosa</i> L.	Roots; Topical application; aqueous and ethanolic extracts; Wistar albino rats of either sex (180–200 g)	EWM, IWM; PVI; Aqueous extract (Murti and Kumar, 2012)	Bark; Methanolic; Female Wistar rats (150–180 g)	AOTS (2000 mg/kg bw); NT/NM (Choudhury and Jadhav, 2013)	IC ₅₀ values of the root bark and heartwood in DPPH radical scavenging activity assay were 5.80 µg/ml and 4.49 µg/ml respectively (Jain et al., 2013).	Ethanolic extract of wood at 200 µg/ml concentration showed 56.41% inhibition in Tyrosinase inhibitory activity using DOPA chrome method (Meechai et al., 2010).
<i>Ficus religiosa</i> L.	Leaves; Topical application; 5% and 10% Hydro alcoholic extract; Wistar albino rats of either sex (150–250 g) Leaves; Topical application; 300 mg/kg ethanolic extract; Male Wistar rats (150–200 g)	EWM, IWM; 5% (w/w) PVI; 10% Hydro alcoholic extract (Roy et al., 2009) EWM, IWM, DSW; FSC; 300 mg/kg ethanolic extract (Charde et al., 2010)	Bark; Methanolic; Swiss albino male mice	AOTS (125, 250 and 500 mg/kg bw); NT/NM (Sreelekshmi et al., 2007)	Ethylacetate root extract scavenged DPPH radical (87.61%) at 250 µg/ml and hydrogen peroxide (70.25%) at 1000 µg/ml (Sharma and Gupta, 2007).	Mushroom tyrosinase inhibitory activity of methanolic extract of seeds showed 53.73% inhibition (Mukherjee et al., 2001)
<i>Holarrhena pubescens</i> Wall. ex G. Don	NE	NE	Seed; Ethanolic; Female albino Wistar rats	AOTS (2000 mg/kg bw); NT/NM (Saha and Subrahmanyam, 2013)	Ethyl acetate fraction of seed showed IC ₅₀ values of 3.38, 15.6, 16.5 and 30.6 µg/ml for the hydroxyl radical scavenging, hydrogen peroxide scavenging, nitric oxide scavenging and inhibition in lipid peroxidation assays respectively (Ali et al., 2011).	NE
<i>Holoptelea integrifolia</i> Planch.	Leaves and stem bark; Topical application; 5% (w/w) methanolic extract; Wistar rats (160–180 g)	EWM, IWM; 0.2% (w/w) NFZ; Methanolic extract (Reddy et al., 2008)	Leaves; Petroleum ether and methanolic; Albino mice of either sex (20–25 g)	AOTS (2000 mg/kg bw); NT/NM (Sutar et al., 2014)	IC ₅₀ values of methanolic extract of leaves and stem bark in DPPH free radical scavenging activity using HPLC method was 50.36 and 37.66 µg/well respectively (Reddy et al., 2008).	NE
<i>Hyptis suaveolens</i> (L.) Poit.	Leaves; Topical and oral application; 500 mg/kg petroleum ether, alcoholic and aqueous extracts; Wistar albino rats of either sex (1500–200 g) and Swiss albino mice of either sex (18–22 g)	EWM, IWM, DSW; NA; Petroleum ether extract (Shenoy et al., 2009b)	Leaves; Petroleum ether, solvent ether, chloroform, alcohol, chloroform and water; Swiss albino mice of either sex (18–22 g)	AOTS (5000 mg/kg bw); NT/NM, LD ₅₀ was found to be more than 5000 mg/kg bw (Shenoy et al., 2009a)	Antioxidant potential of essential oil determined by the DPPH method expressed as IC ₅₀ was 3.72 mg/ml and TEAC value determined by ABTS assay was 65.02 µg/mg (Nantitanon et al., 2007).	NE

<i>Ipomoea carnea</i> Jacq.	Flowers; Topical and oral application; 200 mg/kg/day bw Flavonoid compounds; Male Wistar rats (150–200 g)	EWM, IWM; ST; NA (Ambiga et al., 2007)	Leaves; Aqueous; Wistar albino rats	AOTS (5, 50, 300, 500 and 2000 mg/kg bw); Dose at 2000 mg/kg bw showed the toxic symptoms (Khalid et al., 2011)	n-butanol fraction showed 91.11% inhibition of DPPH free radical scavenging activity with IC ₅₀ value of 74.65 µg/ml, whereas Ethyl acetate soluble fraction showed FRAP value and lipid peroxidation inhibition value of 511.99 µg of trolox equivalents and 61.87%, respectively (Abbasi et al., 2010b).	
<i>Lannea coromandelica</i> (Houtt.) Merr.	Bark; Topical application; 10% (w/w) ethanolic and acetone extracts; Male Wistar rats (150–200 g)	EWM, IWM; 1% FSC; Both (Sathish et al., 2010)	Bark; Methanolic; Mice	AOTS (100 to 1000 mg/kg bw); NT/NM up to 1000 mg/kg bw (Majumder et al., 2013)	Methanolic extract of bark showed antioxidant activity with IC ₅₀ values of 12.32 and 34.72 µg/ml for DPPH and total ROS scavenging methods respectively (Alam et al., 2012).	NE
<i>Lantana camara</i> L.	Leaf; Topical application; 100 mg/kg/day bw aqueous extract; Sprague Dawley rats (200–220 g)	EWM, IWM; NA; Both (Nayak et al., 2009)	Leaves; Methanolic; Adult mice	AOTS (2 g/kg bw); NT/NM (Pour et al., 2011)	Aqueous extract of leaves exhibited antioxidant activity in DPPH radical scavenging assay (IC ₅₀ =42.66 µg/ml) and metal chelating activity assay (IC ₅₀ =1036.4 µg/ml) (Kalita et al., 2012).	NE
<i>Lawsonia inermis</i> L.	Leaf; Topical application; 5% and 10% ethanolic extract; Wistar albino rats (200–250 g) Leaves; Oral and topical application; Petroleum ether, Chloroform, Ethanol and aqueous extracts; Wistar rats of either sex (150–200 g)	IWM; NA; 10% ethanolic extract (Abdulla et al., 2009) EWM, IWM; NA; Ethanolic extract (Sakarkar et al., 2004)	Leaves; Ethanolic; Wistar albino rats of either sexes (200–250 g)	AOTS (100, 500, 1000 and 2000 mg/kg bw), SATS (200, 500 and 1000 mg/kg/day bw); Lethal dose of this plant is 2000 mg/kg bw and doses above 1000 mg/kg bw are toxic (Kaur et al., 2014)	Aqueous (IC ₅₀ =32.27 µg/ml), methanolic (IC ₅₀ =32.87 µg/ml), chloroform (IC ₅₀ =99.92 µg/ml) and hexane (IC ₅₀ =286.91 µg/ml) extracts of whole plant showed DPPH radical scavenging activity, aqueous (IC ₅₀ =12.57 in µg), methanolic (IC ₅₀ =12.59 in µg) and chloroform (IC ₅₀ =41.42 in µg) extracts of whole plant showed activity in ABTS assay, aqueous (IC ₅₀ =313.93 in µg), methanolic (IC ₅₀ =430.80 in µg) and chloroform (IC ₅₀ =1815.67 in µg) extracts of whole plants showed activity in FRAP assay (Guha et al., 2011). DPPH free radical scavenging assay of methanol extract of leaves showed 60.57% inhibition (Veerabadran et al., 2013). 100% methanol, 80% methanol, 100% ethanol and 80% ethanol extract of seed showed 83.6%, 81.3%, 42.2% and 87.5% scavenging in DPPH assay (Anwar and Przybylski, 2012).	More than 50% reduction in tyrosinase activity was achieved by a methanol extract at a concentration of 1140 mg/L (Gholamhoseinian et al., 2009).
<i>Leonotis nepetifolia</i> (L.) R.Br.	Leaves; Topical application; 1 and 2 g ethanolic extracts; Wistar albino rats (150–180 g)	EWM; 2% (w/w) SOF; Ethanolic extract (Nithya et al., 2011)	Stem bark; Methanolic; Swiss albino mice (18–30 g) of either sex	AOTS (10, 100, 1000, 1600, 2900 and 5000 mg/kg bw); NT/NM, LD ₅₀ was found to be 3807.9 mg/kg bw (Ayanwuyi et al., 2009)	DPPH free radical scavenging assay of methanol extract of leaves showed 60.57% inhibition (Veerabadran et al., 2013).	NE
<i>Linum usitatissimum</i> L.	Seed oil; Topical application; 0.75 and 1.5% in Eucerin-vaselin ointment; Male Wistar rats (190–210 g)	EWM; NA; Both (Farahpour et al., 2011)	Fixed seed oil; direct; Swiss albino mice of either sex	AOTS (20 ml/kg bw), SATS (3.0 ml/kg bw); NT/NM (Kaithwas et al., 2011)	100% methanol, 80% methanol, 100% ethanol and 80% ethanol extract of seed showed 83.6%, 81.3%, 42.2% and 87.5% scavenging in DPPH assay (Anwar and Przybylski, 2012).	NE
<i>Litsea glutinosa</i> (Lour.) C.B. Rob.	Whole plant; Topical application; 3% and 5% (w/w) ethanolic extract; Wistar rats of either sex (100–175 g)	EWM, IWM; 0.2% (w/w) NFZ; Both (Devi and Meera, 2010)	Whole plant; Aqueous; Wistar rats (100–175 g) of either sex	AOTS (2000 mg/kg bw); NT/NM (Devi and Meera, 2010)	Antioxidant activities of methanolic stem and leaf extracts in DPPH assay were recorded as 23.02% and 31.42% respectively (Kshirsagar and Upadhyay, 2009).	NE
<i>Mallotus philippensis</i> (Lam.) Mull. Arg.	NA	NA	Leaves; Methanolic; Female mice	AOTS (2000 mg/kg bw); NT/NM (Ramakrishna et al., 2011)	Methanolic extracts of leaf showed 91.33% inhibition in DPPH reduction assay (Ghimire et al., 2011). Total antioxidant activity (TAA) of bark extract was 5.27 mmol Trolox equivalents/g, whereas TAA of other extracts ranged from 0.05 to 1.79 mmol Trolox equivalents/g extract (Arfan et al., 2007).	NE

Table 3 (continued)

Plant species	Wound healing studies		Toxicological studies		Antioxidant activities	Tyrosinase inhibitory activities
	Plant part; Application; Extract; Animal (Weight)	Animal Model; Reference control; Effective treatment [Source for columns 2 & 3]	Plant part; Extract; Animal (Weight)	Animal model (Dose); Toxicity status [Source for columns 4 & 5]		
<i>Martynia annua</i> L.	Leaves; Topical application; 7.1% (w/w) ethanolic extract; Wistar albino rats of either sex (150–200 g):	EWM, IWM; 5% (w/w) PVI; Ethanolic extract (Lodhi and Singhi, 2011)	Leaves; Methanolic; Wistar albino rats (150–200 g)	AOTS (2000 mg/kg bw); NT/NM (Babu et al., 2010)	DPPH scavenging of n-butanol soluble fraction of stem and leaves were 83.62% and 82.88% respectively at concentration of 250 µg/ml. n-butanol soluble fraction of stem exhibited highest TAA (0.187) and highest FRAP value of 149.00 mg/TE (Aziz-ur-rehman et al., 2012). Methanol extract of leaves showed antioxidant properties in RP, DPPH radical scavenging activity, nitric oxide scavenging activity, H ₂ O ₂ radical scavenging activity, superoxide radical scavenging assay, hydroxyl radical scavenging activity and total antioxidant method (Nagda et al., 2009).	NE
<i>Millettia extensa</i> (Benth.) Baker	NE	NE	NE	NE	NE	NE
<i>Mirabilis jalapa</i> L.	NE	NE	Flowers; Hydroethanolic; Female rat	AOTS (2000 mg/kg bw); NT/NM (Augustine et al., 2013)	Crude methanol extract of the bark showed IC ₅₀ value of 598.02 µg/ml in DPPH assay (Rumzhum et al., 2008). EC ₅₀ values for ABTS free radical scavenging activity were 1249 µg/ml and 974 µg/ml for methanolic root and aerial extracts respectively, EC ₅₀ values for DPPH radical scavenging activity were 1679 µg/ml and 3723 µg/ml for methanolic root and aerial extracts respectively (Zachariah et al., 2011).	NE
<i>Mitragyna parvifolia</i> (Roxb.) Korth.	NE	NE	Leaves; Ethanolic; Swiss albino mice (25–30 g)	AOTS (5, 100, 300 and 1500 mg/kg bw); NT/NM (Kaushik et al., 2009a)	DPPH activity of ethanolic leaf extract was 87.6% at a concentration of 500 µg/ml, whereas percentage inhibition of superoxide generation at 500 µg/ml concentration was 65.0% in superoxide radical scavenging assay (Kaushik et al., 2009c).	NE
<i>Momordica dioica</i> Roxb. ex Willd.	NE	NE	Fruits; Aqueous; Wistar rats (150–200 g)	AOTS (5, 10 and 15 times of effective dose (ED) i.e., 200 mg/kg bw; NT/NM even with 15 times of ED (R. Singh et al., 2011)	Ethanolic extract of root scavenge the super oxide generated by ascorbic system at the dose of 12.5 µg/ml, whereas at concentration of 400 µg/ml DPPH radical scavenging activity was recorded as 77.30% (Shreedhara and Vaidya, 2006). <i>In vivo</i> antioxidant and free radical scavenging activities were positive for both ethanolic and aqueous extracts of leaves (Jain et al., 2008). Hydroethanolic extract of leaves at 200 and 400 mg/kg concentration	NE

<i>Mucuna pruriens</i> (L.) DC.	Seeds; Topical application; 1% and 2% methanolic extract; Swiss albino mice (20–25 g);	EWM, IWM; FRA; 2% methanolic extract (Gunde et al., 2013)	Seeds; Methanol; Albino mice of both sexes	AOTS (150, 300, 500, 1000, 2000, 3000 and 4000 mg/kg bw); NT/NM (Manalisha and Chandra, 2012)	significantly reversed the diminution in the level of <i>in vivo</i> antioxidants in rats (Obogwu et al., 2014). <i>In vitro</i> study of seed extract possesses dose dependent protection against superoxide generation, hydroxyl radical production and FeSO ₄ induced lipid peroxidation in normal albino rats (100–150 g bw) of Charles Foster strain (Tripathi and Upadhyay, 2001).	
<i>Nicotiana glauca</i> L.	NE	NE	NE	NE	NE	NE
<i>Oroxylum indicum</i> (L.) Kurz	Root; Topical application; 1% and 2.5% methanolic extract; Albino mice (NA)	BWM; SSD, CHG; Methanolic extract (H. Singh et al., 2011)	Roots; Ethanolic; Wistar male albino rats (150–200 g)	AOTS (5000 mg/kg bw); NT/NM (Tamboli et al., 2011)	IC ₅₀ values in DPPH radical scavenging assay of root extract, root bark extract, stem extract, stem extract, leaf extract and fruit extract was recorded as 158.23, 112.21, 189.36, 149.59, 106.4 and 159.46 µg/ml respectively, Nitric oxide radical scavenging assay of root extract, root bark extract, stem extract, stem extract, leaf extract and fruit extract was recorded as 102.56, 89.36, 88.15, 139.90, 72.05 and 182.25 µg/ml respectively, Superoxide radical scavenging assay of root extract, root bark extract, stem extract, stem extract, leaf extract and fruit extract was recorded as 455.36, 154.96, 681.22, 137.30, 170.87 and 399.86 µg/ml respectively, Hydroxy radical scavenging assay of root extract, root bark extract, stem extract, stem extract, leaf extract and fruit extract was recorded as 179.35, 47.01, 309.27, 33.30, 46.52 and 76.39 µg/ml respectively (Mishra et al., 2010).	NE
<i>Persicaria barbata</i> (L.) H.Hara	NE	NE	Leaves; Alcoholic; Wistar albino rats	AOTS (5, 50, 300 and 2000 mg/kg bw); NT/NM safe up to dose of 2000 mg/kg bw (Sheela et al., 2011)	NE	NE
<i>Phyllanthus amarus</i> Schumacher and Thonn.	Whole plant; Oral application; 200 mg/kg alcoholic extract; Male Wistar albino rats (150–200 g)	EWM, IWM, DSW; 0.17 mg/kg DXM; Alcoholic extract (Devi et al., 2005)	Leaves; Aqueous; Female rats (150–200 g)	AOTS (2000 and 5000 mg/kg bw); NT/NM safe up to dose of 5000 mg/kg bw (Asare et al., 2011)	Aqueous (IC ₅₀ =52.14 µg/ml), methanolic (IC ₅₀ =63.50 µg/ml), chloroform (IC ₅₀ =367.28 µg/ml) and hexane (IC ₅₀ =454.34 µg/ml) extracts of whole plant showed DPPH radical scavenging activity, aqueous (IC ₅₀ =12.58 in µg) and methanolic (IC ₅₀ =16.56 in µg) extracts of whole plants showed activity in ABTS assay, aqueous (IC ₅₀ =392.37 in µg) and methanolic (IC ₅₀ =448.81 in µg) extracts of whole plants showed activity in FRAP assay (Guha et al., 2011).	NE
<i>Pongamia pinnata</i> (L.) Pierre	Leaves; Topical application; 200 mg/kg methanolic extract; Wistar albino rats of either sex (180–230 g)	EWM, IWM; PIO; Methanolic extract (Prasad et al., 2011)	Seed; Crude extracts; Wistar female albino rats (100–120 g)	AOTS (2000 mg/kg bw); NT/NM (Aneela et al., 2011)	Chloroform and butanol extracts of seeds at conc. of 500 µg significantly inhibited the generation of hydroxyl radical and superoxide anions as well as the reaction of lipid peroxidation in rat liver microsomes induced by Fe ²⁺ (Bhatia et al., 2008). <i>In-vitro</i> antioxidant activity of cycloart-23-ene-3β, 25-diol isolated from stem	NE

Table 3 (continued)

Plant species	Wound healing studies		Toxicological studies		Antioxidant activities	Tyrosinase inhibitory activities
	Plant part; Application; Extract; Animal (Weight)	Animal Model; Reference control; Effective treatment [Source for columns 2 & 3]	Plant part; Extract; Animal (Weight)	Animal model (Dose); Toxicity status [Source for columns 4 & 5]		
<i>Premna mollissima</i> Roth	NE	NE	Leaves; Methanolic; Albino mice of either sex	AOTS (100, 500, 1000, 3000 and 5000 mg/kg bw); NT/NM (Mahire et al., 2009)	bark showed dose dependent% reduction against DPPH radical, reducing power, superoxide anion radical scavenging, hydroxyl radical scavenging, metal chelating, hydrogen peroxide scavenging and nitric oxide radical scavenging by B2 and β -tocopherol (Badole et al., 2011). NE	NE
<i>Ranunculus sceleratus</i> L.	NE	NE	NE	NE	NE	NE
<i>Rauvolfia serpentina</i> (L.) Benth. ex Kurz	NE	NE	Root; Methanolic; Male Wistar albino mice (20–30 g)	AOTS (10, 30, 60, 100, 150, 200 and 250 mg/kg bw); Doses at 150, 200 and 250 mg/kg bw showed acute toxicity, therefore, LD ₅₀ was determined as 141.25 mg/kg bw (Azmi and Qureshi, 2012)	NE	NE
<i>Ricinus communis</i> L.	NE	NE	Root; Aqueous and methanolic; Wistar albino rats	AOTS (2000 mg/kg bw), SATS (1000 mg/kg bw); NT/NM (Ilavarasan et al., 2011)	IC ₅₀ values for the leaves extracts ranged from 0.65 to 2.39 μ g/ml in DPPH radical scavenging assay (Wafa et al., 2014). Aqueous methanol extract of dry leaves had IC ₅₀ value of 2.70 μ g/ml in DPPH radical scavenging assay (P.P. Singh et al., 2009).	NE
<i>Scoparia dulcis</i> L.	Roots; Topical application; 5 and 10% (w/w) ethanolic extract; Wistar albino rats of either sex (180–200 g)	EWM, IWM, DSW; 0.2% (w/w) NFZ; 10% Ethanolic extract (Murti et al., 2011)	Whole plant; Methanolic; Mice	AOTS (10, 100, 1000, 1600, 2900 and 5000 mg/kg bw); NT/NM, LD ₅₀ 3807 mg/kg bw (Abdulsalaam et al., 2013)	Freeze dried aqueous extract at 0.01, 0.1 and 1.0 mg/ml concentration showed antioxidant index of 26.92, 34.95 and 48.26, respectively and EC ₅₀ value was 1.08 mg/ml (Ratnasooriya et al., 2005).	NE
<i>Semecarpus anacardium</i> L.f.	NE	NE	Nuts; <i>Serankottai nei</i> -a Siddha preparation by milk extracts of nuts; Wistar strain male albino rats (80–100 g)	AOTS (75–2000 mg/kg bw), SATS (50, 100, 250 and 500 mg/kg bw); NT/NM, dose of 500 mg/kg bw in SATS showed moderate changes (Vijayalakshmi et al., 2000)	DPPH radical scavenger activity of hexane, chloroform, ethyl acetate and methanol extract of stem bark was reported as 103.69, 82.45, 44.03 and 60.23 μ g/ml respectively, Lipid peroxidation activity of hexane, chloroform, ethyl acetate and methanol extract of stem bark was reported as 205.08, 187.43, 102.34 and 165.21 μ g/ml respectively, Nitric oxide radical scavenger activity of hexane, chloroform, ethyl acetate and methanol extract of stem bark was reported as 176.33, 132.43, 80.75 and 119.23 μ g/ml respectively, Superoxide radical scavenger activity of hexane, chloroform, ethyl acetate and methanol extract of stem bark was reported as 89.91, 73.23, 68.55 and 78.21 μ g/ml respectively (Sahoo et al., 2008).	NE

<i>Senna tora</i> (L.) Roxb.	Leaves; Topical application; 5% (w/w) ethanolic extract; Wistar albino rats (150–180 g)	EWM; 0.2% (w/w) NFZ; Ethanolic extract (Jayasutha and Nithila, 2011)	Leaves; Ethanolic; Swiss albino mice of both sexes	AOTS (100, 1000, 10000 mg/kg bw), SATS (500, 1000 mg/kg/day bw); NT/NM (Ambali et al., 2005)	EC ₅₀ values of ethyl acetate, n-butanol and water fraction was recorded as 70.1, 578.0 and 770.0 µg/ml respectively in DPPH assay, whereas ferrous ions chelating activity of ethyl acetate, n-butanol and water fraction was recorded as 70.57%, 13.60% and 30.40% at 0.5 mg/ml concentration, respectively (Zhenbao et al., 2007).	NE
<i>Sesamum indicum</i> L.	NE	NE	Seed oil; direct; Swiss albino mice (25–30 g)	AOTS (oil oral doses of 0.5, 1, 1.5, 2 and 3 g/kg bw); NT/NM (Monteiro et al., 2014)	Sesamol from seed showed IC ₅₀ value of 5.44 µg/ml in DPPH free radical assay (C. M. Kumar et al., 2011)	Sesamol from seed showed tyrosinase inhibition with IC ₅₀ value of 3.2 µM in mushroom tyrosinase assay (C.M. Kumar et al., 2011).
<i>Shorea robusta</i> Gaertn.	Resin; Topical application; 10 and 30% ethanolic extract; Wistar albino rats (150–250 g), Swiss mice (18–20 g)	EWM, IWM; 1% (w/w) FRA; Both (Wani et al., 2012a)	Leaves; Aqueous and methanol; Swiss albino mice (18–20 g) and male wistar rats (150–180 g)	AOTS (0–2500 mg/kg orally (P.O.) and 0–1000 mg/kg (i.p.)); NT/NM with LD ₅₀ of orally fed methanolic and aqueous extract of 2.4 g/kg and 2.7 g/kg bw, while it was 1.2 g/kg and 1.4 g/kg i.p. respectively (Chattopadhyay et al., 2012)	NE	NE
<i>Sida cordata</i> (Burm. F.) Borss. Waalk.	NE	NE	NE	NE	Leaf extract reduced the lipid peroxidation in the liver tissue and restored activities of defense antioxidant enzymes superoxide dismutase (SOD), lipid peroxidation, catalase (CAT), reduced glutathione (GSH) towards normal levels in Wister albino rats (Mistry et al., 2013).	NE
<i>Sida rhombifolia</i> L.	NE	NE	Root; Aqueous; Male and female Sprague-Dawley rats (130–190 g)	AOTS (5,000 mg/kg bw), SATS (300, 600 and 1200 mg/ml bw); NT/NM (Sireeratawong et al., 2008)	Ethanol extracts of root, leaves, whole plant and stem showed 50% inhibitory concentrations of 546.1, 852.8, 983.8, and 1,222.5 µg/ml, respectively in DPPH assay (Dhalwal et al., 2007).	NE
<i>Solanum incanum</i> L.	NE	NE	Fruit; Ethanolic; Female Swiss albino mice (25–30 g)	AOTS (100, 250, 500, 750, 1000 and 2000 mg/kg bw); NT/NM (Indhumathi et al., 2014)	NE	NE
<i>Terminalia arjuna</i> (Roxb. ex DC.) Wight & Arn.	Bark; Topical application; hydroalcoholic extract; Rats (NA)	EWM, IWM; NA; NA (Chaudhari and Menqi, 2006)	Leaves; Methanolic; Adult male Swiss albino mice (18–22 g)	AOTS (900 mg/kg bw), SATS (200 mg/kg bw for 28 days); NT/NM with LD ₅₀ 900 mg/kg bw (Biswas et al., 2011a)	Reducing power of bark extracts at 10 mg/ml concentration were 1.340, 1.600 and 1.740 for 80% ethanol, 80% methanol and 80% acetone extracts respectively (Sultana et al., 2007).	NE
<i>Tridax procumbens</i> (L.) L.	Whole plant; Topical application; aqueous extract, butanol fraction and petroleum ether fraction; Albino rats (NA)	EWM, IWM; NA; NA (Udupa et al., 1995)	Leaves; Ethyl acetate; Swiss albino mice of both sexes each (20–30 g) and albino rats (50–80 g)	AOTS (10, 100, 1000, 1600, 2900 and 5000 mg/kg bw), Short term toxicity study (50, 100, 200, 400 and 800 mg/kg/day bw for 14 days); LD ₅₀ of the extract was 2100 mg/kg bw (Abubakar et al., 2012)	Ethyl acetate and methanol extract of aerial parts showed IC ₅₀ values of 45.95 and 35.62 µg/ml, respectively in DPPH assay, whereas IC ₅₀ values of 24.47 and 18.17 µg/ml were reported for ethyl acetate and methanol extracts of aerial parts in ABTS assay (Jachak et al., 2011).	NE
<i>Typha domingensis</i> Pers.	NE	NE	NE	NE	Superoxide scavenging half-maximal effective concentration (EC ₅₀) of fruit, female flower and male flower extracts was 3.5, 4.8, and 28.2 mg dry matter (DM)/ml, respectively, while nitric oxide scavenging EC ₅₀ of fruit, female flower and male flower extracts was 0.16, 0.65, and 0.95 mg DM/ml, respectively. Iron chelating EC ₅₀ of female flower, male flower and fruit extracts was 4.86, 6.43, and 10.88 mg DM/ml, respectively (Chai et al., 2014).	NE
	NE	NE				NE

Table 3 (continued)

Plant species	Wound healing studies		Toxicological studies		Antioxidant activities	Tyrosinase inhibitory activities
	Plant part; Application; Extract; Animal (Weight)	Animal Model; Reference control; Effective treatment [Source for columns 2 & 3]	Plant part; Extract; Animal (Weight)	Animal model (Dose); Toxicity status [Source for columns 4 & 5]		
<i>Vallisneria spiralis</i> (Roth) Kuntze			Root bark oil; Ethanolic; Albino rats (150–200 g)	Toxicity study (3 and 5 ml/kg bw); NT/NM (Punam et al., 2012)	Ethanol extract in qualitative test of DPPH free radical assay showed antioxidant activity (Karmakar et al., 2011).	
<i>Vanda tessellata</i> (Roxb.) Hook. ex G.Don.	Whole Plant; Topical application; NA; Rats(NA)	EWM; NA (Nayak et al., 2005)	Whole plant with flowers; Water and alcoholic; Swiss mice (25–35 g)	AOTS (0, 0.5, 1.0 and 2 g/kg bw), SATS (200 and 400 mg/kg bw); NT/NM (Kumar et al., 2000)	NE	NE
<i>Verbascum thapsus</i> L.	Flowers; Topical application; 20% chloroform methanol extract; Male rabbits (1500 g)	PIW; 20% ZNO; 20% Chloroform methanol extract (Mehdinezhad et al., 2012)	Leaves; Aqueous	Brine Shrimp and radish seed bioassays Toxic at higher doses(around 1000 mg/L) with LC ₅₀ of < 1000 mg/L (Turker and Camper, 2002)	Aqueous extract of flower showed total antioxidant capacity of 195 µmol Trolox equivalent/g dry wt in a two-stage Trolox based assay (Vanderjagt et al., 2002).	NE
<i>Vitex negundo</i> L.	Leaves; Topical application; 200 mg/kg/day aqueous and ethanolic extracts; Wistar rats of both sexes (150–200 g)	EWM, IWM; 5% PVI; Both extracts (Talekar et al., 2012)	Oil; direct; Wistar rats of both sexes	Acute dermal toxicity study (2000 mg/kg bw), Sub-chronic dermal toxicity study (250, 500 and 1000 mg/kg bw); NT/NM, LD ₅₀ value was over 2000 mg/kg bw (Chattopadhyay et al., 2014)	Aqueous (IC ₅₀ =80.45 µg/ml), Methanolic (IC ₅₀ =93.73 µg/ml), Chloroform (IC ₅₀ =190.58 µg/ml) and hexane (IC ₅₀ =913.76 µg/ml) extracts of whole plant showed scavenging activity in DPPH assay, Aqueous (IC ₅₀ =13.01 in µg) and Methanolic (IC ₅₀ =23.25 in µg) extracts of whole plants showed antioxidant activity in ABTS assay, whereas aqueous (IC ₅₀ =1021.38 in µg) and methanolic (IC ₅₀ =476.91 in µg) extracts of whole plant showed activity in FRAP assay (Guha et al., 2011).	Tyrosinase inhibition activity by (+)-lyoniresinol from roots showed inhibitory activity with IC ₅₀ value of 3.21 µM (Azhar-ul-Haq et al., 2006).
<i>Wrightia arborea</i> (Dennst.) Mabb.	Leaves; Topical application; NA; Wistar albino Rats of either sex (150–200 g)	EWM, IWM; 0.2% NFZ; NA (Devi and Divakar, 2013)	Leaves; Methanolic; Male albino rats (120–140 g)	AOTS (2000 mg/kg bw); NT/NM (Nahar et al., 2013)	Free radical scavenging activity of ethanolic bark extract has shown ability to suppress the oxidation (D.L. Kumar et al., 2011).	NE
<i>Ziziphus nummularia</i> (Burm. f.) Wight & Arn.	Leaves; Topical application; 20 and 30% ethanolic extracts; Wistar albino rats (200 g)	EWM; 10% (w/w) PIO; 30% ethanolic extract (Yusufoglu, 2011)	Leaves; Cyclopeptide alkaloid fraction; Wistar rats of both sexes (180–200 g) and Swiss mice (20–24 g)	AOTS (50, 200 and 300 mg/kg bw); Category-3 having LD ₅₀ 200 mg/kg bw, CNS depression and reduction in locomotor activity in mice treated with 200 and 300 mg/kg bw (Goyal et al., 2013)	Aqueous ethanol extract of peel (200.22 to 40.26 AEAC/100 g DW), pulp (51.30 to 28.56 AEAC/100 g DW) and seed (50.06 to 22.44 AEAC/100 g DW) showed scavenging activity in DPPH assay, whereas, aqueous ethanol extract of peel (372.49 to 3852.88 AEAC/100 g DW), pulp (982.31 to 252.43 AEAC/100 g DW) and seed (639.05 to 219.94 AEAC/100 g DW) also showed activity in FRAP assay (Zhang et al., 2010).	NE

Abbreviations: ABTS=2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) assay, AOTS=acute oral toxicity study, BWM=burn wound model, CHG=chlorhexidine gluconate, CTS=chronic toxicity study, DPPH=2,2-diphenyl-1-picrylhydrazyl assay, DSW=dead space wound model, DXM=dexamethasone, EWM=excision wound model, FRA=framycetin, FRAP=ferric reducing antioxidant powder assay, FSC=framycetine sulfate cream, IWM=incision wound model, NA=not available, NE=not evaluated, NEO=neospirin, NFZ=nitrofurazone, NT/NM=non-toxic/no mortality, PIW=punch incision wound model, PVI=povidone iodine ointment, RP=reducing power, SATS=sub acute toxicity study, SOF=soframycin, SSD=silver sulphadiazine, ST=sulphathiazole, TAC=total antioxidant capacity, ZNO=zinc oxide.

Table 4

Earlier studies showing anti-inflammatory and antimicrobial activities, and active compounds in the plants reported in the present study.

Plant species	Antiinflammatory studies		Microbiological studies		Active compound [Plant part]; Biological activity related to skin problems (DNP, 2014)
	Plant part; Extract; Animal (Weight)	Animal Model (Dose); Reference control; Activity [Source for columns 2 & 3]	Plant part tested; Solvent or extracts used: Tested against (micro-organism)	Effective extract; Most effective against; Phytoconstituents [Source for columns 4 and 5]	
<i>Acalypha indica</i> L.	Whole plant; Methanolic; Swiss albino mice (20–25 g) and Long Evans rats (140–160 g) of either sex	CPE (125 and 250 mg/kg bw); Reference drug, Phenylbutazone (100 mg/kg bw); AIA (Rahman et al., 2010)	Leaves; Water, ethanol and chloroform; Bacteria; <i>Escherichia coli</i> , <i>Salmonella enteritidis</i> , <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> and fungi; <i>Candida albicans</i> , <i>Candida tropicalis</i> , <i>Microsporum canis</i> , <i>Aspergillus fumigatus</i> Whole plant; Ethyl acetate, methanol, Ethanol and diethyl ether; <i>Bacillus subtilis</i> , <i>Escherichia coli</i> , <i>Enterococcus faecalis</i> , <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i> , <i>Staphylococcus epidermis</i> , <i>Klebsiella pneumonia</i> , <i>Proteus vulgaris</i>	Water and ethanol for gram positive bacteria, Antifungal activity in chloroform extract; NA (Somchit et al., 2010) Ethyl acetate; Most of the bacterial strains; Tannins, steroids, saponins, cardiac glycosides, terepens, alkaloids (Poornima and Prabakaran, 2012)	NE
<i>Achyranthes aspera</i> L.	Roots; Alcoholic; Wistar rats Leaves; Topical application of methanolic extract; Male Wistar rats	CPE, CPG (50,100 and 200 mg/kg bw); NA; AIA in both models (S.V. Kumar et al., 2009) CPE; NA; Weak AIA (Khuda et al., 2013)	Roots; Ethyl acetate, acetone, ethanol; <i>Bacillus subtilis</i> , <i>Streptococcus aureus</i> , <i>Salmonella typhimurium</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Salmonella abony</i>	Ethanol; Most of the bacterial strains; NA (Shendkar et al., 2012)	17-Pentatriacontanol [shoot]; Antifungal agent.
<i>Acorus calamus</i> L.	Leaves; Aqueous; On human keratinocyte HaCaT cells.	PPCIN, PGNIN; NA; AIA (Kim et al., 2009)	Rhizome; Ethanol; Bacteria: <i>Pseudomonas sp.</i> , <i>Bacillus sp.</i> , <i>Staphylococcus aureus</i> and fungus: <i>Aspergillus niger</i> , <i>Aspergillus flavus</i> , <i>Trichoderma sp.</i>	Ethanol; <i>Pseudomonas sp.</i> , <i>Staphylococcus aureus</i> , <i>Aspergillus flavus</i> ; Aromatic compounds, ketone, alkaloid, fatty acids like palmitic acid and linoleic acid (Senthilkumar et al., 2010)	4-Acoren-3-one (multi drug resistance MDR inhibitory activity) and 2,5-Dihydroxy-1,4-benzoquinone (mild activity against gram-positive bacteria and mycobacteria); Antibacterial agents. 1-(4-Hydroxyphenyl)-2-propen-1-ol; Superoxide anion generation inhibitor. Inhibits tumor promotion by TPA JWJ26 on ICR mouse skin.
<i>Ageratum conyzoides</i> (L.) L.	Whole plants; Alcoholic; Swiss albino mice (25–30 g) and Wister albino rats of the both sexes (150–200 g)	CPE; Diclofenac sodium (40 mg/kg bw); NA; AIA (M.A. Rahman et al., 2012)	Leaves; methanol, ethanol, distilled water, hexane, ethyl acetate and acetone; <i>Staphylococcus</i> and <i>Escherichia coli</i>	Methanol; <i>Staphylococcus</i> and <i>Escherichia coli</i> ; Alkaloids, tannins, saponins, glycoside, flavonoids, resins and anthraquinone (Onuoha et al., 2013)	NE
<i>Albizia lebbek</i> (L.) Benth.	Bark; Petroleum ether, ethyl acetate and methanolic	CPE (400 mg/kg bw); NA; AIA (Saha and Ahmed, 2009)	Leaves; Petroleum ether, ethyl acetate, methanol; <i>Bacillus subtilis</i> , <i>Escherichia coli</i> , <i>Klebsiella pneumonia</i> , <i>Proteus vulgaris</i> , <i>Pseudomonas aeruginosa</i> , <i>Salmonella typhi</i> , <i>Staphylococcus aureus</i> . Leaves; ethyl acetate; <i>Escherichia coli</i> , <i>Staphylococcus aureus</i> , <i>Pseudomonas aeruginosa</i> , <i>Bacillus cereus</i>	Methanol; <i>Salmonella typhi</i> , ethyl acetate; <i>Escherichia coli</i> ; Flavonoids, tannins, saponins (Bobby et al., 2012) Ethyl acetate; <i>Pseudomonas aeruginosa</i> ; Alkaloids, glycoside, tannins, saponins, flavanoids, carbohydrates (Chulet et al., 2010)	NE
<i>Allium cepa</i> L.	Bulbs; Male albino rats (125–150 g)	CPE; HPE; Reference drug, Diclofenac (5 mg/kg bw); AIA (Riyaz et al., 2012)	Onion oil; 4 gram positive, 4 gram negative and 9 fungal strains	Against all the tested microorganisms; NA (Zohri et al., 1995)	Di-2-propenyl disulfide, protein 1 [seed] and allicepin [bulb]; Antifungal agents. Protein 1 [seed]; Antibacterial agents. Antimicrobial protein 1 in seeds; Antimicrobial agent. Alliucide G, 3-Mercapto-2-methylpentanol, <i>Allium</i> Quercetin trimer, Quercetin-protocatechuic acid heterodimer and <i>Allium cepa</i> Antimicrobial protein 1; Antioxidants.
<i>Amaranthus spinosus</i> L.		CPE (250, 500 and 750 mg/kg bw); Reference drug,	Leaves; hexane, ethyl acetate, dichloromethane and methanol; Grampositive: <i>Staphylococcus</i>		NE

Table 4 (continued)

Plant species	Antiinflammatory studies		Microbiological studies		Active compound [Plant part]; Biological activity related to skin problems (DNP, 2014)
	Plant part; Extract; Animal (Weight)	Animal Model (Dose); Reference control; Activity [Source for columns 2 & 3]	Plant part tested; Solvent or extracts used: Tested against (micro-organism)	Effective extract; Most effective against; Phytoconstituents [Source for columns 4 and 5]	
	Leaves; Petroleum ether and ethanolic; Male or female albino rats (140–160 g)	Ibuprofen (10 mg/kg bw); AIA (Baral et al., 2010)	<i>aureus</i> and <i>Bacillus</i> spp, gram-negative <i>Escherichia coli</i> , <i>Salmonella typhi</i> , <i>Pseudomonas aeruginosa</i> , <i>Proteus mirabilis</i> and <i>Klebsiella pneumoniae</i> and a pathogenic fungus <i>Candida albicans</i>	<i>Salmonella typhi</i> ; flavonoids, steroids, terpenoids, saponins and cardiac glycosides (Maiyo et al., 2010)	
<i>Anisomeles indica</i> (L.) Kuntze	Leaves and stem; Freeze-dried decoction; Rats	CPE, FoPE, APE (125, 250 and 500 mg/kg bw); NA; AIA (Dharmasiri et al., 2002)	Leaves; aqueous, methanolic, ethanolic, Chloroform and ethyl Acetate; <i>Escherichia coli</i> , <i>Staphylococcus aureus</i> , <i>Proteus vulgaris</i> , <i>Pseudomonas aeruginosa</i> and <i>Klebsiella pneumonia</i>	Chloroform; <i>Pseudomonas aeruginosa</i> ; NA (Dixit, 2013)	3,7,11,15(17)-Cembratetraene-16,2:19,6-diolide; Induces cell cycle arrest and apoptosis in human oral squamous cell carcinoma Ca9-22 cells.
<i>Annona squamosa</i> L.	Bark; Caryophyllene oxide isolated from an unsaponified petroleum ether extract; Wistar albino rats of either sex	Caryophyllene oxide (12.5 and 25 mg/kg bw); NA; AIA (Chavan et al., 2010)	Seeds, methanol, petroleum ether, and chloroform; Gram-positive (<i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i>) and Gram-negative (<i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> and <i>Klebsiella pneumoniae</i>)	Methanol, Chloroform: All the test organisms; NA (Aher et al., 2012)	3-(14,17:18,21-Diepoxydotriacontyl)-5-methyl-2(5 H)-furanone; Shows antitumor activity <i>in vitro</i> . 3-(12,15:16,19-Diepoxydotriacontyl)-5-methyl-2(5 H)-furanone; Cytotoxic agent showing activity against human prostate tumour cells. 3-(14,17:18,21-Diepoxydotriacontyl)-5-methyl-2(5 H)-furanone and Parviflorin, Bullatacinone, Mosinone A, Squamostanin A, Annosquatin I, Rollicosin and 2-(18,21-Epoxydotriacontyl)-2,5-dihydro-5-methyl-2(5 H)-furanone; Cytotoxic agent.
<i>Argemone mexicana</i> L.	Leaves; Lyophilized aqueous extract; Male and female white mice (rain NMRI) (25–30 g)	CPE (250 and 500 mg/kg bw); Reference drug, Paracetamol (100 mg/kg bw); Significant AIA (Sourabie et al., 2012)	Leaves, seeds; Crude methanol, aqueous, n hexane, chloroform, methanol; <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Enterococcus</i> sp., <i>Salmonella typhi</i> .	Methanol, Chloroform: All the test organisms; Alkaloids, amino acids, phenolics, fatty acids (Priya and Rao, 2012)	Sanguinarine; Toxic to animals, potent cytotoxic, antifungal, antibacterial and antimicrobial agent.
<i>Azadirachta indica</i> A. Juss.	Roots; Alcoholic; Wistar albino rats of either sex	CPE, CPG (200, 400 and 800 mg/kg bw); Reference drug, Aspirin (100 mg/kg bw); AIA (Patil et al., 2012)	Leaf, Stem; Acetone, ethyl acetate, petroleum ether; <i>Escherichia coli</i> , <i>Shigella</i> sp., <i>Staphylococcus</i> sp., <i>Salmonella</i> sp. Leaves; Aqueous; <i>Escherichia coli</i> and <i>Salmonella</i> sp.	Petroleum ether; <i>Staphylococcus</i> sp., <i>Escherichia coli</i> ; NA (Rahman et al., 2011) Aqueous ; Against all the tested organisms; Alkaloids, steroids, tannin (Susmitha et al., 2013)	Isorhamnetin 3,7-diglycosides; Antioxidant. 10-Undecyn-1-ol [leaf]; Antifungal agent.
			Leaves; Methanol; <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Klebsiella pneumonia</i> , <i>Streptococcus pyrogens</i> , <i>Staphylococcus aureus</i>	Methanol; <i>Pseudomonas aeruginosa</i> ; Reducing sugar, glycosides, tannins, triterpenes (Vasantharaj et al., 2013)	Nimbolide, marnoodin, α -Nimolactone, β -nimolactone and tetrahydro-3,4-furandiol; Antibacterial agents. Nimbolide; Cytotoxic towards human tumor cells. 7-Hydroxy-1-meliacene-3,15-dione; Shows antitumour activity. NE
<i>Bambusa bambos</i> (L.) Voss	Leaves; Methanolic; Albino rats	CPE, IPE; NA; Significant AIA (Muniappan and Sundararaj, 2003)	Leaves; hexane, ethyl acetate, ethanol and hydroalcohol; <i>Escherichia coli</i> , <i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> , <i>Klebsiella pneumonia</i> <i>Pseudomonas aeruginosa</i>	Hydroalcohol as well as ethanol; <i>Escherichia coli</i> ; NA (Sandhiya et al., 2013)	
<i>Bauhinia variegata</i> L.	Bark; Methanolic and aqueous; Albino rats (150–200 g)	CPE, DPE (200 and 250 mg/kg bw); NA; AIA (Bairagi et al., 2012)	Stem bark; Methanol, aqueous, ethanol; <i>Bacillus cereus</i> , <i>Staphylococcus aureus</i> , <i>Klebsiella pneumonia</i> , <i>Escherichia coli</i> , <i>Pseudomonas pseudoalcaligenes</i> , <i>Bacillus subtilis</i> , <i>Pseudomonas</i>	Methanol, ethanol; Gram positive bacteria; Quercitroside, isoquercitroside, rutoside, myricetol glycoside, kaempferol glycoside (Sahu and Gupta, 2012)	3',4',5,7-Tetrahydroxy-3-methoxyflavone; Anti-inflammatory agent.

<i>Boerhavia diffusa</i> L.	Leaves; Ethanolic; Mice	CPE, SPE, HPE, DPE (400 mg/kg bw); NA; AIA (Mahesh et al., 2012)	<i>aeruginosa</i> , <i>Salmonella typhi</i> , <i>Shigella dysenteriae</i> , <i>Vibrio cholera</i>	Ethanol, chloroform, aqueous; More activity towards human pathogenic organisms except <i>Vibrio cholerae</i> ; Quinones, saponins, triterpenoids, flavonoids, alkaloids, steroids, tannins, furanoids, phenols (Umamaheswari et al., 2010)	NE
<i>Brassica juncea</i> (L.) Czern.	Seeds; Petroleum ether and ethanolic; Male Wistar albino rats (150–200 g)	CPE (250 and 500 mg/kg bw); Reference drug, Ibuprofen (10 mg/kg bw); AIA (Sindhoor et al., 2012)	Seed meal; Crude extract and purified phenolic fractions; <i>Staphylococcus aureus</i> , <i>Listeria monocytogenes</i> , <i>Lactobacillus plantarum</i> , <i>Bacillus subtilis</i> , <i>Escherichia coli</i> , <i>Pseudomonas fluorescens</i>	Crude extract and purified phenolic fraction; <i>Staphylococcus aureus</i> , <i>Listeria monocytogenes</i> (Engels et al., 2012)	Cyclobrassinin sulfoxide and brassilexin [leaf]; Phytoalexin.
<i>Buchanania cochinchinensis</i> (Lour.) M. R.Almeida	Kernels; Methanolic; Wistar rats (150–200 g)	CPE (100 and 200 mg/kg bw); Reference drug, Indomethacin (10 mg/kg bw); AIA (Warokar et al., 2010)	Bark; Methanol; Bacteria (<i>Bacillus cereus</i> , <i>Escherichia coli</i> , <i>Proteus vulgaris</i> , <i>Pseudomonas aeruginosa</i> , <i>Bacillus subtilis</i> , <i>Salmonella</i> sp., <i>Staphylococcus aureus</i>) and fungal species (<i>Aspergillus niger</i> , <i>penicillium</i> sp. and <i>Trichoderma viride</i>)	Methanol; <i>Escherichia coli</i> , <i>Staphylococcus aureus</i> ; saponins, flavanoids, steroids, cardiac glycosides, alkaloids, tannins and phenolics (Venkata et al., 2012)	Isorhamnetin 3,7-diglycosides; Antioxidant. NE
<i>Butea monosperma</i> (Lam.) Taub.	Flowers; Ethanolic; Male Wistar albino rats of either sex (150–200 g)	CPE; Reference drug, Diclofenac gel (0.2 g 1%); AIA (Sapkale et al., 2013)	Leaves; Ethanol, aqueous; <i>Staphylococcus aureus</i> , <i>Escherichia coli</i>	Ethanol; both; Alkaloids, carbohydrates, tannins, flavonoids, phenolic compounds, starch (Ramanjaneyulu et al., 2011)	5,7-Dihydroxy-3,4',6-trimethoxyflavone [flower]; Antifungal agent.
<i>Caesulia axillaris</i> Roxb.	NE	NE	Essential oil; <i>Trichophyton tubrum</i> and <i>Microsporum gypseum</i>	Essential oil; <i>Trichophyton tubrum</i> and <i>Microsporum gypseum</i> (Kishore et al., 1993)	NE
<i>Callicarpa macrophylla</i> Vahl	Leaves; Ethanolic and aqueous; Male and female albino rats (120–150 g)	CPE (200 and 400 mg/kg bw); Reference drug, Diclofenac sodium (20 mg/kg bw); AIA (Yadav et al., 2011)	Stem; Ethanol, aqueous; Gram +ve <i>Streptococcus pyogenes</i> , <i>Bacillus</i> , <i>Micrococcus luteus</i> , <i>Staphylococcus epidermidis</i> , <i>Clostridium sporogens</i> , <i>Streptococcus faecalis</i> , <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> and Gram –ve <i>Agrobacterium tumefaciens</i> , <i>Klebsiella pneumonia</i> , <i>Salmonella typhimurium</i> , <i>Pseudomonas aeruginosa</i> , <i>Serratia marcescens</i> , <i>Enterobacteria aerogens</i> , <i>Proteus vulgaris</i> , <i>Escherichia coli</i>	Ethanol; <i>Bacillus subtilis</i> ; NA (V. Yadav et al., 2012)	NE
<i>Calotropis procera</i> (Aiton) Dryand.	Roots; Ether, benzene, chloroform, methanol and aqueous; Wistar rats of either sex (125–175 g)	CPE (200 mg/kg bw); Reference drug, Diclofenac sodium, (25 mg/kg bw); AIA (Babu and Karki, 2011)	Flowers; chloroform, acetone and methanolic; <i>Salmonella paratyphi</i> , <i>Bacillus subtilis</i> , <i>Bacillus thuringiensis</i> , <i>Proteus mirabilis</i> , <i>Proteus vulgaris</i> , <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i> and <i>Escherichia coli</i>	Methanol; against all the pathogens tested; alkaloids, tannins, steroids, glycosides, saponins, phenols and flavonoids (Prabha and Vasantha, 2012)	NE
	Latex; Aqueous; Rats	CPE, FoPE; NA; AIA (Kumar and Basu, 1994)	Root and leaves; water, methanol and ethanol; <i>Pseudomonas aeruginosa</i> , <i>Salmonella typhi</i> , <i>Escherichia coli</i> , <i>Staphylococcus aureus</i> and <i>Streptococcus pyrogenes</i>	Aqueous; alkaloids, flavonoids, tannins, saponins, and cardiac glycosides balsams and volatile oil and steroids (Mainasara et al., 2012)	
<i>Cannabis sativa</i> L.	Seeds; Petroleum oil; Male and female albino rats (100–130 g)	CPE (0.5 and 1 ml/kg bw); Reference drug, Indomethacin (10 mg/kg bw); AIA (Musa et al., 2011)	Leaves; ethanol. Methanol, aqueous benzene; <i>Pseudomonas</i> , <i>Staphylococcus</i> , <i>Escherichia coli</i> , <i>Klebsiella</i>	NA; Tannins, terpenoids and reducing sugars (V. Kumar et al., 2011)	2-dodecanone [oil]; Antifungal agent.
					Cannabidiol and cannabigerolic acid; Antibiotic properties. 1-(3,5-Dihydroxyphenyl)-2-(4-hydroxyphenyl)ethane; Phytoalexin. 3-(3,7-Dimethyl-2,6-octadienyl)-2,4-dihydroxy-6-pentylbenzoic acid; Antibiotic.
<i>Cassia fistula</i> L.	Bark; Methanolic and aqueous; Wistar albino rats (170–200 g) of either sex	CPE (250 and 500 mg/kg bw); Reference drug, Diclofenac sodium (5 mg/kg bw); AIA (Ilavarasan et al., 2005)	Leaves; Hydroalcohol; <i>Staphylococcus aureus</i> , <i>Streptococcus pyogenes</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Aspergillus niger</i> , <i>Aspergillus clavatus</i> , <i>Candida albicans</i>	Hydroalcohol; against all the pathogens tested; Tannins, flavonoids, saponins, triterpenoids, steroids, glycosides, anthraquinones, reducing sugars, carbohydrates, proteins, amino acids (Bhalodia and Shukla, 2011)	2-Hentriacontanone and 5-Nonatetracontanone [pods]; Antibacterial agents.
					Celastrol; Cytotoxic and antibacterial agent.

Table 4 (continued)

Plant species	Antiinflammatory studies		Microbiological studies		Active compound [Plant part]; Biological activity related to skin problems (DNP, 2014)
	Plant part; Extract; Animal (Weight)	Animal Model (Dose); Reference control; Activity [Source for columns 2 & 3]	Plant part tested; Solvent or extracts used; Tested against (micro-organism)	Effective extract; Most effective against; Phytoconstituents [Source for columns 4 and 5]	
<i>Celastrus paniculatus</i> Willd.	Seeds; Alcoholic and methanolic; Albino mice (25–32 g)	CPE (500 mg/kg bw); Reference drug, Diclofenac sodium (50 mg/kg bw); AIA (Parimala et al., 2009)	Oil: Bacterial such as <i>Escherichia coli</i> , <i>Staphylococcus aureus</i> , <i>Bacillus cereus</i> , <i>Salmonella dysenterica</i> , <i>Klebsiella pneumoniae</i> , <i>Proteus vulgaris</i> and Fungi such as <i>Aspergillus niger</i> , <i>Aspergillus flavus</i> , <i>Penicillin</i> sp. and <i>Trichoderma</i> sp. Seed; Aqueous, ethanolic; Fungi such as <i>Fusarium oxysporum</i> , <i>Aspergillus terreus</i> , <i>Curvularia lunata</i> and <i>Alternaria solani</i> and Bacteria such as <i>Bacillus subtilis</i> (strain APR-4) and <i>Escherichia coli</i> (EC-1).	Oil; Weak activity against <i>Proteus vulgaris</i> , <i>Staphylococcus aureus</i> , <i>Salmonella dysenterica</i> (Patel and Trivedi, 1962) Aqueous; <i>Fusarium oxysporum</i> , <i>Aspergillus terreus</i> , <i>Curvularia lunata</i> (Parkash and Sandhu, 2012)	
<i>Chrysopogon zizanioides</i> (L.) Roberty	Root; n-Hexane, chloroform, ethyl acetate and butanol; Male Wistar albino rats (250–300 g)	CPE; Reference drug, Diclofenac sodium (10 mg/kg bw); AIA (Kamble et al., 2013)	Plant; Ethanolic and aqueous; <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , and <i>Bacillus subtilis</i>	Ethanol; <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> ; flavanoids, glycosides, phenols, tannins, saponins and alkaloids (Devprakash et al., 2011)	NE
<i>Cissampelos pareira</i> L.	Roots; Ethanolic; Rats	CPE, FoPE, HPE, CPG (200 and 400 mg/kg bw); NA; Significant AIA (Amresh et al., 2007b)	Roots; Methanol; <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i>	Methanol; <i>Bacillus subtilis</i> ; Tannins, flavonoids, steroids (Kumar et al., 2012)	Cycleanine; Antiinflammatory properties.
<i>Cleome viscosa</i> L.	Flowers; Quercetin 3-O-(2"-acetyl)-glucoside obtained from ethyl acetate fraction	CPE; NA; AIA (Senthamsilvelvi et al., 2012)	Seeds; Petroleum ether, chloroform, ethyl acetate, ethanol and water; <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> and four fungal species <i>Aspergillus niger</i> , <i>Sacromyces cerviceae</i> , <i>Aspergillus flavus</i> , <i>Candida albicans</i>	Chloroform, ethyl acetate, ethanol; all tested microorganisms; NA (Wake et al., 2011)	Grandirubrine; Shows potent cytotoxic activity. Norrufescine; Exhibits cytotoxicity against P388 cells. Isoliquiritigenin; Antitumour promoter, antioxidant, antiinflammatory, antimutagenic and antifungal activity.
<i>Clerodendrum infortunatum</i> L.	Leaves; Methanol; Wistar strain rats of either sex (150–180 g)	CPE, HPE, DPE (250 and 500 mg/kg bw); Reference drug, Phenylbutazone (100 mg/kg bw); AIA (S. Das et al., 2010)	Leaves; Chloroform, ethanol, methanol, iso-amyl alcohol and propanol; <i>Bacillus subtilis</i> , <i>Proteus</i> sp., <i>Staphylococcus aureus</i> , <i>Klebsiella pneumoniae</i> and <i>Salmonella typhi</i>	e Iso-amyl Alcohol; <i>Bacillus subtilis</i> and <i>Staphylococcus aureus</i> ; Glycosides, Terpenoids, Anthraquinones, Flavonoids, Saponins, Tannins, Lignin, Phenol and Alkaloids (Prasad et al., 2012)	2-Amino-9-(4-oxo-2-azetidiny) and nonanoic acid [seed]; Antibacterial agents. NE
<i>Commelina benghalensis</i> L.	Leaves; Hydroethanolic; Wistar albino rats (150–200 g) and Swiss albino mice (25–30 g) of either sex	CPE, CPG, XPE (200 and 400 mg/kg bw); Reference drug, Indomethacin (10 mg/kg bw); AIA (Tiwari et al., 2013)	Whole plant; aqueous, ethanol; <i>Candida albicans</i> , <i>Escherichia coli</i> , <i>Staphylococcus aureus</i>	Ethanol; all tested microorganisms; alkaloids, lactones and coumarins, triterpenes and steroids, resins, reducing agents, phenols and tannins, amino acids, quinones, flavonoids and saponins (Cuellar et al., 2010)	NE
<i>Crotalaria juncea</i> L.	Leaves; Ethanolic; Male albino Wistar rats (150–250 g)	CFAA; NA; Significant AIA (Ashok et al., 2006)	Flower and seeds; Ethanol; <i>Escherichia coli</i> , <i>Enterococcus faecalis</i> , <i>Klebsiella pneumoniae</i> , <i>Pseudomonas aeruginosa</i> , <i>Shigella flexneri</i> , <i>Shigella dysenteriae</i> , <i>Staphylococcus aureus</i> , <i>Vibrio cholerae</i>	Ethanol ; <i>Klebsiella pneumoniae</i> , <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i> ; steroids, triterpenes, flavonoids, phenolics and glycosides (Chouhan and Singh, 2010)	NE
<i>Cryptolepis dubia</i> (Burm. f.) M.R. Almeida	Aerial part; Ethanolic; Rat	CPE, CPG; NA; AIA (Laupattarakasem et al., 2006)	Leaves; Methanol; <i>Staphylococcus aureus</i> , <i>Staphylococcus epidermidis</i> , <i>Salmonella typhi</i> , <i>Salmonella paratyphi</i>	Methanol; Against all bacteria (Mahida and Mohan, 2007)	NE

NE

<i>Curculigo orchioides</i> Gaertn.	Tubers; Methanolic; Adult Wistar albino rats (150–200 g) of either sex	CPE (200 and 400 mg/kg bw); Reference drug, Diclofenac sodium (15 mg/kg bw); AIA (Agrahari et al., 2010b)	Roots; Methanol, acetonitrile, chloroform, hexane; <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i> , <i>Klebsiella pneumoniae</i>	Methanol; <i>Pseudomonas aeruginosa</i> ; Saponins (Singh and Gupta, 2008)	
<i>Cuscuta reflexa</i> Roxb.	Stem; Alcoholic and aqueous; Wistar albino rats of either sex (150–200 g) and albino mice (16–25 g)	HPE (100, 200 and 400 mg/kg bw); Reference drug, Ibuprofen (40 mg/kg bw); AIA (Katiyar et al., 2012)	Stem; Ethanol; Gram positive bacteria <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> and Gram negative bacteria <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> Fungi <i>Penicillium citrium</i> , <i>Aspergillus niger</i>	Ethanol; Gram negative bacteria and fungi; Alkaloids, carbohydrates, some glycosides, flavonoids, tannins, phenolic compounds, steroids (Inamdare et al., 2011)	NE
<i>Cynodon dactylon</i> (L.) Pers.	Whole plant; Chloroform, methanolic	CPE; Reference drug, Indomethacin (5 mg/kg bw); AIA (Yogesh et al., 2013)	Leaves; Acetone, ethanol, propanol; <i>Bacillus subtilis</i> , <i>Enterobacter aerogenes</i> , <i>Escherichia coli</i> , <i>Klebsiella pneumoniae</i> , <i>Proteus vulgaris</i> , <i>Pseudomonas aeruginosa</i> , <i>Salmonella typhi</i> , <i>Shigella</i> sp., <i>Staphylococcus aureus</i> , <i>Streptococcus</i> sp.	Acetone; <i>Enterobacter aerogenes</i> ; Alkaloids, phytosterols, tannins (Hema et al., 2013)	NE
<i>Dalbergia sissoo</i> DC.	Leaves; Ethanolic; Rats	CPE, KPE, NPE, CPG (100, 300 and 1000 mg/kg bw); NA; Significant AIA (Hajare et al., 2001)	Leaves; Chloroform, ethyl acetate, acetone, methanol; <i>Escherichia coli</i> , <i>Staphylococcus aureus</i> , <i>Pseudomonas aeruginosa</i>	Methanol; <i>Staphylococcus aureus</i> , <i>Pseudomonas aeruginosa</i> (Prasad et al., 2014)	4-Methoxydalbergione, 3,5-Dihydroxystilbene; Antifungal and antibacterial agents.
<i>Datura stramonium</i> L.	Leaves; Ethanolic; Albino Wister rats of either sex (100–160 g)	CPE (50, 100 and 200 mg/kg bw); Reference drug, Diclofenac sodium (5 mg/kg bw) (Gupta et al., 2010)	Aerial parts; Methanol; <i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Enterococcus faecalis</i> , <i>Pseudomonas aeruginosa</i>	Methanol; Slight antibacterial property (Eftekhar et al., 2005)	1,2,4,5-Tetrahydroxy-7-methylantraquinone; Antifungal agent. NE
<i>Dendrophthoe falcata</i> (L.f.) Ettingsh.	Leaves; Aqueous and methanolic; Rats	CPE, CPG (300 mg/kg bw); Reference drug, Diclofenac sodium; AIA (Patil et al., 2011)	Stem bark; Aqueous, ethanol; <i>Staphylococcus aureus</i> , <i>Salmonella typhi</i> , <i>Shigella</i> spp., <i>Escherichia coli</i> , <i>Klebsiella pneumoniae</i> , <i>Nisseria gonorrhoeae</i>	Ethanol; <i>Staphylococcus aureus</i> , <i>Salmonella typhi</i> , <i>Shigella</i> spp., <i>Escherichia coli</i> , <i>Klebsiellapneumoniae</i> ; Saponins, tannins, steroids, alkaloids, flavonoids, phenols and glycosides (Shagal et al., 2012)	NE
<i>Desmodium gangeticum</i> (L.) DC.	Aerial parts and roots; Water decoction; Rats	CPE (5, 10 and 20 mg/kg bw); NA; AIA (Rathi et al., 2004)	Aerial parts; Petroleum ether, chloroform and ethanol; <i>Staphylococcus aureus</i> , <i>Staphylococcus pyogenes</i> , <i>Staphylococcus epidermidis</i> , <i>Micrococcus luteus</i> , <i>Bacillus subtilis</i> , <i>Bacillus cereus</i> , <i>Klebsiella pneumoniae</i> , <i>Enterobacter aerogenes</i> , <i>Pseudomonas aeruginosa</i> , <i>Serratia marcescens</i> , and five fungi <i>Candida albicans</i> , <i>Candida tropicalis</i> : dimorphic fungi, <i>Aspergillus fumigatus</i> , <i>Aspergillus niger</i> : systemic fungi, and some infectious bacteria <i>Escherichia coli</i> and <i>Salmonella typhi</i>	NE (Pattanayak and Sunita, 2008)	
<i>Desmodium gangeticum</i> (L.) DC.	Aerial parts and roots; Water decoction; Rats	CPE (5, 10 and 20 mg/kg bw); NA; AIA (Rathi et al., 2004)	Whole plant; Methanol, ethanol, chloroform and aqueous; <i>Klebsiella pneumoniae</i> , <i>Escherichia coli</i> , <i>Salmonella typhi</i> , <i>Streptococcus mutans</i> and <i>Pseudomonas aeruginosa</i>	Methanol; <i>Streptococcus mutans</i> ; NA (Karthikeyan et al., 2012)	2',4',5,7-Tetrahydroxy-6-prenylisoflavanone and desmocarpin; Phytoalexin.
<i>Desmochachya bipinnata</i> (L.) Stapf	Root; Hydroalcoholic; Wister albino rats (160–190 g)	CPE (200, 300 and 400 mg/kg bw); Reference drug, Indomethacin (5 mg/kg bw); Significant AIA (V. Kumar et al., 2010)	Aerial part oil; <i>Staphylococcus aureus</i> , <i>Staphylococcus epidermidis</i> , <i>Escherichia coli</i> and <i>Pseudomonas aeruginosa</i>	Against all the tested organisms; Camphene, isobornyl acetate, tricyclene, (+,-) trans-2,6-gamma-Irone, Caryophyllene diepoxide, β -eudesmol, Eseroline and Calarene (K.A. Kumar et al., 2010)	NE
<i>Dicliptera paniculata</i> (Forssk.) I. Darbysh.	Leaves, stem, flowers, seeds and whole plant; Ethanolic; Sprague-Dawley rats (110–150 g) and albino mice (15–20 g) of either sex	CPE, CPG (50, 100, 200 mg/kg bw); Reference drug, Phenylbutazone (100 mg/kg bw); Dose dependent AIA (Rathi et al., 2003)	Leaves; chloroform, acetone, ethanol and water; <i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> , <i>Pseudomonas aeruginosa</i> and fungal strains <i>Aspergillus niger</i> and <i>Penicillium chrysogenum</i>	Acetone and ethanol; <i>Staphylococcus aureus</i> and <i>Bacillus subtilis</i> ; NA (Hosamani et al., 2011)	NE
<i>Eclipta prostrata</i> (L.) L.	Leaves; Methanolic; Albino Wistar rats of either sex (160–180 g)	CPE (100 and 200 mg/kg bw); Reference drugs, Indomethacin (10 mg/kg bw) and cyproheptadine (8 mg/kg	Whole plant; ethyl acetate, ethanol, petroleum ether; <i>Klebsiella pneumoniae</i> , <i>Escherichia coli</i> , <i>Salmonella typhi</i> and <i>Pseudomonas aeruginosa</i> , <i>Bacillus cereus</i> , <i>micrococcus luteus</i> , <i>Candida albicans</i>	Ethyl acetate; Alkaloids, flavonoids, tannins, saponins (Sharma and Sharma, 2010)	Verazine; Antifungal agent.

Table 4 (continued)

Plant species	Antiinflammatory studies		Microbiological studies		Active compound [Plant part]; Biological activity related to skin problems (DNP, 2014)
	Plant part; Extract; Animal (Weight)	Animal Model (Dose); Reference control; Activity [Source for columns 2 & 3]	Plant part tested; Solvent or extracts used; Tested against (micro-organism)	Effective extract; Most effective against; Phytoconstituents [Source for columns 4 and 5]	
<i>Ehretia laevis</i> Roxb.	NE	bw); AIA (Arunachalam et al., 2009)	NE	NE	NE
<i>Eulaliopsis binata</i> (Retz.) C.E.Hubb.	NE	NE	NE	NE	NE
<i>Euphorbia hirta</i> L.	Whole plant; Aqueous and ethanolic; Male albino Wistar rats (180–250 g)	CPE (100 and 200 mg/kg bw); Reference drug, Diclofenac sodium (50 mg/kg bw); AIA (P. Das et al., 2010)	Whole plant; Methanol, hexane, distilled water; <i>Escherichia coli</i> , <i>Klebsiell pneumonia</i> , <i>Salmonella dysentriae</i> , <i>Salmonella typhi</i> , <i>Proteus mirabilis</i> Roots, stem, bud, leaves; Ethanol, petroleum ether; <i>Staphylococcus aureus</i> , <i>Salmonell typhi</i> , <i>Pseudomonas aeruginosa</i> , <i>Vibrio cholera</i> , <i>Escherichia coli</i>	Aqueous; <i>Escherichia coli</i> , <i>Salmonell typhi</i> ; Tannins, saponins, phenolics, flavonoids, cardiac glycosides, anthroquinones, alkaloids (Abubakar, 2009) Ethanol; <i>Salmonell typhi</i> ; NA (Saravanan et al., 2012)	NE
<i>Euphorbia thymifolia</i> L.	Whole plant; Ethanolic; Wistar albino rats (150–200 g) of either sex	CPE (100 mg/kg bw); Reference drug, Indomethacin (10 mg/kg bw); AIA (Garipelli et al., 2012)	Whole plant; Fresh juice, latex, methanol, ethanol, DCM, aqueous; <i>Bacillus pumilus</i> , <i>Staphylococcus pneumoniae</i> , <i>Escherichia coli</i> , <i>Citrobacter freundii</i> , <i>Klebsiell pneumonia</i> , <i>Candida albicans</i> , <i>Aspergillus niger</i>	Fresh juice; <i>Bacillus pumilus</i> , <i>Staphylococcus aureus</i> , <i>Streptococcus pneumoniae</i> , <i>Escherichia coli</i> , <i>Citrobacter freundii</i> , <i>Klebsiell pneumonia</i> ; Tannins, alkaloids, flavanoids (Hussain et al., 2014)	Tellimagrandin II; Antifungal and antiviral agent.
<i>Ficus benghalensis</i> L.	Bark; Aqueous, alcoholic; chloroform	HRBC [100 and 200 mg/ml]; Reference drug, Diclofenac sodium; AIA (Matpal et al., 2013)	Roots; Aqueous, ethanol; <i>Staphylococcus aureus</i> , <i>Pseudomonas aeruginosa</i> , <i>Klebsiell pneumonia</i>	Ethanol; <i>Staphylococcus aureus</i> ; NA (Murti and Kumar, 2011)	NE
<i>Ficus racemosa</i> L.	Leaves; Rat	CPE, SPE, HPE, DPE (200 and 400 mg/kg bw); Reference drug, Phenylbutazone; AIA (Mandal et al., 2000)	Roots; Aqueous, ethanol; <i>Staphylococcus aureus</i> , <i>Pseudomonas aeruginosa</i> , <i>Klebsiell pneumonia</i>	Ethanol; <i>Staphylococcus aureus</i> ; NA (Murti and Kumar, 2011)	Racemosic acid; Anti-inflammatory and antioxidant properties.
<i>Ficus religiosa</i> L.	Bark; Methanolic; Wistar albino male rats (170–190 g) and Swiss albino male mice (27–5 g)	CPE (125, 250 and 500 mg/kg bw); Reference drug, Indomethacin (5 mg/kg bw); AIA (Sreelekshmi et al., 2007)	Roots; Methanol; <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i>	<i>Bacillus subtilis</i> ; Tannins, flavonoids, steroids (Kumar et al., 2012)	
<i>Holarrhena pubescens</i> Wall. ex G. Don	Seeds; Ethanolic; Albino Wistar rats (180–200 g) of either sex	CPE, CPG (100, 200 and 400 mg/kg bw); Reference drug, Indomethacin (10 mg/kg bw); AIA (Saha and Subrahmanyam, 2013)	Bark and leaves; Diethyl ether and methanol; <i>Escherichia coli</i> , <i>Staphylococcus aureus</i> , <i>Pseudomonas aurignosa</i> and <i>Aspergillus niger</i>	Methanol extract of bark and leaves; <i>Escherichia coli</i> , <i>Staphylococcus aureus</i> , <i>Pseudomonas aurignosa</i> and <i>Aspergillus niger</i> (Ramakrishnaiah and Hariprasad, 2012)	Bergaptol; Anti-inflammatory and antifungal agent.
<i>Holoptelea integrifolia</i> Planch.	Leaves; Ethanolic; Wistar rats (180–200 g) and Swiss albino mice (18–22 g) of either sex	CPE, DPE, HPE, SPE, CPG (250 and 500 mg/kg bw); Reference drug, Indomethacin (10 mg/kg bw); AIA (Kalpana and Upadhyay, 2010)	Bark; Methanol; <i>Bacillus cereus</i> , <i>Bacillus subtilis</i> , <i>Bacillus thuringiensis</i> , <i>Staphylococcus epidermidis</i> , <i>Staphylococcus saprophyticus</i> , <i>Staphylococcus aureus</i> , <i>Corynebacterium hoffmanii</i> , <i>Corynebacterium xerosis</i> , <i>Streptococcus faecalis</i> , <i>Micrococcus luteus</i> , <i>Escherichia coli</i> , <i>Salmonella typhi</i>	Methanol; <i>Micrococcus luteus</i> ; NA (Siddiqui et al., 2012)	NE
<i>Hyptis suaveolens</i> (L.) Poit.	Leaves; Suaveolol and Methyl suaveolate from leaves; Rat	CODE; Reference drug, Indomethacin; AIA (Grassi et al., 2006)	Stem bark; Petroleum ether, benzene, chloroform, methanol and aqueous; <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Bacillus subtilis</i> , <i>Pseudomonas aeruginosa</i>	Chloroform; against all the tested microorganisms; NA (Nadella and Paarakh, 2011)	1,4-Naphthoquinone; Antibacterial agent.
			Whole plant; Aqueous and ethanol; Fungi <i>Canidida albicans</i> , <i>Collectrotrichum capsici</i> , <i>Fusarium oxysporum</i> F. sp. <i>Lycopersici</i> , and four bacteria <i>Klebsiella pneumoneae</i> , <i>Staphylococcus</i>	Ethanol; against all the tested microorganisms; NE volatile oil, starch, proteins, tannins, saponins, fats, alkaloids and glycosides (Pachkore et al., 2011)	

<i>Ipomoea carnea</i> Jacq.	Leaves; Aqueous; Albino Wistar rats of either sex	CPE (250 and 500 mg/kg bw); Reference standard, Etoricoxib intraperitoneal (6 mg/kg bw); AIA (Khalid et al., 2011)	<i>aureus</i> , <i>Escherichia coli</i> and <i>Pseudomonas aeruginosa</i> Leaves; n-hexane, ethyl acetate, acetone, ethanol and acetone; <i>Escherichia coli</i> , <i>Staphylococcus aureus</i> , <i>Salmonella typhimurium</i> , <i>Pseudomonas aeruginosa</i> , <i>Proteus vulgaris</i> , <i>Bacillus cereus</i>	Acetone and ethanol; <i>Proteus vulgaris</i> , <i>Salmonella typhimurium</i> and <i>Pseudomonas aeruginosa</i> ; NA (Adsull et al., 2012)	NE
<i>Lannea coromandelica</i> (Houtt.) Merr.	Bark; Ethanolic; Rat	CPE, DPE; NA; AIA (Singh and Singh, 2006)	Bark; ethanol, aqueous; <i>Streptococcus pyogenes</i> , <i>Staphylococcus aureus</i> and <i>Candida albicans</i>	Both; against all the tested microorganisms; NA (Kaur et al., 2013)	NE
<i>Lantana camara</i> L.	Aerial parts; Aqueous; Albino rats (200–250 g) of either sex	CPE (300 and 500 mg/kg bw); Reference drug, Ibuprofen (100 mg/kg bw); AIA (Gidwani et al., 2009)	Methanol, ethanol and water; leaves; Four Gram positive and Gram negative bacterial isolates <i>Staphylococcus aureus</i> , <i>Pseudomonas aeruginosa</i> , <i>Klebsiella pneumoniae</i> , <i>Bacillus subtilis</i> and two fungal strains <i>Aspergillus fumigatus</i> and <i>Aspergillus flavus</i> Leaves; Methanol, Petroleum ether, water and Chloroform; <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i> and <i>Staphylococcus saprophyticus</i>	Methanol; <i>Staphylococcus aureus</i> and <i>Pseudomonas aeruginosa</i> ; Phenolic, Flavonoids, Tannin, Saponin (Naz and Bano, 2013)	Lantic acid and 2,10-Bisaboladien-1-one; Antibacterial agents.
<i>Lawsonia inermis</i> L.	NA; A pure compound lawsone isolated from the chloroform extract; Rat	500 mg/kg bw; Reference drug, Phenylbutazone (100 mg/kg bw); AIA (Alia et al., 1995)	Leaves; Methanol; Gram positive; <i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> and <i>Staphylococcus epidermidis</i> and Gram negative; <i>Escherichia coli</i> , <i>Shigella flexneri</i> , <i>Pseudomonas aeruginosa</i> bacteria	Methanol; against all the tested microorganisms; tannin, catechin, saponin, steroids, alkaloids, phenol, anthroquinone, protein and reducing sugar (Mary, 2011) Methanol; against all the tested microorganisms; glycosides, phytosterol, steroids, saponins, tannins and flavonoids (Raja et al., 2013)	Lalioside; Antimicrobial agent.
<i>Leonotis nepetifolia</i> (L.) R.Br.	Aerial parts; Hexane, ethyl acetate and methanolic; Male CD-1 mice (20–25 g)	TPAPE; NA; AIA (Parra-Delgado et al., 2004)	Leaves, stem, inflorescence and root; Petroleum ether, Ethyl acetate and methanol; Gram-negative bacteria <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Salmonella typhimurium</i> , <i>Klebsiella pneumoniae</i> , gram-positive bacteria <i>Bacillus cereus</i> , <i>Bacillus subtilis</i> , <i>Micrococcus luteus</i> , <i>Staphylococcus aureus</i> and one fungal strain <i>Candida albicans</i> Leaves, flower, stem; Methanol; <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> , <i>Micrococcus luteus</i> , <i>Vibrio cholera</i> , <i>Shigella flexneri</i> , <i>Shigella dysenteriae</i> .	Methanol; against all the tested microorganisms; alkaloids, steroids, saponins, tannins, flavanoids, coumarins, volatile oils, glycosides, terpenoids, phenols and glycosides (Gnaneswari and Raju, 2012)	NE
<i>Linum usitatissimum</i> L.	Fixed oil; Wistar strain albino rats (100–150 g) and Swiss albino mice (20–25 g)	CPE, PGPE, AAPE, LPE, HPE, BKPE (3.0 and 2.85 g/kg bw); Reference drugs different as per model; AIA (Kaithwas et al., 2011)	Seeds; Petroleum ether, chloroform, aqueous, ethanol; Gram-positive and negative bacteria <i>Staphylococcus aureus</i> , <i>Bacillus cereus</i> , <i>Klebsiella pneumoniae</i> and <i>Pseudomonas aeruginosa</i>	Petroleum ether; against all the tested microorganisms; NA (Al-Bayati, 2007)	Secoisolaricresinol; Anticancer and antioxidant agent.
<i>Litsea glutinosa</i> (Lour.) C.B. Rob.	NA; Aqueous; Wistar rats (100–175 g) of either sex	CPE, HPE, DPE; Reference drug, Indomethacin (10 mg/kg bw); AIA (Devi and Meera, 2010)	Bark; Petroleum ether, Ethanol, aqueous; Gram positive <i>Staphylococcus aureus</i> , Gram negative like <i>Pseudomonas aeruginosa</i> , <i>Salmonella typhi</i> , <i>E-coli</i> and fungal species <i>Aspergillus fumigatus</i> and <i>Candida albicans</i>	Petroleum ether & Ethanolic; <i>Pseudomonas aeruginosa</i> ; Alkaloids, Proteins, Carbohydrates, Cardiac glycosides, Saponins, Tannins & Phenolics (Hosamath, 2011)	Cyclolinopeptide A; Possesses good cytoprotective properties. Cassythicine; Antimicrobial agent.
<i>Mallotus philippensis</i> (Lam.) Mull. Arg.	Whole plant; 11-O-galloylbergenin isolated from ethanolic extract; Rats	CPE (10, 20 and 30 mg/kg bw); NA; AIA (Arfan et al., 2010)	Whole plant; Hexane, chloroform, ethyl acetate, butanol and aqueous; <i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> , <i>Staphylococcus pneumoniae</i> , <i>Pseudomonas aeruginosa</i> , <i>Proteus vulgaris</i> , <i>Salmonella typhi</i> , <i>Aspergillus flavus</i> , <i>Aspergillus niger</i> and <i>Candida albicans</i>	Ethyl acetate, butanol; against all the tested microorganisms; alkaloids, flavonoids, glycosides, phenols, quinines, saponins, tannins and terpenoids (Afzal et al., 2013)	NE
<i>Martynia annua</i> L.	NE	NE	Leaves; Chloroform, ethyl acetate and methanol; <i>Streptococcus faecalis</i> , <i>Streptococcus pyogenes</i> , <i>Enterococcus faecalis</i> , <i>Bacillus subtilis</i> , <i>Bacillus thuringiensis</i> , <i>Klebsiella pneumoniae</i> ,	In Chloroform extract the activity was high in <i>Proteus vulgaris</i> ; In Ethyl acetate <i>Salmonella paratyphi</i> A, B, <i>Proteus mirabilis</i> , <i>P. vulgaris</i> and <i>Klebsiella pneumoniae</i> . In Methanol extract,	NE

Table 4 (continued)

Plant species	Antiinflammatory studies		Microbiological studies		Active compound [Plant part]; Biological activity related to skin problems (DNP, 2014)
	Plant part; Extract; Animal (Weight)	Animal Model (Dose); Reference control; Activity [Source for columns 2 & 3]	Plant part tested; Solvent or extracts used; Tested against (micro-organism)	Effective extract; Most effective against; Phytoconstituents [Source for columns 4 and 5]	
<i>Millettia extensa</i> (Benth.) Baker	NE	NE	NE	<i>Proteus vulgaris</i> , <i>Bacillus subtilis</i> , <i>Salmonella paratyphi</i> .B. and <i>Pseudomonas aeruginosa</i> ; terpenoid, alkaloids, glycosides, steroids, tannins and saponins and moderate quantity of cardiac glycosides, phenols and anthroquinones (Sermakkani and Thangapandian, 2010)	NE
<i>Mirabilis jalapa</i> L.	Leaves; Aqueous; Albino Wistar rats of either sex (150–200 g)	CPE (200 and 400 mg/kg bw); Reference drug, Diclofenac sodium (10 mg/kg bw); AIA (M. Singh et al., 2010)	Leaves; Methanol, ethanol, petroleum ether and aqueous; <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Klebsiella pneumoniae</i> , <i>Pseudomonas aeruginosa</i> , <i>Salmonella typhi</i> and <i>Proteus mirabilis</i>	Ethanol; <i>Pseudomonas aeruginosa</i> ; alkaloids, saponins, tannins, and flavonoids (Akintobi et al., 2011)	2'-O-Methylabronisoflavone and 4-Hydroxy-9-O-methylboeravinone B; Antifungal agents.
<i>Mitragyna parvifolia</i> (Roxb.) Korth.	Leaves; Ethanolic; Male Wistar rats (150–200 g) and Swiss mice (20–25 g) of either sex	CPE (100, 200 and 300 mg/kg bw); Reference drug, Phenylbutazone (80 mg/kg bw); AIA (Gupta et al., 2009)	Leaves; Ethanol; <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> , <i>Escherichia coli</i> and <i>Pseudomonas aeruginosa</i>	Ethanol; <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> and <i>Pseudomonas aeruginosa</i> ; NA (Kaushik et al., 2009b)	<i>Mirabilis jalapa</i> antimicrobial peptides; Antimicrobial and antifungal agent. <i>Mirabilis jalapa</i> Trypsin inhibitors; Trypsin inhibitor. Dihydrocorynantheol; Shows activity against Gram-positive bacteria.
<i>Momordica dioica</i> Roxb. ex Willd.	Fruit pulp; Hexane and ethyl acetate; Wistar albino mice (25–30 g) and rats (150–200 g) of either sex	CPE (50 and 100 mg/kg bw); Reference drug, Diclofenac sodium (5 mg/kg bw); Potential AIA (Ilango et al., 2003)	Fruit pulp; Hexane and ethyl acetate soluble portion of methanolic extract; Gram positive bacteria; <i>Bacillus cereus</i> , <i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> & <i>Staphylococcus epidermidis</i> and seven Gram negative bacteria; <i>Escherichia coli</i> , <i>Klebsiella pneumoniae</i> , <i>Proteus mirabilis</i> , <i>Proteus vulgaris</i> , <i>Pseudomonas aeruginosa</i> , <i>Salmonella typhi</i> and <i>Shigella dysenteriae</i>	Both extracts; <i>Salmonella typhi</i> and <i>Shigella dysenteriae</i> ; secondary metabolites such as steroids, fatty acids in hexane extract (HE) and proteins, saponin glycosides and triterpenes (Ilango et al., 2012)	Corynantheine; Aniviral agent against influenza A. 3,23-Dihydroxy-12-oleanen-28-oic acid; Epicaricogen inhibitor.
<i>Mucuna pruriens</i> (L.) DC.	Leaf; Ethanol; Male Wistar rats (120–150 g)	CPE (200 mg/kg bw); Reference drug, Indomethacin (10 mg/kg bw); Significant AIA (Nagarani et al., 2014)	Root and seeds; hexane, petroleum ether, benzene, methanol and aqueous; <i>Staphylococcus aureus</i> , <i>Klebsiella pneumoniae</i> , <i>Bacillus subtilis</i> , <i>Pseudomonas aeruginosa</i> , <i>Salmonella typhi</i> and <i>Escherichia coli</i>	Benzene, methanol; against all the tested microorganisms; alkaloids, anthraquinones, flavonoids, phenols, tannins, terpenoids and xanthoprotein (Murugan and Mohan, 2011)	NE
<i>Nicotiana glauca</i> Viv.	Aerial parts; Ethanolic; Swiss albino mice of either sex (20–25 g)	CPE, CPG (200, 400 mg/kg bw); Reference drug, Aspirin (150 mg/kg bw); AIA (Bala et al., 2011)	Leaves; Aqueous and methanol extracts; <i>Bacillus cereus</i> , <i>Bacillus fusiformis</i> , <i>Salmonella typhimurium</i> <i>Staphylococcus aureus</i> and <i>Pseudomonas aeruginosa</i>	Aqueous extract showed strongest activity on <i>Bacillus fusiformis</i> and methanol leaf extract also showed strongest activity on <i>Bacillus fusiformis</i> ; alkaloids, saponin, tannin, flavonoides, cardiac glycosides, phenolic compounds, steroids, terpenoides and carbohydrates (K.P. Singh et al., 2010)	NE
	Root bark; Petroleum ether, chloroform, ethyl acetate and	CPE, CPG (100 and 300 mg/kg bw); Reference drug,		Dichloromethane; <i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> ; Flavonoids (bacailein	Aloe emodin (shows antibacterial activity against methicillin resistant <i>Staphylococcus</i>

<i>Oroxylum indicum</i> (L.) Kurz	n-butanol; Swiss mice (18–22 g) and Wistar albino rats (200–250 g) of either sex	Diclofenac sodium (5 mg/kg bw); AIA (Zaveri and Jain, 2010)	Stem bark, root; Dichloromethane; <i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Candida albicans</i> Root, stem; Alcohol; <i>Escherichia coli</i> , <i>Klebsiella</i> spp., <i>Proteus</i> , <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i>	and chrysin) and a naphthoquinone, lapachol (MatAli et al., 1998) Alcohol stem; <i>Staphylococcus aureus</i> , <i>Klebsiella</i> spp., <i>Proteus</i> , <i>Pseudomonas aeruginosa</i> ; Alkaloids, phenols, fats, lipids, waxes (Radhika et al., 2011)	<i>aureus</i> (MRSA)) and oroxylin A (against gram-positive bacteria); Antibacterial agents. Chrysin; Shows antifungal and antiinflammatory activity. Active against <i>Pseudomonas aeruginosa</i> and <i>Candida albicans</i> . Oroxylin A; Shows antiinflammatory properties. Aloe emodin; Antimicrobial, antimutagenic and antiseptic agent. 4',5,6,7-Tetrahydroxyflavone; Antioxidant. Chrysin; Antifungal agent. NE
<i>Persicaria barbata</i> (L.) H.Hara	Aerial parts; Petroleum ether, ethyl acetate and chloroform; Swiss albino mice (20–25 g) and long evans adult rats (130–170 g) of either sex	CPE (200 and 400 mg/kg bw); NE Reference drug, Phenylbutazone (100 mg/kg bw); AIA (Mazid et al., 2009)	NE	NE	NE
<i>Phyllanthus amarus</i> Schumacher and Thonn.	Whole plant; Methanolic; Male Wistar rats (150–200 g)	PAPE, CAP, CPG; Reference drug, Indomethacin; AIA (Mahat and Patil, 2007)	Leaves; Aqueous; <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> , <i>Streptomyces albus</i> , <i>Streptococcus faecalis</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Klebsiella pneumoniae</i> , <i>Proteus vulgaris</i> Leaves; Petroleum ether, benzene, methanol and aqueous; <i>Escherichia coli</i> , <i>Staphylococcus aureus</i> , <i>Klebsiella pneumoniae</i> , <i>Bacillus subtilis</i> , <i>Enterobacter aerogenes</i> , <i>Pseudomonas aeruginosa</i> , <i>Salmonella typhimurium</i> , <i>Salmonella typhi</i> , <i>Staphylococcus epidermidis</i> and <i>Proteus vulgaris</i> Chloroform, methanol, water; <i>Trichophyton rubrum</i> , <i>Trichophyton mentagrophytes</i> , <i>Trichophyton tonsurans</i> , <i>Microsporum gypseum</i> , <i>Microsporum fulvum</i> NE	Aqueous; <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> , <i>Streptomyces albus</i> , <i>Escherichia coli</i> ; Alkaloids, phytosterols, phenolic compounds and tannins, proteins and amino acids, lignins, saponins (Dhandapani et al., 2007) Methanol; Carbohydrates, alkaloids, flavonoids glycosides, steroids, tannins and saponins (Arote et al., 2009)	3,3',4,4',9,9'-Hexahydroxylignan; Cytotoxicity enhancer. Pongaflavone, 3-Me ether, 9,10-di-Ac and 6-O-Methylolivetone (against <i>Mycobacterium tuberculosis</i>); Antibacterial agents. Karanjabiflavone and $\alpha,\beta,2',3,4,4'$ -Hexahydroxychalcone; Antioxidant.
<i>Pongamia pinnata</i> (L.) Pierre	Leaves; Ethanolic; Rat	CPE, HPE, HTPE, PGPE, KPE, FoPE, CPG (100, 300 and 1000 mg/kg bw); NA; AIA (Srinivasan et al., 2001)	NE	Chloroform; against all tested microorganisms; NA (Sharma et al., 2012b)	NE
<i>Premna mollissima</i> Roth	Leaves; Methanolic; Albino male Swiss mice (18–25 g) and Wistar Rat (150–180 g)	CPE, CPG (125, 250 and 500 mg/kg bw); Reference drug, Indomethacin (10 mg/kg bw); AIA (Mahire et al., 2009)	NE	NE	NE
<i>Ranunculus sceleratus</i> L.	Aerial parts; NA	TPAPE, AAPE, CPE; NA; AIA (Prieto et al., 2003)	Leaves; Chloroform, methanol, water; <i>Trichophyton rubrum</i> , <i>Trichophyton mentagrophytes</i> , <i>Trichophyton tonsurans</i> , <i>Microsporum gypseum</i> , <i>Microsporum fulvum</i> Leaves and roots; Ethanol; <i>Escherichia coli</i> , <i>Bacillus subtilis</i> , <i>Salmonella typhi</i> and <i>Staphylococcus aureus</i>	Chloroform; against all tested microorganisms; NA (Sharma et al., 2012b)	NE
<i>Rauvolfia serpentina</i> (L.) Benth. ex Kurz	NE	NE	NE	Ethanol; root extract against <i>S. typhi</i> ; carbohydrates, tannins, saponins, flavonoids, alkaloids and starch soluble compounds are present and four indole alkaloids reserpine, ajmalicine, ajmaline and yohimbine (Deshmukh et al., 2012) Methanol; NA (Naz and Bano, 2012)	NE
<i>Ricinus communis</i> L.	Roots; Methanolic; Wister albino rats	CPE, CPG (250 and 500 mg/kg bw); NA; AIA (Ilavarasan et al., 2006)	Leaf; Methanol, ethanol and water; Gram positive bacteria (<i>Bacillus subtilis</i> and <i>Staphylococcus aureus</i>) as well as Gram negative bacteria (<i>Pseudomonas aeruginosa</i> and <i>Klebsiella pneumoniae</i>) and fungal strains <i>Aspergillus fumigatus</i> and <i>Aspergillus flavus</i> .		Casbene [seed]; Antifungal agent.
<i>Scoparia dulcis</i> L.	Whole plant; Ethanolic; Mice	CPE (0.5 and 1.0 g/kg); NA; AIA (Tsai et al., 2011) CPE, CPG (150 mg/kg bw); Reference drug, Indomethacin	Whole plant; Aqueous and ethanolic; <i>Staphylococcus aureus</i> , <i>Microsporum canis</i> , <i>Candida albicans</i> , <i>Salmonella typhi</i> , <i>Streptococcus species</i> , <i>Escherichia coli</i> and <i>Proteus vulgaris</i>	Ethanol; Saponins, glycoside and carbohydrate (Yisa, 2009) Petroleum Ether; against all the tested organisms; Phenols, Triterpenoids, Steroids,	6,16-Dihydroxy-18-aphidicolanoic acid; Shows antiviral activity towards Herpes simplex and cytotoxic activity to HeLa cells. Scopadulcic acid A; Antitumour agent. 3-Pentadecyl-1,2-benzenediol; Antibacterial agent, cytotoxic to human cancer cells.

Table 4 (continued)

Plant species	Antiinflammatory studies		Microbiological studies		Active compound [Plant part]; Biological activity related to skin problems (DNP, 2014)
	Plant part; Extract; Animal (Weight)	Animal Model (Dose); Reference control; Activity [Source for columns 2 & 3]	Plant part tested; Solvent or extracts used; Tested against (micro-organism)	Effective extract; Most effective against; Phytoconstituents [Source for columns 4 and 5]	
<i>Semecarpus anacardium</i> L. f.	Nuts; Milk and Clarified butter; Adult male Wistar rats (180 and 200 g)	(10 mg/kg bw); AIA (Ramprasath et al., 2005)	Nut; Petroleum Ether; <i>Escherichia coli</i> , <i>Micrococcus luteus</i> , <i>Salmonella typhi</i> , <i>Bacillus subtilis</i> and <i>Klebsiella pneumonia</i> Nut; Alcoholic; <i>Aspergillus fumigates</i> ; <i>Candida albicans</i>	Alkaloids, Flavonoids, Saponins, Tannins and Anthraquinones (Bagewadi et al., 2012) Alcoholic; NA (Sharma et al., 2002)	
<i>Senna tora</i> (L.) Roxb.	Leaves; Methanolic; Rats	CPE, HPE, SPE, DPE (400 mg/kg bw); NA; AIA (Maity et al., 1998)	Leaves; Hexane, chloroform and methanol, ethyl acetate; Bacteria <i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> , <i>Pseudomonas aeruginosa</i> <i>Proteus vulgaris</i> , <i>Escherichia coli</i> , Fungi <i>Candida albicans</i> , <i>Aspergillus niger</i> , <i>Fusarium oxysporum</i> and <i>Rhizopus oryzae</i>	Hexane; <i>Escherichia coli</i> Ethyl acetate <i>Aspergillus niger</i> ; Phenolics, flavonoids, tannins and anthraquinones (John et al., 2011)	2-Acetyl-1,6,8-trihydroxy-3-methylnaphthalene (strong activity against methicillin-resistant <i>Staphylococcus aureus</i>) and toralactone (against methicillin-resistant <i>Staphylococcus aureus</i>) [seed]; Antibacterial agents. 1,2,6,8-Tetrahydroxy-3-methylantraquinone; Antioxidant. Toralactone [seed]; Antimicrobial agent. 2,3',4,4',5-Pentahydroxy-7,9':7',9'-diepoxy lignan [Seed], Sesamolol [Seed]; Antioxidant. 2-Chloro-5,8-dihydroxy-3-prenyl-1,4-naphthoquinone [Root]; Antifungal agent.
<i>Sesamum indicum</i> L.	Seed; Oil; Male Wistar rats (180–220 g) and male Swiss albino mice (25–30 g) Seed; oil; Wister albino rats (150–200 g) and Swiss albino mice (25–30 g)	CPE (100, 200 and 300 mg/kg bw); NA; AIA (Monteiro et al., 2014) CPE (5 and 10 ml/g bw); Reference drug, Diclofenac sodium (10 mg/kg bw); AIA (Saleem et al., 2011)	NE	NE	
<i>Shorea robusta</i> Gaertn.	Resin; Ethanolic; Male albino rats (150–250 g)	CPE, CPG (30, 100 and 300 mg/kg bw); Reference drug, Etoricoxib (10 mg/kg bw); AIA (Wani et al., 2012b)	Oleoresin; Aqueous, methanol, petroleum and benzene; <i>Bacillus subtilis</i> , <i>Bacillus licheniformis</i> , <i>Bacillus coagulans</i> , <i>Bacillus cereus</i> , <i>Staphylococcus aureus</i> , <i>Staphylococcus epidermidis</i> , <i>Staphylococcus griseus</i> , <i>Escherichia coli</i> , <i>Proteus vulgaris</i> , <i>Pseudomonas fluorescens</i> , <i>Aspergillus flavus</i> , <i>Aspergillus niger</i> , <i>Candida albicans</i> , <i>Penicillium chrysogenum</i> Floral parts; Aqueous; <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> , <i>Klebsiella pneumonia</i> , <i>Serratia marcescens</i>	Aqueous; <i>Bacillus coagulans</i> , <i>Escherichia coli</i> , <i>Bacillus cereus</i> ; Alkaloids, carboxylic acids, fatty acids, phenols, saponins, steroids (Murthy et al., 2011) Aqueous; All; Tannins, flavanoids, cardiac glycosides, steroids (Duddukuri et al., 2011)	Chebulinic acid [Seed]; Antioxidant. NE
<i>Sida cordata</i> (Burm. F.) Borss. Waalk.	NE	NE	NE	NE	NE
<i>Sida rhombifolia</i> L.	Leaves; Hydroalcoholic; Rat	CPE (400 mg/kg bw); Reference drug, Indomethacin (10 mg/kg bw); AIA (Khalil et al., 2006)	NE	NE	NE
<i>Solanum incanum</i> L.	NE	NE	NE	NE	Spirosol-5-en-3-ol; Cytotoxic agent.
<i>Terminalia arjuna</i> (Roxb. ex DC.) Wight & Arn.	Leaves; Methanolic; Adult male Swiss albino mice (20 ± 2 g)	CPE, DPE, HPE (100, 200 mg/kg bw); Reference drug, Indomethacin (10 mg/kg bw); AIA (Biswas et al., 2011b)	Stem barks; Methanol; <i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Klebsiella pneumoneae</i> , <i>Salmonella typhi</i> , <i>Proteus mirabilis</i> , <i>Micrococcus</i> sp.	Methanol; <i>Staphylococcus aureus</i> ; Alkaloids, glycosides, flavonoids, flavanols, phenols, saponins, terpenoids (Patil and Gaikwad, 2011)	2,3,23-Trihydroxy-12-oleanen-28-oic acid; Skin tumour proliferation inhibitor.
<i>Tridax procumbens</i> (L.) L.	Leaves; Aqueous; Male and female albino rats (130–170 g)	CPE (400 mg/kg bw); Reference drug, Ibuprofen; AIA (Awasthi et al., 2009)	Leaves; Methanol, ethyl acetate; <i>Escherichia coli</i> , <i>Klebsiella pneumoniae</i> , <i>Salmonella typhi</i> , <i>Bacillus cereus</i> , <i>Staphylococcus aureus</i>	Ethyl acetate; <i>Staphylococcus aureus</i> ; Alkaloids, flavanoids, glycosides (Bharathi et al., 2012) NE	Punicalin; Moderate cytotoxic agent. 1,2,3,4,6,7-Naphthalenehexol; Antioxidant. NE
			NE	NE	NE

<i>Typha domingensis</i> Pers.	Leaves; Petroleum ether and methanolic; Albino Wistar rats (100–150 g)	CPE, HPE (100, 200, 400 mg/kg bw); Reference drug, Diclofenac sodium (10 mg/kg bw); AIA (Pawar et al., 2011)			
<i>Vallaris solanacea</i> (Roth) Kuntze	Leaves; Methanolic; Albino rats of either sex (150–200 g)	CPE (250, 500 mg/kg bw); Reference drug, Indomethacin (100 mg/kg bw); AIA (Joshi et al., 2014)	Chloroform, ethanol, Petroleum ether; Gram positive- <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> and gram negative – <i>Salmonella typhi</i> and <i>Escherichia coli</i> , Fungus like <i>Candida albicans</i> and <i>Aspergillus niger</i>	Petroleum ether; <i>Salmonella typhi</i> , <i>Escherichia coli</i> , <i>Aspergillus niger</i> and <i>Candida albicans</i> ; NA (Vagdevi et al., 2011)	NE
<i>Vanda tessellata</i> (Roxb.) Hook. ex G.Don.	NE	NE	Leaves; Chloroform, petroleum ether, ethylacetate, acetone and methanol; <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> , <i>Escherichia coli</i> , <i>Klebsiella pneumoniae</i> , <i>Proteus mirabilis</i> and <i>Candida albicans</i>	Ethyl acetate; <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Klebsiella pneumoniae</i> and <i>Candida albicans</i> (Gupta and Katewa, 2012)	NE
<i>Verbascum thapsus</i> L.	NE	NE	NE	NE	Aucubigenin; Antibacterial and phytotoxic agent. 11,13(18)-Oleanadiene-3,16,23,28-tetrol; Antiinflammatory properties.
<i>Vitex negundo</i> L.	Seeds; Chloroform; Sprague-Dawley male rats (100–160 g)	CPE (500 mg/kg); Reference drug, Ibuprofen (50 mg/kg bw); AIA (Chawla et al., 1992)	Leaves; Benzene, chloroform, ethanol, water-ethanol (50:50), water; <i>Staphylococcus aureus</i> , <i>Proteus vulgaris</i> , <i>Bacillus subtilis</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Aspergillus niger</i> , <i>Aspergillus flavus</i> , <i>Candida albicans</i>	Water-ethanol (50:50); showed maximum antimicrobial, Water; showed maximum antifungal activity; α -pinene, camphene, caryophyllene, citral, glycosides like negundoside, nishinadaside, other hydroxybenzoylmussaenosidic acid derivatives (Aswar et al., 2009)	Vitegnoside and Vitelignin A [leaf]; Antifungal agents.
<i>Wrightia arborea</i> (Dennst.) Mabb.	Leaves; Methanolic; Male albino rats (120–140 g)	FPE, CPE (100, 200 mg/kg bw); Reference drug, Indomethacin (10 mg/kg bw); AIA (Nahar et al., 2013)	Bark; Petroleum ether, chloroform, acetone and methanol; <i>Bacillus subtilis</i> , <i>Bacillus megaterium</i> , <i>Escherichia coli</i> , <i>Klebsiella planticola</i> , <i>Micrococcus luteus</i> , <i>Pseudomonas aeruginosa</i> <i>Staphylococcus aureus</i> and <i>Salmonella typhi</i>	Chloroform ; <i>Micrococcus luteus</i> , <i>Staphylococcus aureus</i> , <i>Salmonella typhi</i> and <i>Pseudomonas aeruginosa</i> ; alkaloids, phenolics, saponins, flavonoids and tannins (Khyade and Vaikos, 2011)	NE
<i>Ziziphus nummularia</i> (Burm. f.) Wight & Arn.	Leaves; Alkaloids; Wistar rats of both sexes (180–200 g) and Swiss mice (20–24 g)	CPE, HPE, DPE, SPE, CPG, CIP (10, 20, 30 mg/kg bw oral); NA; 30 mg/kg bw shows anti-oedematogenic effect (Goyal et al., 2013)	Leaves; Aqueous, ethanol; <i>Staphylococcus aureus</i> , <i>Streptococcus pyogenes</i> , <i>Bacillus subtilis</i> , <i>Pseudomonas aeruginosa</i> , <i>Aspergillus niger</i> , <i>Aspergillus flavus</i> , <i>Candida albicans</i> , <i>Trichophyton rubrum</i>	Ethanol; <i>Staphylococcus aureus</i> and fungi <i>Trichophyton rubrum</i> ; NA (Gautam et al., 2011)	NE

Abbreviations: AAPE=arachidonic acid (AA) induced paw edema model, AIA=anti-inflammatory activity, APE=adjuvant induced paw edema model, BKPE=Bradykinin induced paw edema model, CAP=carrageenan air pouch model, CFAA=complete Freund's adjuvant (CFA) induced arthritis model, CIP=carrageenan induced peritonitis model, CODE=Croton oil induced dermatitis of the mouse ear model, CPE=carrageenan induced paw edema model, CPG=cotton pellet granuloma model, DPE=dextran induced paw edema model, FoPE=formaldehyde induced paw edema model, FPE=formaline induced paw edema model, HPE=histamine induced paw edema model, HRBC=human blood cell membrane stabilization method, HTPPE=5-hydroxytryptamine induced paw edema model, IPE=immunologically induced paw edema model, KPE=Kaolin induced paw edema model, LPE=leukotriene induced paw edema model, NE=not evaluated, NPE=Nystatin induced paw edema model, PAPE=phlogistic agents induced paw edema model, PGIN=peptidoglycan (PGN) induced inflammatory reaction, PGPE=prostaglandin E2 induced paw edema model, PPCIN=polyinosinic: polycytidylic acid (polyI:C) induced inflammatory reaction, SPE=serotonin induced paw edema model, TPAPE=tetradecanoylphorbol acetate (TPA) induced paw edema model, XEE=xylene induced ear edema model.

Quercetin, the active ingredient, has been shown in *in vitro* studies to decrease fibroblast proliferation, inflammation, extracellular matrix deposition, and stabilized mast cells (Pawlikowska-Pawlega and Gawron, 1995; Ho et al., 2006; Cho et al., 2010). A prospective, double-blinded, split-scar study on 20 Asians who had new Pfannenstiel's cesarean section scars was conducted to assess efficacy of onion gel containing 12% *Allium cepa* L. (Erase gel, ABCA Pharma Lab Co., Ltd., Nonthaburi, Thailand) and results showed improvement of scar height and scar symptoms (Chanprapaph et al., 2012). The randomized, controlled, single-blind study on 44 healthy male and female aged 18–70 years evaluated the appearance of new dermal scars after eight weeks of once-daily application of non-prescription proprietary onion extract gel formulation (Mederma®, Merz Pharmaceuticals, LLC, Greensboro, North Carolina), the study showed that this proprietary onion extract gel is safe and significantly improves scar appearance after four weeks of once-daily application (Draeos et al., 2012).

Amaranthus spinosus L. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). The methanol extracts of *Amaranthus spinosus* L. leaves reveals significant antitumor effects in cancers of breast, colorectal, liver and normal cell lines (Rajasekaran et al., 2014). Ethanolic extract of this plant was found to be non-toxic to Wistar albino rats even at high doses of 2000 mg/kg bw (S.B. Mishra et al., 2012). Preclinical trials have shown this plant as a promising candidate for development of some skin care products. However human clinical trial related to any skin disease has not been conducted on this plant.

Anisomeles indica (L.) Kuntze has shown antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Methanolic extract and purified ovatodiolide from whole plant of *Anisomeles indica* (L.) Kuntze has shown tyrosinase inhibitory property suggesting that they could be applied as a type of dermatological whitening agent in skin care products (Huang et al., 2012). Ovatodiolide also showed cytotoxicity causing apoptosis in producing reactive oxygen species and down-regulation of FLICE inhibitory prote in leading to cell cycle arrest towards oral squamous cell carcinoma (Hou et al., 2009). It possesses significant antiinflammatory activity by inhibiting the enhanced production of nitric oxide (NO) radicals and pro-inflammatory cytokines (TNF- α and IL-12) induced by LPS/IFN- γ on murine peritoneal macrophages (Hsieh et al., 2008). A study by Hsu et al. (2012a) demonstrated that the hexane extract of this plant can induce apoptosis of FaDu human pharynx squamous cancer cells by down regulating the expressions of Bcl-2 and Bcl-xL proteins, up-regulating the expressions of Bax and Bak proteins, and activating caspase-9 and caspase-3. In another study Hsu et al. (2012b) demonstrated that hexane extract of this plant significantly inhibited migration and invasion of FaDu cells in a dose-dependent manner under non-cytotoxic concentrations by suppressing the expression of metalloproteinase-9 and metalloproteinase-2. Acute oral toxicity study of aerial parts of this plant on Wistar rats showed that ethyl acetate and chloroform extract of this plant is non-toxic even at high doses of 2000 mg/kg bw (Sundriyal et al., 2013). However human clinical trial related to any skin disease has not been conducted on this plant.

Annona squamosa L. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Seeds of this plant contain a caustic resin that is a toxic irritant principal and can be used to get rid of hair lice (Perry and Metzger, 1980). A clinical trial showed that the seed and the leaf can treat hair lice with minor side effects or irritation (Puapattanakul, 1980). In unblinded, controlled clinical trial the application of cream based on custard apple seed extract killed > 90% of lice (Tiangda et al., 2000). Crude extract of seeds of this plant was non-toxic at lower dose (300 mg/kg bw) but toxic at

higher dose of 2000 mg/kg bw to Wistar rats (Aneela et al., 2011). *Argemone mexicana* L. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4) but it is generally orally toxic (Table 3). However, in the present study seed paste and latex of this plant are reported to be used topically (Table 1). Acetone, methanol, ethanol and aqueous leaf extracts of *Argemone mexicana* L. showed antipseudomonal activity against multidrug resistant *Pseudomonas aeruginosa* isolated from clinical samples (Sahu et al., 2012). Aqueous extract of aerial parts showed anti-parasite activity against the chloroquine resistant K1 strain of *Plasmodium falciparum* with IC₅₀ value of 5.89 μ g/ml (Schrader et al., 2012). In a randomized, controlled clinical trial 89% patients recovered clinically although parasite clearance was achieved in 9% patients (Schrader et al., 2012). However human clinical trial related to any skin disease has not been conducted on *Annona squamosa* L. and *Argemone mexicana* L.

Azadirachta indica A. Juss. has shown wound healing, antiinflammatory, antioxidant, antimicrobial and tyrosinase inhibitory properties in the earlier studies (Tables 3 and 4). There are large numbers of Ayurvedic skin care products made from this plant available in the market. It is part of polyherbal drug of Goodcare Pharma Pvt. Ltd. made of 3 herbs known as “Neem Guard capsules”, which is used in skin diseases. Clinical study was done on 57 patients with symptoms of skin diseases and this capsules were found to be effective in lowering IgE level in case of skin diseases and was effective in some selective diseases like acne, contact dermatitis, urticaria, allergic dermatitis and wet eczema (Allayurveda.com, 2014). *Azadirachta indica* A. Juss. is also a part of polyherbal Ayurvedic formulation of The Himalaya Drug Company made of 8 herbs known as “Anti-dandruff Hair cream”, which is recommended for the treatment of dandruff. The open, non-comparative, phase III clinical trial of this cream was done on 50 patients to evaluate the clinical efficacy and safety (short- and long-term) of this cream in the management of dandruff (Agarwal et al., 2009). The study observed significant symptomatic and clinical improvement of dandruff in 6 weeks and concluded that this cream is effective and safe in the management of dandruff (Agarwal et al., 2009). *Azadirachta indica* A. Juss. along with 6 other herbs is a part of another Ayurvedic formulation which was tested for treating acne vulgaris (Lalla et al., 2001). A Randomized, double-blind, placebo-controlled Phase II clinical trials were conducted on 53 patients for 4 weeks to test efficacy of this formulations and it was observed that combination of use of internal and external preparations showed better efficacy as compared to the used of oral formulations alone (Lalla et al., 2001). In a limited clinical trial oral administration of 5 g of an aqueous leaf paste of this plant enabled diabetic patients to reduce their dosages of insulin up to 30 to 50% without a significant effect on the blood glucose levels (Shukla et al., 1973). Azadirachtin obtained from *Azadirachta indica* A. Juss. possess anti-tumor properties and has the potential to target NF- κ B (Thoh et al., 2010). Nimbolide, another compound derived from leaves and flowers of this plant has shown numerous biological activities including anti-cancer activity (H.G. Kumar et al., 2010).

Bambusa bambos (L.) Voss has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Acute oral toxicity study on this plant revealed that aqueous extract of leaves of this plant was safe to Swiss albino mice even up to dose of 1000 mg/kg bw (Kundu et al., 2011). *Bauhinia variegata* L. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). This plant also has trypsin inhibitory properties (de Souza et al., 2005). This plant was non-toxic to experimental animals up to oral dose of 2000 mg/kg bw (Table 3). 3',4',5,7-Tetrahydroxy-3-methoxyflavone present in this plant is an

antiinflammatory agent (DNP, 2014). Water fraction of *Bauhinia variegata* L. leaf showed pronounced cytotoxic effect against DU-145, HOP-62, IGR-OV-1, MCF-7 and THP-1 human cancer cell lines with 90–99% cell growth inhibitory activity, whereas ethyl acetate fraction also produced considerable cytotoxicity against MCF-7 and THP-1 cell lines (Mishra et al., 2013). The healing potential of lectin of *Bauhinia variegata* L. and its recombinant isoform was evaluated on mice and study indicated that the lectin and its recombinant isoform possess pro-healing properties and may be employed in the treatment of acute skin wounds (Neto et al., 2011). Ethanolic extract of *Bauhinia variegata* L. has shown potential chemopreventive property against N-nitrosodiethylamine induce liver tumor and human cancer cell lines (Rajkapoor et al., 2006). Chemopreventive potential of bark of this plant against DMBA-induced skin papillomagenesis was studied in mice which showed a significant reduction in the skin tumors in treated animals as compared to the control group of animals (Singh and Kale, 2010). However human clinical trial related to any skin disease has not been conducted on *Bambusa bambos* (L.) Voss and *Bauhinia variegata* L.

Boerhavia diffusa L. has shown antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Aqueous extract of leaves of this plant was non-toxic to Wistar albino rats and albino mice even up to dose of 2000 mg/kg bw (Orisakwe et al., 2003). Aqueous methanol extract of *Boerhavia diffusa* L. reduced metastases formation by B167-10 melanoma cells whereas Punarnavine a compound from this plant enhanced immune response against metastatic progress of B16F-10 melanoma cells in mice (Manu and Kuttan, 2007). Clinical trial conducted on 50 patients newly diagnosed with pulmonary tuberculosis showed that clinical recovery rate of patients treated with this plant was faster than in the control (Kant et al., 2001). *Brassica juncea* (L.) Czern. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). This plant has trypsin inhibitory properties (Mandal et al., 2002). Acute dermal toxicity-fixed study showed that leaf extracts of this plant are non-toxic even at concentration of 2000 mg/kg bw topically (Malan et al., 2011). *Buchanania cochinchinensis* (Lour.) M.R.Almeida has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Methanolic seed extracts of this plant are non-toxic to Swiss albino mice and Wistar rats even at dose of 2000 mg/kg bw (Warokar et al., 2010; Singh and Bothara, 2012). In a study by Pattnaik et al. (2013), methanolic root extract of *Buchanania cochinchinensis* (Lour.) M.R.Almeida showed significant wound healing potential. Topical application of *Buchanania cochinchinensis* (Lour.) M.R.Almeida (10% w/w) ointment significantly increased (40.84%) the tensile strength in the incision wound model (Pattnaik et al., 2013). However human clinical trials related to any skin disease have not been conducted on *Boerhavia diffusa* L., *Brassica juncea* (L.) Czern. and *Buchanania cochinchinensis* (Lour.) M.R.Almeida.

Butea monosperma (Lam.) Taub. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). This plant has trypsin inhibitory properties (Pandey and Jamal, 2010). Toxicity study on powder suspension of seeds of this plant on Charles Foster strain albino rats showed that it has toxic effect when administered in powder form (Donga et al., 2011). *Butea monosperma* (Lam.) Taub. is part of topical herbal formulation of The Himalaya Drug Company made of 2 herbs known as “Hair Loss Cream”, which is recommended for preventing hair loss and promoting new growth of hair in men and women. Open, phase IV clinical trial was conducted on 20 patients (15 male and 5 females) age ranging between 21 and 71 years and cream was found to be safe and effective in prevention of hair loss (Ravichandran et al., 2008). *Butea monosperma* (Lam.) Taub. has

antiinflammatory (Shahavi and Desai, 2008) and antifungal (Yadava and Tiwari, 2007) activities which could be reason for its efficacy.

Caesulia axillaris Roxb. has shown antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). In a toxicological study on mice, LD₅₀ value of essential oil of this plant was reported as 9166.6 µl/kg bw (P.K. Mishra et al., 2012). Essential oil of *Caesulia axillaris* Roxb. has exhibited its fungitoxicity against *Aspergillus flavus* at its minimum inhibitory concentration of 1300 mg/L and fungitoxic principle of the oil was standardized as γ-asarone which showed fungitoxicity against the test fungus at 500 mg/L (Varma et al., 2002). However human clinical trial related to any skin disease has not been conducted on this plant. *Callicarpa macrophylla* Vahl has shown antiinflammatory and antimicrobial properties in the earlier studies (Tables 3 and 4). Betulinic acid (BA), a pure compound from *Callicarpa macrophylla* Vahl has been reported to be a selective inducer of apoptosis in tumor cells and it also exhibits antiinflammatory and immunomodulatory properties (Aggarwal et al., 2011). Treatment of cells with Betulinic acid suppressed NF-κB-dependent reporter gene expression and the production of NF-κB-regulated gene products such as COX-2 and MMP-9 induced by inflammatory stimuli (Takada and Aggarwal, 2003). Further pre-clinical, toxicological and clinical studies are required to assess safety and efficacy of this plant in treatment of skin diseases.

Calotropis procera (Aiton) Dryand. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Its milky sap is caustic and irritant on skin and can cause swelling and ulceration (Morton, 1962). Irritant effects were observed in six contact dermatitis patients patch tested with the leaf crushed in normal saline (Singh et al., 1978). Latex of this plant was reported as toxic to black rat (Pahwa and Chatterjee, 1988). In another study it was observed that aqueous extract of root bark of this plant is relatively safe as compared to hydroalcoholic extract of root bark (Ouedraogo et al., 2013). *Calotropis procera* (Aiton) Dryand. along with 21 other plants is part of a polyherbal Ayurvedic formulation of The Himalaya Drug Company known as “Muscle & Joint Rub”. Phase III clinical trial of Muscle & Joint Rub showed that it is effective and safe in the management of muscle sprains, contusions and inflammatory musculoskeletal disorders (Rajanna and Kolhapure, 2005). Anti-inflammatory effect of this plant is known to be caused by inhibiting PGE2 (Arya and Kumar, 2005).

Cannabis sativa L. has shown antiinflammatory, antioxidant, antimicrobial and tyrosinase inhibitory properties in the earlier studies (Tables 3 and 4). Leaf extracts of this plant were non-toxic to female Wistar albino rats even up to dose of 4000 mg/kg bw (Zade et al., 2013), whereas seed oil was toxic to wistar albino rat (Dahab et al., 2013). The dust from the dried fibre may cause irritation and pruritus and its oil can also cause skin infection (Slaviero, 1916; Schwartz et al., 1957). Tetrahydrocannabinol is the compound present in this plant which is toxic to the nervous system (Duke, 2012). There are some biologically active compounds in *Cannabis sativa* L., such as cannabinoids which is antibacterial (Appendino et al., 2008; Radwan et al., 2009) and antifungal (Radwan et al., 2009), cannabinoid ester which is antibacterial as well as antifungal (Ahmed et al., 2008), and cannabidiol has anti-inflammatory properties (BenShabat et al., 2009). *Cassia fistula* L. has shown wound healing, antiinflammatory, antioxidant, antimicrobial and tyrosinase inhibitory properties in the earlier studies (Tables 3 and 4). This plant has trypsin inhibitory properties (Wijaya et al., 2000). Pod pulp, seed extract and stem bark extract of this plant were recorded to be non-toxic to experimental animals (Table 3). A clinical trial was conducted on 81 children aged between 4 and 13 years to assess the laxative effect of *Cassia fistula* L. emulsion (CFE) with mineral oil on

Pediatric functional constipation (FC) and results showed that CFE was most effective in the 3 week treatment of children with FC (Mozaffarpur et al., 2012). Further preclinical, toxicological and clinical studies are required to assess safety and efficacy of this plant in treatment of skin diseases.

Celastrus paniculatus Willd. has shown wound healing, anti-inflammatory and antioxidant properties in the earlier studies (Tables 3 and 4). Seed oil is rubefacient (Nadkarni, 1976). Whole plant and seed extracts of this plant were not toxic even at high doses of 2000 and 5000 mg/kg bw, respectively (Table 3). *Celastrus paniculatus* Willd. along with 21 other plants is part of a polyherbal Ayurvedic formulation of The Himalaya Drug Company known as “Muscle & Joint Rub”. Phase III clinical trial of Muscle & Joint Rub showed that it is effective and safe in the management of muscle sprains, contusions and inflammatory musculoskeletal disorders (Rajanna and Kolhapure, 2005). Since the studied product was a combination of different plants, it is not clear what effect this plant had on the studied disease. Celastrol, a plant-derived triterpene has antioxidant and antiinflammatory activity and in low nanomolar concentrations celastrol was found to suppress the production by human monocytes and macrophages of the pro-inflammatory cytokines TNF- α and IL-1 β (Allison et al., 2001). Additional clinical studies are required before a firm conclusion can be made.

Chrysopogon zizanioides (L.) Roberty has shown antiinflammatory, antioxidant, antimicrobial and tyrosinase inhibitory properties in the earlier studies (Tables 3 and 4). Oil of this plant was found to be non-toxic to albino mice and albino rats with LD₅₀ value of 2985.38 mg/kg bw (Tripathi et al., 2006). Its oil can produce dermatitis in hypersensitive individuals (Greenberg and Lester, 1954). *Chrysopogon zizanioides* (L.) Roberty is also a part of polyherbal formulation of The Himalaya Drug Company made of 16 herbs known as “Anti-Wrinkle cream”, which is used in management of facial skin wrinkles. Prospective, open, non-comparative, phase III clinical trial was conducted on 25 patients of both sexes, aged from 35 to 65 years with wrinkled facial skin and significant improvement in the facial skin wrinkles after a week's application in almost all persons and no clinically significant adverse reactions were observed (Ravichandran et al., 2005). Oil of *Chrysopogon zizanioides* (L.) Roberty is also a part of polyherbal formulation of The Himalaya Drug Company made of oils of 3 plants known as “Baby Powder”, which is used in the management of infantile hyperhidrosis, miliaria rubra, and bad body odor. A prospective clinical trial on 20 infants (birth weight of more than 2500 g) of age 1–12 months suffering from hyperhidrosis, miliaria rubra, and bad body odor showed significant improvement in 3 days time and complete recovery after a week's application confirming that it is clinically effective and safe (Chatterjee et al., 2005a). *Chrysopogon zizanioides* (L.) Roberty oil is also a part of polyherbal formulation of The Himalaya Drug Company made of 5 plants known as “Nourishing Baby Oil”, which is used in infantile dry skin. Prospective, open, phase III clinical trial was conducted on infants between 1 and 12 months of age and was found to be effective and safe in infantile xerosis (Chatterjee et al., 2005c; Parthasarathy et al., 2009). *Chrysopogon zizanioides* (L.) Roberty is part of a polyherbal formulation of The Himalaya Drug Company made of 7 herbs known as “Anti-dandruff Shampoo”, which is recommended for the treatment of dandruff. Prospective, open, non-comparative, phase III clinical trial was conducted on 35 patients suffering from moderate to severe form of dandruff to evaluate the clinical efficacy and safety (short- and long-term) of the shampoo in the management of dandruff (Ravichandran et al., 2004b). The study observed a significant reduction in the mean scores of itching and white scales of dandruff and subjective evaluation revealed remarkable symptomatic and clinical improvement in 2 weeks period

(Ravichandran et al., 2004b). Since all the clinically tested Ayurvedic products were having this plant in combination with other plants it is not clear what effect this plant had on diseases. Additional clinical studies are required before a firm conclusion can be made. The active compounds of *Chrysopogon zizanioides* (L.) Roberty are valencene, 9-octadecenamide, 2,6,10,15,19,23-hexamethyl-2, 6,10,14,18,22-tetracosahexaene, 1,2-benzendicarboxylic acid, di-isooctyl ester and terpenoids such as 3 monoterpenes, 2 sesquiterpenes and 1 triterpene (Huang et al., 2004). According to Chatterjee et al. (2005a, 2005b, 2005c), *Chrysopogon zizanioides* (L.) Roberty is a potent emollient and acts as a moisturizing and soothing agent.

Cissampelos pareira L. has shown antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Hydroethanolic extract of whole plant was non-toxic to rats even at dose of 2 g/kg bw (Amresh et al., 2008). A clinical trial of *Cissampelos pareira* L. showed anti-dengue activity of its extracts (Bhatnagar et al., 2011). Cycleanine, a compound isolated from *Cissampelos pareira* L. has antiinflammatory properties (DNP, 2014). Isoliquiritigenin, a type of natural phenol isolated from *Cleome viscosa* L. is antitumor promoter, antioxidant, antiinflammatory, antimutagenic and antifungal agent (DNP, 2014) and it is under experimentation phase testing for use as a cancer treatment and as an aide for cocaine addiction. *Cleome viscosa* L. has shown wound healing, antiinflammatory, antioxidant, antimicrobial and tyrosinase inhibitory properties in the earlier studies (Tables 3 and 4). Seeds and leaves have rubefacient and vesicant properties (Chopra and Bhadwar, 1940; Behl et al., 1966), whereas bark is irritant and acrid (Nadkarni, 1976). Ethanolic extract of leaves and whole plant was non-toxic to Wistar albino rats even at dose of 2 g/kg bw (Panduraju et al., 2011). Further clinical trials are needed to assess efficacy of *Cissampelos pareira* L. and *Cleome viscosa* L. to treat any skin disease.

Clerodendrum infortunatum L. has shown antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Methanolic extract of leaves of this plant was non-toxic to Swiss albino mice even at dose of 2000 mg/kg bw (Das et al., 2011). A study on anticancer activity of methanol extract of *Clerodendrum infortunatum* L. against Ehrlich's ascites carcinoma (EAC) bearing Swiss albino mice and isolation of bioactive terpenoids from it showed that it has significant anticancer activity, which can be attributed to the presence of oleanolic acid and clerodinin A (Sannigrahi et al., 2012). There is one US patent on this plant, which provides a composition having an extract from *Clerodendrum infortunatum* L., and a method of obtaining the extract, a method of treating cancer by administering a composition having an extract of *Clerodendrum infortunatum* L. to a person in need thereof, a method of activating caspase-3 in a cell by administering a composition having an extract of *Clerodendrum infortunatum* L. to a cell and method of inducing apoptosis in a cell by administering a composition having an extract of *Clerodendrum infortunatum* L. to a cell (Giri et al., 2014). *Commelina benghalensis* L. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Hydroethanolic extract of leaves of this plant was non-toxic to Wistar albino female rats even at dose of 2000 mg/kg bw (Tiwari et al., 2013). Human clinical trial related to any skin disease has not been conducted on *Clerodendrum infortunatum* L. and *Commelina benghalensis* L.

Crotalaria juncea L. has shown antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Ethanolic extract of leaves of this plant was non-toxic to male albino Wistar rats even at dose of 2000 mg/kg bw (Ashok et al., 2006). *Cryptolepis dubia* (Burm.f.) M.R.Almeida has shown antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Whole plant and leaf extracts of this plant

were non-toxic to mice, rat and rabbit (Minh and Tuan, 2013). *Curculigo orchioideis* Gaertn. has shown wound healing, anti-inflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Hydroalcoholic extract of rhizome of this plant was non-toxic to Wistar rats even at dose of 2000 mg/kg bw (Asif and Kumar, 2010). According to Chinese Pharmacopoeia (2010), *Curculigo orchioideis* Gaertn. has a certain degree of toxicity, and the clinical dosage recommended for adults is 3–9 g daily. Curculigoside treated human umbilical vein endothelial cell injury induced by H₂O₂ in *in vitro* study with active concentrations of 0.5, 5 and 10 μ mol/ml (Wang et al., 2010). However human clinical trial related to any skin disease has not been conducted on *Crotalaria juncea* L., *Cryptolepis dubia* (Burm.f.) M.R.Almeida and *Curculigo orchioideis* Gaertn.

Cuscuta reflexa Roxb. has shown anti-inflammatory, antioxidant, antimicrobial and tyrosinase inhibitory properties in the earlier studies (Tables 3 and 4). Aqueous and alcoholic extract of stem of this plant was non-toxic to albino mice of both sex even at dose of 2000 mg/kg bw (Katiyar et al., 2012). *Cynodon dactylon* (L.) Pers. has shown wound healing, anti-inflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Chloroform and ethanolic extracts of whole plant was non-toxic to mice even at dose of 2000 mg/kg bw (Yogesh et al., 2013). *Dalbergia sissoo* DC. has shown anti-inflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Alcoholic extract of bark of this plant was non-toxic to Wistar rats even at dose of 3000 mg/kg bw (Mojahid-ul-Islam and Elhddad, 2012). Alcohol extract of aerial parts showed dose dependent inhibitory effect on the motility of isolated rabbit duodenum, pronounced bronchodilation, as well as significant anti-inflammatory, antipyretic, analgesic and estrogen-like activities (Sarg et al., 1999). However human clinical trial related to any skin disease has not been conducted on *Cuscuta reflexa* Roxb. and *Dalbergia sissoo* DC.

Datura stramonium L. has shown anti-inflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). *Datura stramonium* L. poisoning is very common in India usually involving seeds (Gaire and Subedi, 2013) and symptoms of poisoning include agitation, seizures, blurred vision, dry mouth, delirium, mydriasis, photophobia, extreme thirst, tachycardia, nausea, vomiting, decreased bowel sounds, difficulty in swallowing and speaking, hyperthermia, hypertension, loss of consciousness and coma (Oberndorfer et al., 2002). As per study conducted on male rats by Bouzidi et al. (2011), a single dose of acute toxicity of 100 mg/kg of *Datura stramonium* L. includes decrease in the weight of the liver, spleen and brain and significant increases in the levels of red blood cells (RBC), hematocrit (HCT), hemoglobin (HGB), and white blood cells (WBC). *Dendrophthoe falcata* (L.f.) Ettingsh. has shown anti-inflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Ethanolic extract of leaves of this plant showed low level of general toxicity with LC₅₀ value of 100 μ g/ml in Brine shrimp lethality assay for general toxicity on Swiss albino mice of either sex (Hasan et al., 2006). However human clinical trial related to any skin disease has not been conducted on *Datura stramonium* L. and *Dendrophthoe falcata* (L.f.) Ettingsh.

Desmodium gangeticum (L.) DC. has shown anti-inflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Extracts of aerial parts of this plant were non-toxic to albino Wistar rats even at dose of 2000 mg/kg bw (Bisht and Bhattacharya, 2013). *Desmostachya bipinnata* (L.) Stapf has shown anti-inflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Alcoholic and aqueous extracts of roots of this plant were non-toxic to female albino mice even at dose of 2000 mg/kg bw (Hegde et al., 2010). A clinical study of *Desmostachya bipinnata* (L.) Stapf along with *Imperata*

cylindrica Beauv., which are part of Trinapanchamoola, a well known diuretic was conducted on 29 healthy volunteers and results showed that use of these herbs led to a percentage increase in urine volume as compared to placebo group (N.T. Shah et al., 2012). Different concentrations of 70% methanolic extract of the roots of this plant have shown effective antioxidant activities in different reactive oxygen species scavenging assays including DPPH, nitric oxide, hydrogen peroxide and hydroxyl radical scavenging activities (Rahate et al., 2012). Besides that *in-vitro* cytotoxic study on the human cervical cancer cell lines (HeLa), human laryngeal epithelial carcinoma cells (Hep-2) and BIH 3T3 using 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) showed inhibition in concentration dependent manner of range between 25 and 400 μ g/ml on all the cell lines (Rahate et al., 2012). *Dicliptera paniculata* (Forssk.) I.Darbysh. has shown anti-inflammatory and antimicrobial properties in the earlier studies (Tables 3 and 4). Aqueous, methanolic and butanolic extracts of leaves and stem of this plant were non-toxic to Wistar rats even at dose of 5000 mg/kg bw (Abdulazeez et al., 2010). However human clinical trial related to any skin disease has not been conducted on *Desmodium gangeticum* (L.) DC., *Desmostachya bipinnata* (L.) Stapf and *Dicliptera paniculata* (Forssk.) I.Darbysh.

Eclipta prostrata (L.) L. has shown anti-inflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Ethanolic extract of whole plant was non-toxic to Swiss albino mice and Wistar albino rats even at dose of 4 g/kg bw (Md.A. Rahman et al., 2012). The active compounds of *Eclipta prostrata* (L.) L. are triterpenoid glucosides such as daucosterol, stigmaterol-3-O-glucoside and ecliptasaponin C (Zhang and Chen, 1996), which could be involved in beneficial effects of extract of this plant on skin diseases. Immunomodulatory effect of daucosterol, a beta-sitosterol glycoside, against disseminated candidiasis caused by *Candida albicans* were studied and it was showed that it protects mice against disseminated candidiasis by the CD4+ Th1 immune response (Lee et al., 1997). *Eclipta prostrata* (L.) L. also has potent anti-inflammatory (Ferreira et al., 2000), analgesic (Sawant et al., 2004) and immunomodulatory activities (Jaythirtha and Mishra, 2004). Its leaves contain α -terthienyl which is phototoxic to human skin (Gommers and Voor in't Holt, 1976). *Eclipta prostrata* (L.) L. is part of a polyherbal formulation of The Himalaya Drug Company made of 5 herbs known as "Anti-dandruff Hair Oil", which is recommended for the treatment of dandruff. Open, non-comparative, non-randomized, phase III clinical trial was done on 25 patients of both sexes, from the age group of 20–45 years to evaluate the clinical efficacy and safety (short- and long-term) of this oil in the management of dandruff (Vyjayanthi et al., 2004). The study showed that there was highly significant reduction in the mean score for itching and white scales at the end of 2 weeks and oil is effective and safe in the management of dandruff (Vyjayanthi et al., 2004). Further clinical studies are required to assess efficacy of this plant in treatment of skin diseases.

Ehretia laevis Roxb. has shown antioxidant properties in the earlier studies (Table 3). Further preclinical, toxicological and clinical studies are required to assess safety and efficacy of *Ehretia laevis* Roxb. and *Eulaliopsis binata* (Retz.) C.E.Hubb. in treatment of skin diseases. *Euphorbia hirta* L. has shown wound healing, anti-inflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Its sap is described as irritant (Chopra and Bhadwar, 1940; Souder, 1963; Behl et al., 1966). Patch test of crushed leaves of this plant in small quantity of normal saline elicited positive reaction in 2 out of 18 contact dermatitis patients (Singh et al., 1978). Aqueous and ethanolic extracts of fruits suspended in 0.5% w/v Sodium-CMC were found to be toxic to adult Swiss albino mice even at low doses and mortality was observed at high doses of 2500 and 3000 mg/kg bw (P. Das et al., 2010). However human clinical trial related to any skin disease has

not been conducted on this plant. *Euphorbia thymifolia* L. has shown antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Its latex is vesicant to the skin of many individuals and can cause dermatitis (Pammel, 1911; Souder, 1963). Alcohol, chloroform and aqueous extracts of whole plant were non-toxic to albino mice even at dose of 2000 mg/kg bw (Mamatha et al., 2014). Clinical trials with triturated fresh leaves of this plant have proved its efficacy in paronychia (common nail bed infection) when applied at early stage (Mali and Panchal, 2013). Further clinical studies are required to assess efficacy of this plant in treatment of skin diseases.

Ficus benghalensis L. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Ethanol and aqueous extracts of bark of this plant were non-toxic to female albino Wistar rats even at dose of 2000 mg/kg bw (Garg and Paliwal, 2011). This plant is a part of a polyherbal Ayurvedic formulation of The Himalaya Drug Company made of 7 herbs known as “Anti-dandruff Shampoo”, which is recommended for the treatment of dandruff. Prospective, open, non-comparative, phase III clinical trial was conducted on 35 patients suffering from moderate to severe form of dandruff to evaluate the clinical efficacy and safety (short- and long-term) of the shampoo in the management of dandruff (Ravichandran et al., 2004b). The study observed a significant reduction in the mean scores of itching and white scales of dandruff and subjective evaluation revealed remarkable symptomatic and clinical improvement in 2 weeks period (Ravichandran et al., 2004b). The principal compounds of *Ficus benghalensis* L. are ketones such as tetratriacontene, 6-heptatriacontene, pentatriacontan, beta-sitosterol- α -D-glucose and meso-inositol (Subramanian and Misra, 1978). *Ficus racemosa* L. has shown wound healing, antiinflammatory, antioxidant, antimicrobial and tyrosinase inhibitory properties in the earlier studies (Tables 3 and 4). Methanolic extract of bark of this plant was non-toxic to female Wistar rats even at dose of 2000 mg/kg bw (Choudhury and Jadhav, 2013). A clinical trial on leaves of *Ficus racemosa* L. showed antihyperglycemic action showing support for the treatment of Type 2 Diabetes Mellitus in traditional system of medicines (Urooj and Ahmed, 2013). Efficacy of a proprietary herbal preparation consisting of *Ficus racemosa* L. and 5 other plants was evaluated on 28 patients of persistent post prandial hyperglycemia, which showed persistent fall in fasting and post prandial blood glucose levels (Dubey et al., 1993). A clinical trial of polyherbal ointment of which *Ficus racemosa* L. was one of the constituents on 15 burn patients showed that ointment is highly efficacious in controlling *Candida albicans* infection and helped in quicker epithelialization and burns were completely healed in 8–26 days of treatment (Bhatt and Kora, 1984). Racemosic acid, a compound from *Ficus racemosa* L. showed potent inhibitory activity against COX-1 and 5-LOX *in vitro* with IC₅₀ values of 90 and 18 μ M, respectively (Li et al., 2004). Cytotoxic effects of the extracts of *Ficus racemosa* L. were investigated *in vitro* using the ATP-based luminescence assay and results showed no cytotoxicity on the cell lines skin fibroblasts (1BR3), human Caucasian hepatocyte carcinoma (Hep G2) and human Caucasian promyelocytic leukaemia (HL-60) (Li et al., 2004).

Ficus religiosa L. has shown wound healing, antiinflammatory, antioxidant, antimicrobial and tyrosinase inhibitory properties in the earlier studies (Tables 3 and 4). Methanolic extract of bark of this plant was non-toxic to Swiss albino male mice at dose of 500 mg/kg bw (Sreelekshmi et al., 2007). A clinical study of Ashvattha Kshirpaka prepared with 10 g powder of root bark, stem bark, fruit and tender leaf buds of *Ficus religiosa* L. on 44 patients erectile dysfunction showed encouraging results (Virani et al., 2010). Bergaptol a compound isolated from *Ficus religiosa* L. has antiinflammatory and antifungal properties (DNP, 2014). *Holarrhena pubescens* Wall. ex G.Don has shown antiinflammatory,

antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Stem bark of *Holarrhena pubescens* Wall. ex G.Don is part of an Ayurvedic drug known as Sunder Vati used for treatment of acene vulgaris. *Holarrhena pubescens* Wall. ex G.Don has astringent activity (Nadkarni, 1976). Ethanolic extract of seed of this plant was non-toxic to female Wistar rats even at dose of 2000 mg/kg bw (Saha and Subrahmanyam, 2013). Randomised, double-blind, placebo-controlled clinical trial of this drug was done on 42 male and 42 female patients between 18 and 28 years and Sunder Vati was found to be orally effective in the treatment of acne vulgaris (Pranipe and Kulkarni, 1995). A clinical study on stem bark extracts of *Holarrhena pubescens* Wall. ex G.Don in the form of *Kutaja tvak churna* showed healing activity in patients suffering from bleeding piles (Pal et al., 2009). In a clinical study it was observed that a daily intake of the bark powder for 15 days completely cured patients suffering from amebiasis (Shahabuddin et al., 2006).

Holoptelea integrifolia Planch. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Methanol and petroleum ether extracts of leaves of this plant were non-toxic to albino mice of either sex even at dose of 2000 mg/kg bw (Sutar et al., 2014). *Hyptis suaveolens* (L.) Poit. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). This plant also has trypsin inhibitory properties (Aguirre et al., 2004). Extracts of leaves of this plant were non-toxic to Swiss albino mice of either sex and LD₅₀ was found to be more than 5000 mg/kg bw (Shenoy et al., 2009b). *Ipomoea carnea* Jacq. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Aqueous extract of leaves of this plant showed toxicity symptoms to Wistar albino rats at dose of 2000 mg/kg bw (Khalid et al., 2011). Preliminary clinical trial covering 195 patients of three age groups (70 patients for 15–25 years age, 80 patients of 26–35 years age and 45 patients of 36–50 years age) have confirmed the efficacy of this plant in curing vaginal and cervical leucorrhea, with a success rate of about 76% (Atique et al., 2009). However human clinical trial related to any skin disease has not been conducted on *Holoptelea integrifolia* Planch., *Hyptis suaveolens* (L.) Poit. and *Ipomoea carnea* Jacq.

Lansea coromandelica (Houtt.) Merr. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Methanolic extract of bark of this plant was non-toxic to mice even at dose of 1000 mg/kg bw (Majumder et al., 2013). *Lantana camara* L. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Contact with this plant can cause dermatitis to some susceptible individuals (Morton, 1962). Methanolic extract of leaves of this plant was non-toxic to adult mice even at dose of 2 g/kg bw (Pour et al., 2011). *Lawsonia inermis* L. has shown wound healing, antiinflammatory, antioxidant, antimicrobial and tyrosinase inhibitory properties in the earlier studies (Tables 3 and 4). This plant also has trypsin inhibitory properties (Yogisha et al., 2002). Ethanolic extract of leaves of this plant was toxic to Wistar albino rats of either sex with toxicity above dose of 1000 mg/kg bw and lethal dose at 2000 mg/kg bw (Kaur et al., 2014). *Leonotis nepetifolia* (L.) R.Br. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Trichomes of leaves can cause dermatitis in susceptible individuals (Gowanloch and Brown, 1943). Methanolic extract of stem bark of this plant was non-toxic to Swiss albino mice with LD₅₀ value of 3807.9 mg/kg bw (Ayanwuyi et al., 2009). However human clinical trial related to any skin disease has not been conducted on *Lansea coromandelica* (Houtt.) Merr., *Lantana camara* L., *Lawsonia inermis* L. and *Leonotis nepetifolia* (L.) R.Br.

Linum usitatissimum L. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). This plant also has trypsin inhibitory properties (Cierpicki and Otlewski, 2000). Its oil can produce a follicular eruption (Barnes, 1931) and can be irritant and occasionally sensitizing (Schwartz et al., 1957). Fixed seed oil of this plant was non-toxic to Swiss albino mice of either sex even at dose of 20 ml/kg bw (Kaithwas et al., 2011). It has been found that *Linum usitatissimum* L. seed lignan reduces mammary tumor growth in the later stages of carcinogenesis (Thompson et al., 1996). According to Bommareddy et al. (2009), flaxseed meal and oil are effective chemo-preventive agents. Daily consumption of 50 g flaxseed for four weeks lowers the LDL cholesterol by 8% in young healthy adults (Cunnane et al., 1995). Basing on human trials before 2011 Health Canada's Food Directorate concluded that a claim linking consumption of ground whole flaxseed and blood cholesterol lowering was warranted (Health Canada, 2014; Shim et al., 2014).

Litsea glutinosa (Lour.) C.B.Rob. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Aqueous extract of whole plant was non-toxic to Wistar rats even at dose of 2000 mg/kg bw (Devi and Meera, 2010). *Mallotus philippensis* (Lam.) Mull.Arg. has shown antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Methanolic extract of leaves of this plant was non-toxic to female mice even at dose of 2000 mg/kg bw (Babu et al., 2010). Rottlerin or mallotoxin, a principal phloroglucinol constituent of this plant appears to have great potential for being used in chemotherapy because it affects several cell machineries involved in survival, apoptosis, autophagy and invasion (Maioli et al., 2012). A clinical trial on 40 patients exhibiting symptoms of parasitic infection was undertaken to assess efficacy of this plant on intestinal parasites and results showed symptomatic relief without any adverse effect (Bharadwaj et al., 2013). *Martynia annua* L. has shown antiinflammatory and antimicrobial properties in the earlier studies (Tables 3 and 4). Methanolic extract of leaves of this plant was non-toxic to Wistar albino rats even at dose of 2000 mg/kg bw (Babu et al., 2010). *Mirabilis jalapa* L. has shown antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Hydroethanolic extract of flowers of this plant was non-toxic to female rat even at dose of 2000 mg/kg bw (Augustine et al., 2013). There are trypsin inhibitors present in the seeds of *Mirabilis jalapa* L. (Kowalska et al., 2007). Human clinical trial related to any skin disease has not been conducted on *Litsea glutinosa* (Lour.) C.B.Rob., *Mallotus philippensis* (Lam.) Mull.Arg., *Martynia annua* L., *Milletia extensa* (Benth.) Baker. and *Mirabilis jalapa* L. Pharmacological, toxicological, preclinical and clinical studies are required on *Milletia extensa* (Benth.) Baker to prove its efficacy in treatment of skin diseases.

Mitragyna parvifolia (Roxb.) Korth. has shown antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Ethanolic extract of leaves of this plant was non-toxic to Swiss albino mice even at dose of 1500 mg/kg bw (Kaushik et al., 2009a). *Momordica dioica* Roxb. ex Willd. has shown antiinflammatory, antioxidant, antimicrobial and tyrosinase inhibitory properties in the earlier studies (Tables 3 and 4). Aqueous extract of fruits of this plant was non-toxic to Wistar rats even at dose of 15 times more than effective dose i.e., 200 mg/kg bw (R. Singh et al., 2011). *Mucuna pruriens* (L.) DC. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). This plant also has trypsin inhibitory properties (Prakash et al., 2001). Trichomes on pods of this plant can penetrate skin and cause itching and irritation that may last for some time (Allen, 1943). Active pruritogenic principle in this plant is a proteolytic enzyme, mucunain (Arthur and Shelly, 1955; Shelly and Arthur, 1955). Methanolic extract of seeds of this plant was non-toxic to

albino mice of both sexes even at dose of 4000 mg/kg bw (Manalisha and Chandra, 2012). Randomised, controlled, double blind crossover clinical trial of eight Parkinson's disease patients with a short duration L-dopa response and on period dyskinesia was conducted to assess clinical effects and levodopa (L-dopa) pharmacokinetics following two different doses of *Mucuna pruriens* (L.) DC. preparation (Katzenschlanger et al., 2004). The results showed the rapid onset of action and longer on time without concomitant increase in dyskinesias on mucuna seed powder formulation suggest that this natural source of L-dopa might possess advantages over conventional L-dopa preparations in the long term management of Parkinson's disease (Katzenschlanger et al., 2004). *Nicotiana plumbaginifolia* Viv. has shown antimicrobial properties in the earlier studies (Table 3). However human clinical trial related to any skin disease has not been conducted on *Mitragyna parvifolia* (Roxb.) Korth., *Momordica dioica* Roxb. ex Willd., *Mucuna pruriens* (L.) DC. and *Nicotiana plumbaginifolia* Viv. Pharmacological, toxicological, preclinical and clinical studies are required on *Nicotiana plumbaginifolia* Viv. to prove its efficacy in treatment of skin diseases.

Oroxylum indicum (L.) Kurz has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Ethanolic extract of roots of this plant was non-toxic to Wistar male albino rats even at dose of 5000 mg/kg bw (Tamboli et al., 2011). Baicalein from *Oroxylum indicum* (L.) Kurz showed anti-cancer potential in leukemia cells through inducing cell cycle arrest and apoptosis (Roy et al., 2007). Two compounds Chrysin and Oroxylin A isolated from *Oroxylum indicum* (L.) Kurz have antiinflammatory properties (DNP, 2014). *Persicaria barbata* (L.) H.Hara has shown antiinflammatory properties in the earlier studies (Tables 3 and 4). Alcoholic extract of leaves of this plant was non-toxic to Wistar albino rats even at dose of 2000 mg/kg bw (Sheela et al., 2011). However clinical trials related to any skin disease has not been conducted on *Oroxylum indicum* (L.) Kurz and *Persicaria barbata* (L.) H.Hara.

Phyllanthus amarus Schumach. and Thonn. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Aqueous extract of leaves of this plant was non-toxic to female rats even at dose of 5000 mg/kg bw (Asare et al., 2011). Extract from *Phyllanthus amarus* Schumach. and Thonn. and some purified lignans from this plant such as niranthin, nirtetralin and phyltetralin exhibit *in vivo* and *in vitro* anti-inflammatory properties, which are probably mediated through its direct ability to interact with platelet activating factor receptor binding sites (Kassuya et al., 2006). A clinical trial on 50 patients between 25 and 60 years with hepatitis-C suggested that therapy with *Phyllanthus amarus* Schumach. and Thonn. increased antioxidants and reduced lipid peroxidation of hepatic cellular and intracellular membranes and protected liver damage due to free radicals in hepatitis-C (Nikam et al., 2011). Another clinical trial showed that *Phyllanthus amarus* Schumach. and Thonn. has inhibitory effects on HIV *in vitro* and *in vivo* (Notka et al., 2004). Owing to the impressive preclinical therapeutic potential, the plant extracts have been evaluated in human trial for the treatment of HIV, jaundice, hypertension and diabetes (Patel et al., 2011) where they have show promising properties. However further clinical studies are required to prove its efficacy in treatment of skin diseases.

Pongamia pinnata (L.) Pierre has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Crude extract of seeds of this plant was non-toxic to Wistar female albino rats even at dose of 2000 mg/kg bw (Aneela et al., 2011). *Pongamia pinnata* (L.) Pierre is part of polyherbal Ayurvedic formulation of The Himalaya Drug Company made of 8 herbs known as "Anti-dandruff Hair cream". The open, non-comparative, phase III clinical trial of this cream was done on 50 patients to evaluate the clinical efficacy and safety (short- and

long-term) of this cream in the management of dandruff (Agarwal et al., 2009). The study observed significant symptomatic and clinical improvement of dandruff in 6 weeks and concluded that this cream is effective and safe in the management of dandruff (Agarwal et al., 2009). *Ranunculus sceleratus* L. has shown anti-inflammatory and antimicrobial properties in the earlier studies (Tables 3 and 4). Fresh plant is intensely irritating and can produce violent blistering, particularly of the lips and tongue, and also of skin (Weber, 1937; Chopra and Bhadwar, 1940). Chloroform, methanol and water extracts of leaves of *Pongamia pinnata* (L.) Pierre and *Ranunculus sceleratus* L. were evaluated for anti-ringingworm activity against five strains of dermatophytes viz., *Trichophyton rubrum*, *Trichophyton mentagrophytes*, *Trichophyton tonsurans*, *Microsporum gypseum* and *Microsporum fulvum* and chloroform extracts were found to be highly active against all the tested fungi (Sharma et al., 2012a). *Premna mollissima* Roth has shown anti-inflammatory properties in the earlier studies (Table 4). Methanolic extract of leaves of this plant was non-toxic to albino mice of either sex even at dose of 5000 mg/kg bw (Mahire et al., 2009). However clinical trials related to any skin disease have not been conducted on *Ranunculus sceleratus* L. and *Premna mollissima* Roth.

Rauvolfia serpentina (L.) Benth. ex Kurz has shown antimicrobial properties in the earlier studies (Table 4). Methanolic extract of root of this plant was toxic at high doses to male Wistar albino mice and LD₅₀ value was determined as 141.25 mg/kg bw (Azmi and Qureshi, 2012). It is one of the important ingredients of Ayurvedic formulation known as “Divya Muka Vati”, which is used to cure high blood pressure. In a double-blind placebo-controlled investigation, 14 healthy adult male volunteers were studied to assess and compare the urinary effects of acute single doses of two antihypertensive formulations containing *Rauvolfia serpentina* (L.) Benth. ex Kurz and it was observed that even a single dose of these formulations is effective (Leary et al., 1986). Moreover reserpine has shown significant anti-inflammatory activity in animal models (Mokhtarian and Griffin, 1984; Lam and Ferrell, 1991).

Ricinus communis L. has shown anti-inflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Castor oil is said to produce contact dermatitis to some susceptible individual (Karabow, 1927; Coca et al., 1931). Methanolic and aqueous extracts of roots of this plant were non-toxic to Wistar albino rats even at dose of 2000 mg/kg bw (Ilavarasan et al., 2011). In a study on guinea-pig eyelid, ricinolein a compound from this plant showed both pro-inflammatory and anti-inflammatory properties that were observed upon acute and repeated application of this compound, respectively (Vieira et al., 2001). *Ricinus communis* L. along with 21 other plants is part of a polyherbal Ayurvedic formulation of The Himalaya Drug Company known as “Muscle & Joint Rub”. Phase III clinical trial of Muscle & Joint Rub showed that it is effective and safe in the management of muscle sprains, contusions and inflammatory musculoskeletal disorders (Rajanna and Kolhapure, 2005).

Scoparia dulcis L. has shown wound healing, anti-inflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Methanolic extract of whole plant was non-toxic to mice with LD₅₀ value of 3807 mg/kg bw (Abdulsalaam et al., 2013). Nishino et al. (1993), reported that Scopadulcic acid B (SDB), a tetracyclic diterpenoid isolated from this plant inhibited the effects of tumor promoter 12-O-tetradecanoylphorbol-13-acetate *in vitro* and *in vivo*, and potency of Scopadulcic acid B was proved to be stronger than that of other natural antitumor-promoting terpenoids, such as glycyrrhetic acid. SDB has shown antiviral activity against Herpes simplex virus type 1, antitumor activity in various human cell lines and direct inhibitory activity against porcine gastric (H+), K(+)-ATPase (Riel et al., 2002). *Semecarpus anacardium* L.f. has shown anti-inflammatory, antioxidant and

antimicrobial properties in the earlier studies (Tables 3 and 4). Dermatitis may be caused to some susceptible individuals by contact with timber of this plant (Hausen, 1973). Milk preparation of nuts of this plant was non-toxic to Wistar male albino rats even at dose of 2000 mg/kg bw (Vijayalakshmi et al., 2000). 3-Pentadecyl-1,2-benzenediol isolated from this plant is antibacterial agent and cytotoxic to human cancer lines (DNP, 2014). A variety of nut extract preparations of this plant are effective against many diseases, viz., arthritis, tumors, infections etc. and are non-toxic even at high dose of 2000 mg/kg (Premalatha, 2000).

Senna tora (L.) Roxb. has shown wound healing, anti-inflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). This plant also has trypsin inhibitory properties (Tripathi et al., 2011). Ethanolic extract of leaves of this plant was non-toxic to Swiss albino mice of both sexes even at a high dose of 10,000 mg/kg bw (Ambali et al., 2005). Antisporiatic activity of three flavonoids, namely luteolin-7-O- β -glucopyranoside, quercetin-3-O- β -D-glucuronide and formononetin-7-O- β -D-glucoside isolated from the ethanolic extract of leaves were investigated using UV-B induced photodermatitis model on healthy male Wistar rats (120–170 g) and Swiss albino mice (25–30 g) by Vijayalakshmi and Geetha (2014). The ethanolic extract (400 mg/kg) and three flavonoids exhibited a significant percentage reduction of relative epidermal thickness when compared with a positive control, and study concluded that the flavonoids from *Senna tora* (L.) Roxb. leaves have significant antipsoriatic activity (Vijayalakshmi and Geetha, 2014).

Sesamum indicum L. has shown anti-inflammatory, antioxidant and tyrosinase inhibitory properties in the earlier studies (Tables 3 and 4). Seed oil of this plant was non-toxic up to oral dose of 3 g/kg bw to Swiss albino mice (Ambali et al., 2005). The randomized clinical trial on 41 patients with type 2 diabetes showed that *Sesamum indicum* L. seeds have favorable effects in decreasing cardiovascular disease risk factors in Type 2 diabetic patients (Mirmiran et al., 2013), whereas a controlled open-label clinical trial on effect of defatted *Sesamum indicum* L. seeds on type 2 diabetic women has showed glycemic control and weight in diabetic women by use of defatted *Sesamum indicum* L. seeds (Figueiredo, Modesto-Filho, 2008). *Sesamum indicum* L. is also a part of polyherbal Ayurvedic formulation of The Himalaya Drug Company made of 8 herbs known as “Anti-dandruff Hair cream”. The open, non-comparative, phase III clinical trial of this cream was done on 50 patients to evaluate the clinical efficacy and safety (short- and long-term) of this cream in the management of dandruff (Agarwal et al., 2009). The study observed significant symptomatic and clinical improvement of dandruff in 6 weeks and concluded that this cream is effective and safe in the management of dandruff (Agarwal et al., 2009). *Sesamum indicum* L. is rich source of linolenic acid, sesamol (475 mg/100 g), a potent natural antioxidant, and alpha-tocopherol (32 mg/100 g), the most active form of vitamin E (Hahn et al., 2009), which may be main reason for the beneficial effect of oil of this plant on treating skin diseases.

Shorea robusta Gaertn. has shown wound healing, anti-inflammatory and antimicrobial properties in the earlier studies (Tables 3 and 4). Aqueous and methanolic extracts of leaves of this plant were non-toxic to Swiss albino mice and male wistar rats with orally fed LD₅₀ values of 2.4 g/kg and 2.7 g/kg bw, respectively (Chattopadhyay et al., 2012). A study by Chattopadhyay et al. (2012) showed that aqueous extract of leaves of this plant at 40 μ g/ml significantly inhibited NO₂, and release of prostaglandin E₂, tumor necrosis factor- α , interleukins-1 β and interleukin-6 from lipopolysaccharide-stimulated human monocytic cell lines suggesting mechanism of action of anti-inflammatory property of this plant. It is a part of polyherbal formulation of The Himalaya Drug Company made of 4 herbs known as “FootCare Cream”, which is used in management of facial skin wrinkles. Prospective, open, non-comparative, phase III clinical trial

was conducted on 50 patients with mean age of 44.26 years and significant reduction in number of cracks and other symptoms such as scaling, pruritus, wrinkles, pain, pigmentation and laxity of skin was seen (Dange and Grandhi, 2009). Since the studied product was polyherbal formulation, it is not clear what effect exactly *Shorea robusta* Gaertn. had on the studied disease. Alopecia affects 20% of men and has an androgenic etiology in 95% of cases, i.e. high testosterone concentration and a high rate of conversion to DHT by 5- α -reductase (Mas-Chamberlin et al., 2005). Oleanolic acid is the prime alkaloids present in *Shorea robusta* Gaertn. (Dange and Grandhi, 2009). The oleanolic acid is shown to inhibit 5- α -reductase with a dose effect enabling inactivation of 54% of testosterone conversion to dihydrotestosterone at a concentration as low as 9 ppm (Mas-Chamberlin et al., 2005). Chebulinic acid present in seed of this plant is antioxidant compound (DNP, 2014). Further clinical trials are needed to assess efficacy of this plant in the treatment of skin diseases.

Sida cordata (Burm. F.) Borss. Waalk. has shown antioxidant properties in the earlier studies (Table 4). *Sida rhombifolia* L. has shown antiinflammatory and antioxidant properties in the earlier studies (Tables 3 and 4). Aqueous root extract of this plant was non-toxic even up to dose of 5000 mg/kg bw in acute oral toxicity test conducted on Sprague-Dawley rats (Indhumathi et al., 2014). *Solanum incanum* L. roots are pounded and applied topically on leucoderma by the Tharu community. Ethanolic fruit extracts of this plant has not shown toxicity even up to dose of 2000 mg/kg bw in acute oral toxicity test on female swiss albino mice (Indhumathi et al., 2014). A study by Wu et al. (2011) on topical treatment of SR-T100 extracted from *Solanum incanum* L. on UVB-induced cutaneous squamous cell carcinoma (SCC) of hairless mice and actinic keratoses of human, indicated that SR-T100 induces apoptosis of SCC cells via death receptors and the mitochondrial death pathway. The high efficiency of SR-T100 in this preclinical trial suggested that SR-T100 is a highly promising herb for actinic keratoses of human and related disorders (Wu et al., 2011). Clinical trials related to any skin disease have not been conducted on *Sida cordata* (Burm. F.) Borss. Waalk, *Sida rhombifolia* L. and *Solanum incanum* L., and there is need for further preclinical, toxicological and clinical studies on these plants to prove scientific basis for use of these plants in treatment of skin diseases.

Terminalia arjuna (Roxb. ex DC.) Wight & Arn. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). *Terminalia arjuna* (Roxb. ex DC.) Wight & Arn. also has trypsin inhibitory properties (Rai et al., 2008). In a study on male swiss albino mice, methanolic extract of leaves of this plant was found to be non-toxic with LD₅₀ value of 900 mg/kg bw (Biswas et al., 2011a). Several clinical studies have assessed efficacy of *Terminalia arjuna* (Roxb. ex DC.) Wight & Arn. in cardiac disorders (Paarakh, 2010). In a clinical study this plant has shown to decrease platelet activation which shows its antithrombotic properties (Malik et al., 2009). In a clinical study on 20 patients of angiographically proven coronary artery diseases, ethanolic bark extract of *Terminalia arjuna* (Roxb. ex DC.) Wight & Arn. decreased platelet activation suggesting that it possess antithrombotic properties (Malik et al., 2009). *Terminalia arjuna* (Roxb. ex DC.) Wight & Arn. along with 6 other herbs is a part of Ayurvedic formulations which were tested for treating acne vulgaris (Lalla et al., 2001). A Randomized, double-blind, placebo-controlled Phase II clinical trials were conducted on 53 patients for 4 weeks to test efficacy of these formulations and it was observed that combination of use of internal and external preparations showed better efficacy as compared to the used of oral formulations alone (Lalla et al., 2001). Since the product studied was a combination of different plants, it is not clear what exact effect *Terminalia arjuna* (Roxb. ex DC.) Wight & Arn. had on the studied disease but positive effect of the studied formulation could be due to synergistic effect of all the 6 herbs used in it. Although

preclinical trials have shown this plant is promising candidate for development of skin care products but additional human clinical trials are required before moving any further in that direction.

Tridax procumbens (L.) L. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). The methanol extract of this plant exhibited high antifungal activity against clinically important human skin pathogens such as *Microsporum fulvum*, *Microsporum gypseum*, *Trichophyton mentagrophytes*, *Trichophyton rubrum*, *Trichosporon beigelii* and *Candida albicans* with low MIC values and GC-MS analysis revealed the presence of 9,12-octadecadienoic acid ethyl ester, 5 α -cholestane, hexadecanoic acid ethyl ester and 9-octadecenoic acid ethyl ester as major constituents (Policegoudra et al., 2014). In a study on Swiss albino mice and albino rats, ethyl acetate extract of leaves of this plant showed no toxicity with LD₅₀ value of 2100 mg/kg bw (Abubakar et al., 2012). *Typha domingensis* Pers. has shown antiinflammatory and antioxidant properties in the earlier studies (Tables 3 and 4). *Vallisneria spiralis* (Roth) Kuntze has shown antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Root bark oil extract of this plant was found to be non-toxic to albino rats at concentration of 5 ml/kg bw (Punam et al., 2012). Clinical trials related to any skin disease have not been conducted on *Typha domingensis* Pers. and *Vallisneria spiralis* (Roth) Kuntze, and further toxicological, preclinical and chemical studies are required on these plants to identify active compounds in these plants and their efficacy in treatment of skin diseases.

Vanda tessellata (Roxb.) Hook. ex G.Don. has shown antiinflammatory and antimicrobial properties in the earlier studies (Tables 3 and 4). Alcoholic and aqueous extracts of whole plant were found to be non-toxic even up to 2 g/kg bw in a study on Swiss mice (Kumar et al., 2000). Topical application of extract of *Vanda tessellata* (Roxb.) Hook. ex G.Don. to rats at a dose of 150 mg/kg/day for 10 days showed wound healing properties in excision wound model (Nayak et al., 2005). *Vanda tessellata* (Roxb.) Hook. ex G.Don. along with 21 other plants is part of a polyherbal Ayurvedic formulation of The Himalaya Drug Company known as "Muscle & Joint Rub". Phase III clinical trial of Muscle & Joint Rub showed that it is effective and safe in the management of muscle sprains, contusions and inflammatory musculoskeletal disorders (Rajanna and Kolhapure, 2005). Since the studied product was combination of different plants, it is not clear what exact effect *Vanda tessellata* (Roxb.) Hook. ex G.Don. had on the studied disease. Additional human clinical studies are required to assess efficacy of this plant in treating skin diseases.

Verbascum thapsus L. has shown antiinflammatory and antioxidant properties in the earlier studies (Tables 3 and 4). Woolly leaves of this plant are generally irritant to skin of some susceptible individuals (White, 1887). Aqueous extract of leaves of this plant was tested for toxicity in Brine Shrimp and radish seed bioassays and extract showed toxicity at higher doses (around 1000 mg/L) with LC₅₀ of < 1000 mg/L (Turker and Camper, 2002). Verbascoside compound present in *Verbascum thapsus* L. has proven antiinflammatory action (Speranza et al., 2009) and 11,13 (18)-Oleanadiene-3,16,23,28-tetrol compound present in this plant has antiinflammatory properties (DNP, 2014). Clinical trials on 171 children aged 5–18 years have shown that *Verbascum thapsus* L. significantly improve ear pain associated with Otitis Media when treated with the infusion (Sarrell et al., 2003). *Verbascum thapsus* L. has also shown positive anti-tubercular properties in clinical trials (Carthy and Mahony, 2011). Clinical trials related to any skin disease have not been conducted on this plant.

Vitex negundo L. has shown wound healing, antiinflammatory, antioxidant, antimicrobial and tyrosinase inhibitory properties in the earlier studies (Tables 3 and 4). It has also shown trypsin

inhibitory properties (Gacche et al., 2008). Oil of this plant was non-toxic with LD₅₀ value of over 2000 mg/kg (Chattopadhyay et al., 2014) and had some important antifungal, antibacterial and antioxidant compounds (DNP, 2014) (Tables 3 and 4). The flavonoid-rich fraction (5,7,3'-trihydroxy, 6,8,4'-trimethoxy flavones) of *Vitex negundo* L. was found to antagonize the androgenic action of testosterone propionate on the male reproductive system of castrated prepubertal and intact adult dog (Bhargava, 1989). The principal compounds of *Vitex negundo* L. are flavone vitexcarpin (Diaz et al., 2003), nedundins A and B, diasyringaresinol, lyoniresinol, vitrofolal E and F (Azhar-UI-Haq et al., 2004), 6-phenyldihydronaphthalene-type lignan, vitedoin A, phenyl naphthalene-type ligand alkaloids, vitedoamine A, trinorlabdane-type diterpene and vitedoin (Ono et al., 2004). Mature leaves of *Vitex negundo* L. were found to have antihistaminic, membrane stabilizing, anti-inflammatory, analgesic (possibly mediated via prostaglandin synthesis inhibition) astringent and antioxidant activities (Dharmasiri et al., 2003). There are many polyherbal Ayurvedic skin care products with *Vitex negundo* L. in market. Extract of *Vitex negundo* L. is part of polyherbal formulation of The Himalaya Drug Company made of 4 herbs known as "Acne-N-Pimple Cream" recommended for the management of acne vulgaris. Phase III clinical trial of this cream on 26 patients suffering from acne vulgaris showed that application of cream reduced number of blackheads and whiteheads, in number of inflamed pustules and overall inflammation (Ravichandran et al., 2004a). *Vitex negundo* L. oil is one of the key ingredients of polyherbal "antiseptic cream" of The Himalaya Drug Company used for treatment of cuts, wounds, burns, rashes, sores and fungal skin infections. Open clinical trial of this cream was conducted on 50 patients in the age group of 20 to 50 years and study showed that this cream accelerated the rate of healing, reduced pus and odor from wounds, removed necrosis and inflammation, reduced wound size and prevented hypertrophic scarring (Indra Kumar, 2002). Oil of *Vitex negundo* L. is part of another polyherbal formulation of The Himalaya Drug Company made of 4 herbs known as "Diaper Rash Cream" recommended for the management of Infantile Irritant Diaper Dermatitis (IIDD). A prospective phase III clinical trial on 15 infants (birth weight of more than 2500 g) suffering from IIDD showed complete recovery from the clinical manifestations of IIDD, after a week's application (Chatterjee et al., 2005b). *Vitex negundo* L. contains Casticin, isoorientin, chrysophenol D, luteolin, p-hydroxybenzoic acid and D-fructose, which have anti-inflammatory and analgesic activities and it also contains bioflavonoids, which act through inhibition of prostaglandin biosynthesis resulting in both central and peripheral analgesic action (Telang et al., 1999). *Vitex negundo* L. along with 21 other plants is part of a polyherbal Ayurvedic formulation of The Himalaya Drug Company known as "Muscle & Joint Rub". Phase III clinical trial of Muscle & Joint Rub showed that it is effective and safe in the management of muscle sprains, contusions and inflammatory musculoskeletal disorders (Rajanna and Kolhapure, 2005). Water extract in combination with matra basti as 500 mg tablets showed relief from signs and symptoms of sciatica i.e., pain, weakness, numbness, and other discomforts in clinical studies on 119 patients in the age group of 20–60 years (Ali et al., 2010). A clinical trial of external (massage for 30 minutes) application of Nirgundi taila (*Vitex negundo* L. oil) was conducted on 40 children suffering from Saisaveeyavata (Poliomyelitis) and was found that it is effective in management of Poliomyelitis (Nair et al., 1988). Vitexins from this plant have cytotoxic effect on various types of cancer cell lines and also have antitumor activity on tumor xenograft models including breast, prostate, liver and cervical cancers (Zhou et al., 2009).

Wrightia arborea (Dennst.) Mabb. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Methanolic extract of leaves of this plant was found to be non-toxic even at high dose of 2000 mg/kg bw in acute oral toxicity study on albino rats (Nahar et al., 2013). A clinical study was conducted to evaluate the efficacy of coconut

oil extract of *Wrightia arborea* (Dennst.) Mabb. leaves on psoriasis patients with complaints of erythema, scaling, itching, dryness and roughness of skin as well as cracks, fissures and pain in palms and soles and results indicated a definite relief to the patients due to the indigenous drug by improving the clinical symptoms of disease (Kharmawphlang and Sarma, 2003). This is one of the potential plants for the development of skin care products in future. However, further clinical studies humans are required to assess its efficacy.

Ziziphus nummularia (Burm. f.) Wight & Arn. has shown wound healing, antiinflammatory, antioxidant and antimicrobial properties in the earlier studies (Tables 3 and 4). Cyclopeptide alkaloids isolated from the leaves of this plant are potent and efficacious analgesic agent and analgesic activity of these alkaloids is mediated by the peripheral as well as central pathways (Goyal et al., 2013). In acute oral toxicity study, LD₅₀ value of cyclopeptide alkaloids fraction was reported as 200 mg/kg bw and CNS depression and reduction in locomotor activity in mice treated with 200 and 300 mg/kg bw was observed (Goyal et al., 2013). However clinical trials related to any skin disease have not been conducted on this plant.

4. Conclusions

Medicinal plants and their traditional formulations have always been a part of social life of Tharu community, which have proved to be very useful in tackling various health related issues along with their key role in curing different skin diseases. The present study has revealed significant information on various medicinal plants used to treat skin diseases by Tharu community. Literature review has confirmed most of the claims made by the Tharu community regarding treatment of various skin diseases by the reported plants. Ayurveda is one of the most ancient medical traditions practiced in India, Sri Lanka and other South Asian countries and its literature describes over 200 herbs, minerals and fats to maintain and enhance the health and beauty of the skin (Datta et al., 2011). Some of the plants reported in the present study are also used in Ayurvedic system of medicine for treatment of various skin problems viz., *Calotropis procera* (Aiton) Dryand., *Azadirachta indica* A. Juss., *Brassica juncea* (L.) Czern., *Buchanania cochinchinensis* (Lour.) M.R.Almeida, *Butea monosperma* (Lam.) Taub., *Eclipta prostrata* (L.) L., *Euphorbia thymifolia* L., *Lawsonia inermis* L., *Mallotus philippensis* (Lam.) Mull.Arg., *Senna tora* (L.) Roxb., *Sesamum indicum* L. and *Shorea robusta* Gaertn. Information on use of these plants to treat skin problems is given in Dravyaguna Vijnana, section of Ayurveda dealing with drug sources, which is more than 1000 years old knowledge. Clinical trials conducted on polyherbal preparations made from these plants have shown their efficacy in treating various skin diseases, which has been discussed in detail in Section 3.2. Review of literature revealed that these are the most explored plants in the earlier pharmacological and clinical studies and some commercial skin care products based on these plants are already available in market.

Various tests on traditional medicinal plants show the effectiveness of traditional herbs against micro-organisms and turn into the bases of contemporary medicine (Evans et al., 2002). Literature review revealed that most of these plants have shown positive antimicrobial properties against wide spectrum of bacteria and fungi in earlier studies. Besides that most of the recorded plant species have shown antiinflammatory activities in the experimental animal models (Table 4). Further microbiological and antiinflammatory studies are recommended on the remaining plants to assess their antimicrobial and antiinflammatory properties. These plants can be evaluated in future studies for their skin care related activities. It is assumed that plants rich in ample

variety of secondary metabolites or phytochemicals, such as tannins, flavanoids, terpenoids, alkaloids, polyphenols are generally superior in their antimicrobial activities (Cowan, 1999; Maiyo et al., 2010). It can be observed in Table 4 that most of the recorded plants are rich in these secondary metabolites, which may have imparted them these positive antimicrobial properties. Information about the important compound isolated from the recorded plants, which have shown proven biological activity (i.e., Phytoalexin, antioxidant, antitumor, antibacterial, anti-inflammatory, antimicrobial, antifungal, antiviral and cytotoxic) have been given in Table 4, which will help in understanding reasons for efficacy of these plants in treatment of skin diseases. Generally it has been suggested that many herbal medicines have very effective wound healing properties because they encourage the repair mechanisms in the natural way (Shenoy et al., 2009a). In many studies, the formulations prepared using natural active ingredients showed therapeutic activity similar to the synthetic drugs and were effective for antibacterial and wound healing activity (Dhanekula et al., 2013).

Some of the plants reported in the present study have shown oral toxicity in earlier toxicological studies viz., *Annona squamosa* L., *Argemone mexicana* L., *Azadirachta indica* A. Juss., *Butea monosperma* (Lam.) Taub., *Calotropis procera* (Aiton) Dryand., *Cannabis sativa* L., *Datura stramonium* L., *Euphorbia hirta* L., *Ipomoea carnea* Jacq., *Lawsonia inermis* L., *Rauvolfia serpentina* (L.) Benth. ex Kurz and *Ricinus communis* L. Except *Azadirachta indica* A. Juss. all the other plants which have shown oral toxicity were used topically for treatment of skin diseases by the studied community. In case of *Azadirachta indica* A. Juss, stem bark is toxic at higher concentrations, but leaves are generally non-toxic which were used internally by the Tharu community for treatment of body infection.

Malanogenesis plays an important role in protecting the skin from sun-related injuries and is principally responsible for skin color, but at the same time abnormal hyperpigmentation such as freckles, chloasma, and lentigines can be serious esthetic problems (Han et al., 2012). Tyrosinases are widely used to regulate hyperpigmentation in cosmetic industry as it is one of the major targets in screening inhibitors of melanin synthesis (Nagarani et al., 2014). Flavonoid derived compounds are competitive inhibitors of tyrosinase substrates (Jeong et al., 2009). Besides that compounds having higher antioxidant activity and radical scavenging ability may be linked with tyrosinase inhibitory effects (Peng et al., 2013). Some of the plants reported in the present study have shown tyrosinase inhibitory properties (Table 3) viz., *Allium cepa* L., *Anisomeles indica* (L.) Kuntze, *Azadirachta indica* A. Juss., *Cannabis sativa* L., *Cassia fistula* L., *Chrysopogon zizanioides* (L.) Roberty, *Cleome viscosa* L., *Cuscuta reflexa* Roxb., *Ficus racemosa* L., *Ficus religiosa* L., *Lawsonia inermis* L., *Momordica dioica* Roxb. ex Willd., *Sesamum indicum* L. and *Vitex negundo* L. Trypsin inhibitors have important role in skin care. Some of the recorded plants have trypsin inhibitory properties viz., *Achyranthes aspera* L., *Allium cepa* L., *Albizia lebbeck* (L.) Benth., *Bauhinia variegata* L., *Brassica juncea* (L.) Czern., *Butea monosperma* (Lam.) Taub., *Cassia fistula* L., *Hyptis suaveolens* (L.) Poit., *Lawsonia inermis* L., *Linum usitatissimum* L., *Mirabilis jalapa* L., *Mucuna pruriens* (L.) DC., *Senna tora* (L.) Roxb., *Terminalia arjuna* (Roxb. ex DC.) Wight & Arn. and *Vitex negundo* L. Some of these plants can also be targeted for development of new skin care products.

The literature review and discussion on the reported plants in Section 3.2 has revealed that products from very few of the reported plants are available in market, while most of the reported plants are still under preclinical or clinical trials. There are various known phytochemicals and antibiotic, antibacterial, antiviral and antifungal agents present in these plants which may be synthesized or transformed to make pharmaceuticals. Some of the reported plants have shown promising results in preclinical trails

and there is a need of clinical trials to see their safety and efficacy in treating various skin diseases. These plants may be targeted for development of new medicines, ointments or drugs for the treatment of skin diseases. However further toxicological, preclinical and clinical studies are needed to validate claims about little worked out plant species reported in the present study viz., *Sida cordata* (Burm. F.) Borss. Waalk., *Millettia extensa* (Benth.) Baker, *Caesulia axillaris* Roxb., *Ehretia laevis* Roxb., *Vanda tessellate* (Roxb.) Hook. Ex G.Don. and *Eualaliopsis binata* (Retz.) C.E. Hubb. Further studies on these plants are recommended to assess their potential in development of new skin care products.

Acknowledgment

Authors are thankful to the traditional healers and local folk of Tharu community for providing valuable information and sharing their knowledge with us. One of the authors Dr. Jyotsana Sharma is thankful to University Grants Commission, India for providing financial support under Dr. D.S. Kothari Postdoctoral Fellowship (13-811/2013(BSR)).

References

- Abbasi, A.M., Khan, M.A., Ahmad, M., Zafar, M., Jahan, S., Sultana, S., 2010a. Ethnopharmacological application of medicinal plants to cure skin diseases and in folk cosmetics among the tribal communities of North-West Frontier Province, Pakistan. *Journal of Ethnopharmacology* 128, 322–335.
- Abbasi, M.A., Zafar, A., Riaz, T., Aziz-ur-Rehman, Arshad, S., Shahwar, D., Jahangir, M., Siddiqui, S.Z., Shahzadi, T., Ajaib, M., 2010b. Evaluation of comparative antioxidant potential of aqueous and organic fractions of *Ipomoea carnea*. *Journal of Medicinal Plants Research* 4, 1883–1887.
- Abdelouahab, B., Nadia, M., Nabila, K., 2011. Acute toxicity study of *Datura stramonium* seeds in rat. *Research Opinions in Animal & Veterinary Sciences* 1, 434–440.
- Abdulazeez, A.M., Awasum, C.A., Dogo, Y.S., Abiayi, P.N., 2010. Effect of *Peristrophe bicalyculata* on blood pressure, kidney and liver functions of two kidney one clip (2K1C) hypertensive rats. *British Journal of Pharmacology and Toxicology* 1, 101–107.
- Abdulla, M.A., Hassandarvish, P., Ali, H.M., Noor, S.M., Mahmoud, D.H., Bashah, N.S.A., Rokik, R.H., Khalil, W.I., 2009. Acceleration of wound healing potential of *Lantana camara* leaf extract in experimental rats. *Research Journal of Medical Sciences* 3, 75–79.
- Abdulsalaam, I.A., Ehinmidu, J.O., Igbadi, E.O., 2013. Evaluation of antibacterial properties, acute toxicity and immuno-stimulatory potential of *Scoparia dulcis*. *Nigerian Journal of Biotechnology* 26, 21–25.
- Abubakar, A., Ogbadanyi, E.O., Okogun, J.I., Gbodi, T.I., Tifin, U.F., 2012. Acute and sub chronic toxicity of *Tridax procumbens* in experimental animals. *Journal of Environmental Science Toxicology and Food Technology* 1, 19–27.
- Abubakar, E.M.M., 2009. Antibacterial activity of crude extracts of *Euphorbia hirta* against some bacterial associated with enteric infections. *Journal of Medicinal Plants Research* 3, 498–505.
- Adhikari, A., Devkota, H.P., Takano, A., Masuda, K., Nakane, T., Basnet, P., Skalko-Basnet, N., 2008. Screening of Nepalese crude drugs traditionally used to treat hyperpigmentation: *in vitro* tyrosinase inhibition. *International Journal of Cosmetic Science* 30, 353–360.
- Adhikari, B.S., Babu, M.M., Saklani, P.L., Rawat, G.S., 2010. Medicinal plants diversity and their conservation status in Wildlife Institute of India (WII) Campus, Dehradun. *Ethnobotanical Leaflets* 14, 46–83.
- Adsull, V.B., Khatiwora, E., Torane, R., Deshpande, N.R., 2012. Antimicrobial activities of *Ipomoea carnea* leaves. *Journal of Natural Products and Plant Resources* 2, 597–600.
- Afzal, M., Wang, Z., Ali, F., Song, Z., Cox, R., Khan, S., 2013. Preliminary phytochemical screening and antimicrobial activities of various fractions of *Mallotus philippensis* Muell. *Journal of Medicinal Plants Research* 7, 3066–3070.
- Agarwal, U., Pande, P.S., Patki, P.S., Mitra, S.K., 2009. Evaluation of the clinical efficacy and safety of anti-dandruff hair cream in the treatment of dandruff. *The Antiseptic* 106, 37–39.
- Aggarwal, B.B., Prasad, S., Reuter, S., Kannappan, R., Yadev, V.R., Park, B., Kim, J.H., Gupta, S.C., Phromnoi, K., Sundaram, C., Prasad, S., Chaturvedi, M.M., Sung, B., 2011. Identification of novel anti-inflammatory agents from ayurvedic medicine for prevention of chronic diseases: “Reverse Pharmacology” and “Bedside to Bench” approach. *Current Drug Targets* 12, 1595–1653.
- Agnihotri, A., Singh, V., 2013. Effect of *Tamarindus indica* Linn. and *Cassia fistula* Linn. stem bark extracts on oxidative stress and diabetic conditions. *Acta Poloniae Pharmaceutica* 70, 1011–1019.

- Agrahari, A.K., Panda, S.K., Meher, A., Padhan, A.R., 2010a. Screening of wound healing activity of *Curculigo orchoides* Gaertn root tubers' methanolic extract. *International Journal of Pharmaceutical and Applied Sciences* 1, 91–95.
- Agrahari, A.K., Panda, S.K., Meher, A., Padhan, A.R., 2010b. Studies on the anti-inflammatory properties of *Curculigo orchoides* Gaertn. Root tubers. *International Journal of Pharmaceutical Sciences and Research* 1, 139–143.
- Aguilar, A., Camacho, J.R., Chino, S., Jacquez, P., Lopez, M.E., 1994. *Herbario Medicinal del Instituto Mexicano del Seguro Social*. Instituto Mexicano del Seguro Social, Mexico, D.F.
- Aguirre, C., Valdes-Rodriguez, S., Mendoza-Hernandez, G., Rojo-Dominguez, A., Blanco-Labra, A., 2004. A novel 8.7 kDa pectinase inhibitor from chan seeds (*Hyptis suaveolens* L.) inhibits proteases from the larger grain borer *Prostephanus truncatus* (Coleoptera: Bostrichidae). *Comparative Biochemistry and Physiology – Part B: Biochemistry and Molecular Biology* 138, 81–89.
- Aher, P.S., Shinde, Y.S., Chavan, P.P., 2012. *In vitro* evaluation of antibacterial potential of *Annona squamosa* against pathogenic bacteria. *International Journal of Pharmaceutical Sciences and Research* 3, 1457–1460.
- Ahmed, S.A., Ross, S.A., Slade, D., Radwan, M.M., Zulfikar, F., Elshohly, M.A., 2008. Cannabinoid ester constituents from high-potency *Cannabis sativa*. *Journal of Natural Products* 71, 536–538.
- Akintobi, O.A., Agunbiade, S.O., Okonko, I.O., Ojo, O.V., 2011. Antimicrobial evaluation and phytochemical analysis of leaf extracts of *Mirabilis jalapa* against some human pathogenic bacteria. *Nature and Science* 9, 45–53.
- Alam, B., Hossain, S., Habib, R., Rea, J., Islam, A., 2012. Antioxidant and analgesic activities of *Lannea coromandelica* Linn. Bark extract. *International Journal of Pharmacology* 8, 224–233.
- Al-Bayati, F.A., 2007. Antibacterial activity of *Linum usitatissimum* L. seeds and active compound detection. *Rafidain Journal of Science* 18, 27–36.
- Ali, K.M., Chatterjee, K., De, D., Maiti, S., Pathak, T.K., Ghosh, D., 2011. Evaluation of antioxidant activity of seed of *Holarrhena antidysenterica*: an approach through different *in vitro* models. *Journal of Natural Pharmaceuticals* 2, 115–118.
- Ali, M., Shukla, V.D., Dave, A.R., Bhatt, N.N., 2010. A clinical study of Nirugundi Ghana Vati and Matra Basti in the management of Gridhrasi with special reference to sciatica. *AYU* 31, 456–460.
- Alia, B.H., Bashir, A.K., Tanira, M.O., 1995. Anti-inflammatory, antipyretic, and analgesic effects of *Lawsonia inermis* L. (henna) in rats. *Pharmacology* 51, 356–363.
- Allayurveda.com, 2014. Evaluation of NEEM GUARD (A Herbal Preparation) as a Versatile Healer. (http://allayurveda.com/research_neemguard.asp) (accessed 21.09.14).
- Allen, P.H., 1943. Poisonous and injurious plants of Panama. *American Journal of Tropical Medicine* 23, 3–76.
- Allison, A.C., Cacabelos, R., Lombardi, V.R., Alvarez, X.A., Vigo, C., 2001. Celastrol, a potent antioxidant and anti-inflammatory drug, as a possible treatment for Alzheimer's disease. *Progress in Neuro-psychopharmacology and Biological Psychiatry* 25, 1341–1357.
- Ambali, S., Silas, A., Ayo, J.O., Ibrahim, N.D.G., Obalowu, A.A., Salami, S.O., 2005. Toxicity studies on the methanolic extract of the leaves of *Cassia tora* Linn in mice. *Nigerian Veterinary Journal* 27, 54–61.
- Ambiga, S., Narayanan, R., Gowri, D., Sukumar, D., Madhavan, S., 2007. Evaluation of wound healing activity of flavanoids from *Ipomoea carnea* Jacq. *Ancient Science of Life* 26, 45–51.
- Amresh, G., Rao, C.V., Singh, P.N., 2007a. Antioxidant activity of *Cissampelos pareira* on benzo(a)pyrene-induced mucosal injury in mice. *Nutrition Research* 27, 625–632.
- Amresh, G., Reddy, G.D., Rao, C.V., Singh, P.N., 2007b. Evaluation of anti-inflammatory activity of *Cissampelos pareira* root in rats. *Journal of Ethnopharmacology* 110, 526–531.
- Amresh, G., Singh, P.N., Rao, C.V., 2008. Toxicological screening of traditional medicine Laghupatha (*Cissampelos pareira*) in experimental animals. *Journal of Ethnopharmacology* 116, 454–460.
- Anandakirouchenane, E., Chandiran, I.S., Kanimozhi, V., Kadalmani, B., 2013. Antioxidant and protective effect of *Curculigo orchoides* on liver, pancreas and kidney tissue in alloxan induced diabetic experimental rats. *Drug Invention Today* 5, 192–200.
- Anela, S., De, S., Kanthal, L.K., Choudhury, N.S.K., Das, B.L., Sagar, K.V., 2011. Acute oral toxicity studies of *Pongamia pinnata* and *Annona squamosa* on albino wistar rats. *International Journal of Research in Pharmacy and Chemistry* 1, 820–824.
- Anusuya, N., Gomathi, R., Manian, S., Sivaram, V., Menon, A., 2012. Evaluation of *Basella rubra* L., *Rumex nepalensis* Spreng. and *Commelina benghalensis* L. for antioxidant activity. *International Journal of Pharmacy and Pharmaceutical Sciences* 4, 714–720.
- Anwar, F., Przybylski, R., 2012. Effect of solvents extraction on total phenolics and antioxidant activity of extracts from flaxseed (*Linum usitatissimum* L.). *Acta Scientiarum Polonorum. Technologia Alimentaria* 11, 293–301.
- Appendino, G., Gibbons, S., Giana, A., Pagani, A., Grassi, G., Stavri, M., Smith, E., Rahman, M.M., 2008. Antibacterial Cannabinoids from *Cannabis sativa*: a structure–activity study. *Journal of Natural Products* 71, 1427–1430.
- Arfan, M., Amin, H., Karamac, M., Kosinska, A., Shahidi, F., Wiczowski, W., Amarowicz, R., 2007. Antioxidant activity of extracts of *Mallotus philippinensis* fruit and bark. *Journal of Food Lipids* 14, 280–297.
- Arfan, M., Amin, H., Khan, N., Khan, I., Saeed, M., Khan, M.A., Fazal-ur-Rehman, 2010. Analgesic and anti-inflammatory activities of 11-O-galloylbergenin. *Journal of Ethnopharmacology* 131, 502–504.
- Arora, N., Pandey-Rai, S., 2014. GC–MS analysis of the essential oil of *Celastrus paniculatus* Willd. seeds and antioxidant, anti-inflammatory study of its various solvent extracts. *Industrial Crops and Products* 61, 345–351.
- Arote, S.R., Dahikar, S.B., Yeole, P.G., 2009. Phytochemical screening and antibacterial properties of leaves of *Pongamia pinnata* Linn. (Fabaceae) from India. *African Journal of Biotechnology* 8, 6393–6396.
- Arthur, R.P., Shelly, W.B., 1955. The role of proteolytic enzymes in the production of pruritus in man. *Journal of Investigative Dermatology* 25, 341–346.
- Arunachalam, G., Subramanian, N., Pazhani, G.P., Ravichandran, V., 2009. Anti-inflammatory activity of methanolic extract of *Eclipta prostrata* L. (Asteraceae). *African Journal of Pharmacy and Pharmacology* 3, 97–100.
- Arung, E.T., Kusuma, I.W., Shimizu, K., Komndo, R., 2011. Tyrosinase inhibitory effect of quercetin 4-O-β-D-glucopyranoside from dried skin of red onion (*Allium cepa*). *Natural Product Research* 25, 256–263.
- Arya, K.R., Agarwal, S.C., 2008. Folk therapy for eczema, bone fracture, boils, sores and gingivitis in Taragtal province of Uttaranchal. *Indian Journal of Traditional Knowledge* 7, 443–445.
- Arya, S., Kumar, V.L., 2005. Antiinflammatory efficacy of extracts of latex of *Calotropis procera* against different mediators of inflammation. *Mediators of Inflammation* 2005, 228–232.
- Asare, G.A., Addo, P., Bugyei, K., Gyan, B., Adjei, S., Out-Nyarko, L.S., Wiredu, E.K., Nyarko, A., 2011. Acute toxicity studies of aqueous leaf extract of *Phyllanthus niruri*. *Interdisciplinary Toxicology* 4, 206–210.
- Ashafa, A.O.T., Orekoya, L.O., Yakubu, M.T., 2012. Toxicity profile of ethanolic extract of *Azadirachta indica* stem bark in male Wistar rats. *Asian Pacific Journal of Tropical Biomedicine* 2, 811–817.
- Ashok, P., Rajani, G.P., Arulmozhi, S., Hulkoti, B., Desai, B.G., Rajendra, R., 2006. Anti-inflammatory and anti-ulcerogenic effect of *Crotalaria juncea* Linn. in albino rats. *Iranian Journal of Pharmacology and Therapeutics* 5, 141–144.
- Asif, M., Kumar, A., 2010. Acute toxicity study and in-vivo anti-inflammatory activity of different fractions of *Curculigo orchoides* Gaertn. rhizome in albino Wistar rats. *Iranian Journal of Pharmaceutical Sciences* 6, 191–198.
- Aswar, P.B., Khadabadi, S.S., Kuchekar, B.S., Rajurkar, R.M., Saboo, S.S., Javarkar, R.D., 2009. *In-vitro* evaluation of anti-bacterial and anti-fungal activity of *Vitex negundo* (Verbenaceae). *Ethnobotanical Leaflets* 13, 962–967.
- Atal, C.K., Dubey, R.K., Singh, J., 1985. Biochemical basis of enhanced drug bioavailability by piperine: evidence that piperine is a potent inhibitor of drug metabolism. *Journal of Pharmacology and Experimental Therapeutics* 232, 258–262.
- Atigari, D.V., Gundamaraju, R., Sabbithi, S., Chaitanya, B.K., Ramesh, C., 2012. Evaluation of antiepileptic activity of methanolic extract of *Celastrus paniculatus* Willd whole plant in rodents. *International Journal of Pharmaceutical and Phytopharmacological Research* 2, 20–25.
- Atique, A., Beigh, S.Y., Iqbal, M., 2009. Evaluation of a toxic plant *Ipomoea carnea* for curing leucorrhoea. *Journal of Tropical Medicinal Plants* 10, 185–188.
- Augustine, B.B., Dash, S., Lahkar, M., Lihite, R.J., Samudrala, P.K., Pitta, S., 2013. Effect of *Mirabilis jalapa* Linn. flowers in experimentally induced arthritis and consecutive oxidative stress. *International Journal of Pharmacy & Pharmaceutical Sciences* 5, 190–193.
- Awasthi, S., Irshad, M., Das, M.K., Ganti, S.S., Moshahid, A.R., 2009. Anti-Inflammatory activity of *Calotropis gigantea* and *Tridax procumbens* on carrageenin-induced paw edema in rats. *Ethnobotanical Leaflets* 13, 568–577.
- Ayanwuyi, L.O., Yaro, A.H., Adamu, H.Y.S., 2009. Studies on anticonvulsant activity of methanol capitulum extract of *Leonotis nepetifolia* Linn. *Nigerian Journal of Pharmaceutical Sciences* 8, 73–79.
- Ayyanar, M., Ignacimuthu, S., 2009. Herbal medicines for wound healing among tribal people in Southern India: ethnobotanical and scientific evidences. *International Journal of Applied Research in Natural Products* 2, 29–42.
- Azhar-ul-Haq, Malik, A., Anis, I., Khan, S.B., Ahmed, E., Ahmed, Z., 2004. Enzymes inhibiting lignans from *Vitex negundo*. *Chemical and Pharmaceutical Bulletin* 52, 1269–1272.
- Azhar-ul-Haq, Malik, A., Khan, M.T.H., Anwar-ul-Haq, Khan, S.B., Ahmad, A., Choudhary, M.I., 2006. Tyrosinase inhibitory lignans from the methanol extract of the roots of *Vitex negundo* Linn. and their structure–activity relationship. *Phytomedicine* 13, 255–260.
- Aziz-ur-rehman, Ahmad, S., Riaz, T., Abbas, A., Abbasi, M.A., Siddiqui, S.Z., Ajaib, M., 2012. *Martynia annua*: comparative antioxidant potential of its stem and leaves. *Asian Journal of Chemistry* 24, 3335.
- Azmi, M.B., Qureshi, S.A., 2012. Methanolic root extract of *Rauwolfia serpentina* improves the glucose tolerance in wistar mice. *Journal of Food and Drug Analysis* 20, 484–488.
- Babu, A.R.S., Karki, S.S., 2011. Antiinflammatory activity of various extracts of roots of *Calotropis procera* against different inflammation models. *International Journal of Pharmacy & Pharmaceutical Sciences* 3, 191–194.
- Babu, B.H., Mohana, L., Saravana Kumar, S.A., 2010. Studies on phytochemical and anticonvulsant property of *Martynia annua* Linn. *International Journal of Phytopharmacology* 1, 82–86.
- Badole, S.L., Zanwar, A.A., Khopade, A.N., Bodhankar, S.L., 2011. *In vitro* antioxidant and antimicrobial activity cycloart-23-ene-2β,25-diol (B2) isolated from *Pongamia pinnata* (L. Pierre). *Asian Pacific Journal of Tropical Medicine* 4, 910–916.
- Bafna, A., Mishra, S., 2010. Antioxidant and immunomodulatory activity of the alkaloidal fraction of *Cissampelos pareira* Linn. *Scientia Pharmaceutica* 78, 21–31.
- Bagewadi, Z.K., Siddanagouda, R.S., Baligar, P.G., 2012. Phytochemical screening and evaluation of antimicrobial activity of *Semecarpus anacardium* nuts. *International Journal of Pharmacology and Pharmaceutical Technology* 1, 68–74.

- Bairagi, S.M., Aher, A.A., Nema, N., Nimase, P.K., 2012. Anti-inflammatory evaluation of methanol extract and aqueous fraction of the bark of *Bauhinia variegata* (Leguminosae). *International Journal of Research in Pharmacy and Chemistry* 2, 77–82.
- Bala, V., Debnath, A., Shill, A.K., Bose, U., 2011. Anti-inflammatory, diuretic and antibacterial activities of aerial parts of *Mucuna pruriens* Linn. *International Journal of Pharmacology* 7, 498–503.
- Baral, M., Chakraborty, S., Chakraborty, P., 2010. Evaluation of anthelmintic and anti-inflammatory activity of *Amaranthus spinosus* Linn. *International Journal of Current Pharmaceutical Research* 2, 44–47.
- Barnes, M.H., 1931. Linseed dermatitis. *Journal of Industrial Hygiene* 13, 49.
- Basma, A.A., Zakaria, Z., Latha, L.Y., Sasidharan, S., 2011. Antioxidant activity and phytochemical screening of the methanol extracts of *Euphorbia hirta* L. *Asian Pacific Journal of Tropical Medicine*, 386–390.
- Begum, D., Nath, S.C., 2000. Ethnobotanical review of medicinal plants used for skin diseases and related problems in Northeastern India. *Journal of Herbs Spices and Medicinal Plants* 7, 55–93.
- Behl, P.N., Captain, R.M., Bedi, B.M.S., Gupta, S., 1966. Skin-Irritant and Sensitizing Plants Found in India. P.N. Behl, Irwin Hospital, New Delhi.
- BenShabat, S., Eftekhari, F., Habibi, Z., Massarat, S., Malekzadeh, R., 2009. Antibacterial activity of twenty Iranian plant extracts against clinical isolates of *Helicobacter pylori*. *Iranian Journal of Basic Medical Sciences* 12, 105–111.
- Bhalodia, N.R., Shukla, V.J., 2011. Antibacterial and antifungal activities from leaf extracts of *Cassia fistula* L. An ethnomedicinal plant. *Journal of Advanced Pharmaceutical Technology and Research* 2, 104–109.
- Bhandary, M.J., Chandrashekar, K.R., 2002. Glimpses of ethnic herbal medicine of coastal Karnataka. *Ethnobotany* 14, 1–12.
- Bhanot, A., Shri, R., 2010. A comparative profile of methanol extracts of *Allium cepa* and *Allium sativum* in diabetic neuropathy in mice. *Pharmacognosy Research* 2, 374–384.
- Bharadwaj, S., Nayak, S., Joshi, V.K., 2013. An open comparative randomized clinical study of plant based drugs *Embelia robusta* Roxb and *Mallotus philippinensis* Muell on intestinal parasites. *Journal of Gastrointestinal & Digestive System* 3, 1–5.
- Bharathi, V., Varalakshmi, B., Gomathi, S., Shanmugapriya, A., Karpaga, T., 2012. Antibacterial activity of *Tridax procumbens* Linn. *International Journal of Pharma Sciences and Research* 3, 364–367.
- Bhardwaj, M., Surekha, Duhan, J.S., 2011. Free radical scavenging and antimutagenic potential of acetone, chloroform and methanol extracts of leaf of *Argemone mexicana*. *International Journal of Pharmacy and Bio Sciences* 2, 455–464.
- Bhargava, S.K., 1989. Antiandrogenic effects of a flavonoid-rich fraction of *Vitex negundo* seeds: a histological and biochemical study in dogs. *Journal of Ethnopharmacology* 27, 327–339.
- Bhat, P., Hegde, G., Hegde, G.R., 2012. Ethnomedicinal practices in different communities of Uttara Kannada district of Karnataka for treatment of wounds. *Journal of Ethnopharmacology* 143, 501–514.
- Bhat, P., Hegde, G.R., Hegde, G., Mulgund, G.S., 2014. Ethnomedicinal plants to cure skin diseases – an account of the traditional knowledge in the coastal parts of Central Western Ghats, Karnataka. *Indian Journal of Ethnopharmacology* 151, 493–502.
- Bhatia, G., Puri, A., Maurya, R., Yadav, P.P., Khan, M.M., Khanna, A.K., Saxena, J.K., 2008. Anti-dyslipidemic and antioxidant activities of different fractions of *Pongamia pinnata* (lin.) fruits. *Medicinal Chemistry Research* 17, 281–289.
- Bhatnagar, P.K., Katiyar, C.K., Khanna, N., Upadhyay, D.J., Swaminathan, S., Srinivas, K., Sharma, N., Kanaujia, A., Sood, R., Singhal, S., Shukla, G., Duggar, R., Pareek, P. K., Singh, Y., Khan, S., Raut, R., 2011. Anti Dengue Activity of *Cissampelos pareira* Extracts US20120107424 (published on 30 November).
- Bhatt, R.M., Kora, S., 1984. Clinical and experiment study of Panchavalkal and Shatavari on burn wound sepsis-bacterial and fungal. *Journal of the National Integrated Medical Association* 26, 131–133.
- Bhattacharyya, D., Sur, T.K., Lyle, N., Jana, U., Debnath, P.K., 2011. A clinical study on the management of generalized anxiety disorder with *Vaca* (*Acorus calamus*). *Indian Journal of Traditional Knowledge* 10, 668–671.
- Bigoniya, R., Agrawal, S., Verma, N.K., 2013. Potential wound healing activity of *Euphorbia hirta* Linn. total flavonoid fraction. *International Journal of Pharmaceutical Sciences Review and Research* 22, 149–156.
- Bisht, R., Bhattacharya, S., 2013. Effect of various extracts of *Desmodium gangeticum* on Streptozotocin-nicotinamide induced type-2 diabetes. *Asian Journal of Plant Science & Research* 3, 28–34.
- Biswas, M., Karan, T.K., Bhattacharya, S., Kumar, R.B.S., Ghosh, A.K., Haldar, P.K., 2011a. Acute and sub-chronic toxicity of *Terminalia arjuna* leaf in swiss albino mice. *Pharmacologyonline* 1, 366–371.
- Biswas, M., Biswas, K., Karan, T.K., Bhattacharya, S., Ghosh, A.K., Haldar, P.K., 2011b. Evaluation of analgesic and anti-inflammatory activities of *Terminalia arjuna* leaf. *Journal of Phytology* 3, 33–38.
- Bobby, M.N., Wesely, E.G., Johnson, M.A., 2012. *In vitro* anti-bacterial activity of leaves extracts of *Albizia lebbek* Benth against some selected pathogens. *Asian Pacific Journal of Tropical Biomedicine* 2, S859–S862.
- Bommareddy, A., Zhang, X., Schrader, D., Kaushik, R.S., Zeman, D., Matthees, D.P., Chandradhar, D., 2009. Effects of dietary flaxseed on intestinal tumorigenesis in Apc^{Min} Mouse. *Nutrition and Cancer* 61, 276–283.
- Bouzidi, A., Mahdeb, N., Kara, N., 2011. Toxicity studies of alkaloids of seeds of *Datura stramonium*. *The Israel Medical Association Journal* 3, 538–539.
- Cai, Y., Luo, Q., Sun, M., Corke, H., 2004. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sciences* 74, 2157–2184.
- Campanati, A., Savelli, A., Sandroni, L., Marconi, B., Giuliano, A., Giuliodori, K., Ganzetti, G., Offidani, A., 2010. Effect of allium cepa-allantoin-pentaglycan gel on skin hypertrophic scars: clinical and video-capillaroscopic results of an open-label, controlled, nonrandomized clinical trial. *Dermatologic Surgery* 36, 1439–1444.
- Carthy, M.E., Mahony, J.M.O., 2011. What's in a name? Can Mullein weed beat TB? Where modern drugs are failing? (Article ID 239237). *Evidence-Based Complementary and Alternative Medicine*, 7.
- Chai, T., Mohan, M., Ong, H., Wong, F., 2014. Antioxidant, iron-chelating and anti-glucosidase activities of *Typha domingensis* Pers (Typhaceae). *Tropical Journal of Pharmaceutical Research* 13, 67–72.
- Chakraborty, A., Brantner, A., Mukainaka, T., Nobukuni, Y., Kuchide, M., Konoshima, T., Tokuda, H., Nishino, H., 2002. Cancer chemopreventive activity of *Achyranthes aspera* leaves on Epstein–Barr virus activation and two-stage mouse skin carcinogenesis. *Cancer Letters* 177, 1–5.
- Chakraborty, M.K., Bhattacharjee, A., 2006. Some ethnomedicinal uses for various diseases in Purulia district, West Bengal. *Indian Journal of Traditional Knowledge* 5, 554–558.
- Chamarro, G., Carduno, L., Martinez, E., Madrigal, E., Tamariz, J., Salazar, M., 1998. Dominant lethal study of α -asarone in male mice. *Toxicology Letters* 99, 71–72.
- Chandramohan, J.B., Reddy, D.N., 2006. Burn wound healing activity of *Euphorbia hirta*. *Ancient Science of Life* 25, 16–18.
- Chanprapaph, K., Tanrattanakorn, S., Wattanakrai, P., Wongkitisophon, P., Vachiramon, V., 2012. Effectiveness of onion extract gel on surgical scars in Asians (Article ID 212945). *Dermatology Research and Practice*, 6.
- Charde, R.M., Dhongade, H.J., Charde, M.S., Kasture, A.V., 2010. Evaluation of antioxidant, wound healing and anti-inflammatory activity of ethanolic extract of leaves of *Ficus religiosa*. *International Journal of Pharmaceutical Sciences and Research* 1, 73–82.
- Chatterjee, S., Pramanick, N., Chattopadhyay, S., Munian, K., Kolhapure, S.A., 2005a. Evaluation of the efficacy and safety of “Baby Powder” in infantile hyderhidrosis, miliaria rubra and bad body odor. *The Antiseptic* 102, 126–128.
- Chatterjee, S., Pramanick, N., Chattopadhyay, S., Munian, K., Kolhapure, S.A., 2005b. Evaluation of the efficacy and safety of “Diaper Rash Cream” in the management of infantile irritant diaper dermatitis (IIDD). *The Antiseptic* 102, 251–255.
- Chatterjee, S., Pramanick, N., Chattopadhyay, S., Munian, K., Kolhapure, S.A., 2005c. Evaluation of the efficacy and safety of “Nourishing Baby Oil” in infantile xerosis. *The Antiseptic* 102, 179–182.
- Chattopadhyay, D., Mukherjee, H., Bag, P., Ojha, D., Kumar, K.A., Dutta, S., Haldar, P. K., Chatterjee, T., Sharon, A., Chakraborti, S., 2012. Inhibition of NO₂, PGE₂, TNF- α , and iNOS Expression by *Shorea robusta* L.: an ethnomedicine used for anti-inflammatory and analgesic activity (Article ID 254849). *Evidence-Based Complementary and Alternative Medicine*, 14.
- Chattopadhyay, P., Banerjee, S., Pathak, M.P., Agnihotri, A., Karmakar, S., Goyary, D., Dhiman, S., Veer, V., 2014. Acute and subchronic dermal toxicity of *Vitex negundo* essential oil. *Cutaneous and Ocular Toxicology* 33, 6–21.
- Chaudhari, M., Menqi, S., 2006. Evaluation of phytoconstituents of *Terminalia arjuna* for wound healing activity in rats. *Phytotherapy Research* 20, 799–805.
- Chavan, M.J., Wakte, P.S., Shinde, D.B., 2010. Analgesic and anti-inflammatory activity of caryophyllene oxide from *Annona squamosa* L. bark. *Phytomedicine* 17, 149–151.
- Chawla, A.S., Sharma, A.K., Handa, S.S., 1992. Chemical investigation and anti-inflammatory activity of *Vitex negundo* seeds. *Journal of Natural Products* 55, 163–167.
- Chinnasamy, N., Harishankar, N., Kumar, P.U., Rukmini, C., 1993. Toxicological studies on debitterized neem oil (*Azadirachta indica*). *Food and Chemical Toxicology* 31, 297–301.
- Chinese Pharmacopoeia, 2010. The Pharmacopoeia of the People's Republic of China. English Edition, 3 Volumes. Chinese Pharmacopoeia Commission, China.
- Chitra, V., Dharani, P., Pavan, K.K., Narayana, R.A., 2009. Wound healing activity of alcoholic extract of *Buchanania lanzan* in albino rats. *International Journal of ChemTech Research* 1, 1026–1031.
- Cho, J.W., Cho, S.Y., Lee, S.R., Lee, K.S., 2010. Onion extract and quercetin induce matrix metalloproteinase-1 *in vitro* and *in vivo*. *International Journal of Molecular Medicine* 25, 347–352.
- Chopda, M.Z., Mahajan, R.T., 2009. Wound healing plants of Jalgaon District of Maharashtra State, India. *Ethnobotanical Leaflets* 13, 1–32.
- Chopra, R.N., Bhadwar, R.L., 1940. Poisonous plants of India. *Indian Journal of Agricultural Sciences* 10, 1–44.
- Choudhary, S.M., Mishra, N., Upadhyay, T.S., Upadhyay, R., 2011. Indigenous Knowledge of using Medicinal Plants in Treating Skin diseases by Tribal's in Central Narmada Valley of Madhya Pradesh (India). *Bulletin of Environment, Pharmacology and Life Sciences* 1, 60–63.
- Choudhury, P.K., Jadhav, S., 2013. Acute toxicity studies of methanolic bark extract of *Ficus racemosa* and root extract of *Cissampelos pareira*. *International Journal of Pharma and Bio Sciences* 4, 1050–1055.
- Chouhan, H.S., Sahu, A.N., Singh, S.K., 2011. Fatty acid composition, antioxidant, anti-inflammatory and antibacterial activities of seed oil from *Crotalaria juncea* Linn. *Journal of Medicinal Plants Research* 5, 984–991.
- Chouhan, H.S., Singh, S.K., 2010. Antibacterial activity of seed and flower parts of *Crotalaria juncea* Linn. *American–Eurasian Journal of Scientific Research* 5, 212–215.
- Chulet, R., Pradhan, P., Sharma, K.S., Jhajharia, K.M., 2010. Phytochemical screening and anti microbial activity of *Albizia lebbek*. *Journal of Chemical and Pharmaceutical Research* 2, 476–484.

- Cierpicki, T., Otlewski, J., 2000. Determination of a high precision structure of a novel protein, *Linum usitatissimum* trypsin inhibitor (LUTI), using computer-aided assignment of NOESY cross-peaks. *Journal of Molecular Biology* 302, 1179–1192.
- Coca, A.F., Walzer, M., Thommen, A.A., 1931. Asthma and Hay Fever in Theory and Practice. Charles C. Thomas, Springfield, IL, pp. 260–261.
- Cowan, M.M., 1999. Plant products as antimicrobial agents. *Clinical Microbiology Reviews* 12, 564–582.
- Crooke, W., 1896. The Tribes and Castes of the North Western Province and Oudh, Calcutta (Rep. Ed., titled “The Tribes and Castes of the North Western India”, vols. 1–4, 1975, Delhi).
- Cuellar, C., Armando, O., Dennis, O., 2010. Preliminary phytochemical and antimicrobial evaluation of the fresh and dried whole plant extracts from *Commelina benghalensis*. *Revista Colombiana de Ciencia Animal* 2, 104–116.
- Cunnane, S.C., Hamadeh, M.J., Liede, A.C., Thompson, L.U., Wolever, T.M., Jenkins, D. J., 1995. Nutritional attributes of traditional flaxseed in healthy young adults. *The American Journal of Nutrition* 61, 62–68.
- Dahab, M.M., Musa, E.M., Osman, E.A., Koko, W.S., Almgoul, A.Z., Khogali, M.S., 2013. Antigiardial activity and toxicological exploration of *Cannabis sativa* extracts. *Journal of Forest Products and Industries* 2, 24–29.
- Dahare, K.D., Jain, A., 2010. Ethnobotanical studies on plant resources of Tahsil Multai, district Betul, Madhya Pradesh, India. *Ethnobotanical Leaflets* 14, 694–705.
- Dange, S.V., Grandhi, S., 2009. Clinical efficacy of Foot Care Cream in the management of foot cracks. *Indian Medical Journal* 103, 350–353.
- Dangwal, L.R., Sharma, A., Rana, C.S., 2010. Ethno-medicinal plants of the Garhwal Himalaya used to cure various diseases: a case study. *New York Science Journal* 3, 28–31.
- Das, S., Haldarm, P.K., Pramanik, G., Suresh, R.B., 2010. Evaluation of Anti-Inflammatory activity of *Clerodendron infortunatum* Linn. extract in rats. *Global Journal of Pharmacology* 4, 48–50.
- Das, P., Mekap, S., Pani, S., Sethi, R., Nayak, P., 2010. Pharmacological evaluation of anti-inflammatory activity of *Euphorbia hirta* against carrageenan induced paw edema in Rats. *Der Pharmacia Lettre* 2, 151–154.
- Das, S., Bhattacharya, S., Biswas, M.A., Kar, B., Kumar, R.B.S., Pramanik, G., Haldar, P. K., 2011. Acute and sub-chronic toxicity study of *Clerodendron infortunatum* leaf in adult male albino mice. *American-Eurasian Journal of Scientific Research* 6, 188–191.
- Dash, G.K., Murthy, P.N., 2011a. Wound healing effects of *Ageratum conyzoides* Linn. *International Journal of Pharma and Bio Sciences* 2, 369–382.
- Dash, G.K., Murthy, P.N., 2011b. Evaluation of *Argemone mexicana* Linn. leaves for wound healing activity. *Journal of Natural Products and Plant Resources* 1, 46–56.
- Dash, S.S., Misra, M.K., 1999. Taxonomic survey and systematic census of economic plants of Narayana Patna Hills of Koraput Dist. Orissa. *Journal of Economic and Taxonomic Botany* 23, 473–498.
- Datta, H.S., Mitra, S.K., Patwardhan, B., 2011. Wound healing activity of topical application forms based on Ayurveda. Evidence Based Complementary and Alternative Medicine 2011, 134378.
- de Souza, A.F., Torquato, R.J., Tanaka, A.S., Sampaio, C.A., 2005. Cloning, expression and characterization of *Bauhinia variegata* trypsin inhibitor BvTI. *Biological Chemistry* 386, 1185–1189.
- Deshimaru, M., Watanabe, A., Suematsu, K., Hatano, M., Terada, S., 2003. Purification, amino acid sequence, and cDNA cloning of trypsin inhibitors from onion (*Allium cepa* L.) bulbs. *Bioscience, Biotechnology, and Biochemistry* 67, 1653–1659.
- Deshmukh, S.R., Dhanashree, A.S., Patil, B.A., 2012. Extraction and evaluation of indole alkaloids from *Rauwolfia serpentina* for their antimicrobial and anti-proliferative activities. *International Journal of Pharmacy and Pharmaceutical Sciences* 4, 329–334.
- Devi, P., Meera, R., 2010. Study of antioxidant, anti-inflammatory and wound healing activity of extracts of *Litsea glutinosa*. *Journal of Pharmaceutical Sciences and Research* 2, 155–163.
- Devi, S.A., Ganjewala, D., 2011. Antioxidant activities of methanolic extracts of sweet-flag (*Acorus calamus*) leaves and rhizomes. *Journal of Herbs, Spices and Medicinal Plants* 17, 1–11.
- Devi, S.L., Divakar, M.C., 2013. Wound healing activity studies of *Wrightia arborea* Phytosome in rats. *Hygeia – Journal for Drugs and Medicines* 4, 87–94.
- Devi, V., Shanbhag, T.V., Bairy, K.L., Rao, N., Shenoy, S., 2005. Effect of *Phyllanthus niruri* on wound healing in rats. *Indian Journal of Physiology and Pharmacology* 49, 487–490.
- Devprakash, Srinivasan, K.K., Subburaju, T., Singh, S.K., 2011. Antimicrobial activity of alcoholic and aqueous extracts of *Vetiveria zizanioides*. *Journal of Pharmacy Research* 4, 1343–1344.
- Dewan, S.M.R., Amin, M.N., Adnan, T., Uddin, S.M.N., Shahid-Ud-Daula, A.F.M., Sarwar, G., Hossain, M.S., 2013. Investigation of analgesic potential and *in vitro* antioxidant activity of two plants of Asteraceae family growing in Bangladesh. *Journal of Pharmacy Research* 6, 599–603.
- Dhalwal, K., Deshpande, Y.S., Purohit, A.P., 2007. Evaluation of *in vitro* antioxidant activity of *Sida rhombifolia* (L.) ssp. *retusa* (L.). *Journal of Medicinal Food* 10, 683–688.
- Dhandapani, R., Lakshmi, D., Balakrishnan, V., Jayakumar, A., 2007. Preliminary phytochemical investigation and antibacterial activity of *Phyllanthus amarus* Schum & Thorn. *Ancient Science of Life* 28, 1–5.
- Dhanekula, S., Srinivas, P., Mamidi, S., 2013. Formulation and evaluation of herbal gels for antimicrobial and wound healing activity. *Journal of Natural Pharmaceuticals* 4, 141–148.
- Dharmasiri, M.G., Jayakody, J.R., Galhena, G., Liyanage, S.S., Ratnasooriya, W.D., 2003. Anti-inflammatory and analgesic activities of mature fresh leaves of *Vitex negundo*. *Journal of Ethnopharmacology* 87, 199–206.
- Dharmasiri, M.G., Ratnasooriya, W.D., Thabrew, M.I., 2002. Anti-inflammatory activity of decoctions of leaves and stem of *Anisomeles indica* at preflowering and flowering stages. *Pharmaceutical Biology* 40, 433–439.
- Diallo, A., Ekl-Gadegheku, K., Agbonon, A., Aklilikou, K., Creepy, E.E., Gbeassor, M., 2010. Acute and sub-chronic (28-day) oral toxicity studies of Hydroalcohol leaf extract of *Ageratum conyzoides* L. (Asteraceae). *Tropical Journal of Pharmaceutical Research* 9, 463–467.
- Diaz, F., Chavez, D., Lee, D., Mi, Q., Chai, H.B., Tan, G.T., Kardono, L.B., Riswan, S., Fairchild, C.R., Wild, R., Farnsworth, N.R., Cordell, G.A., Pezzuto, J.M., Kinghorn, A.D., 2003. Cytotoxic flavone analogues of vitexicarpin, a constituent of the leaves of *Vitex negundo*. *Journal of Natural Products* 66, 865–867.
- Dixit, P., 2013. A comparative screening of antibacterial activity of *Anisomeles indica* with *Mentha piperita* against human pathogenic micro-organisms. *Indian Journal of Fundamental and Applied Life Sciences* 3, 85–88.
- DNP, 2014. Dictionary of Natural Products 22.2. Taylor & Francis Group (accessed on 10.03.14).
- Donga, S., Shukla, V.J., Ravishanker, B., Ashok, B.K., Mishtry, I.U., 2011. Chronic toxicity study of *Butea monosperma* (Linn.) Kuntze seeds in albino rats. *AYU* 32, 120–125.
- Draeos, Z.D., Baumann, L., Fleischer, A.B., Plaum, S., Avakian, E.V., Hardas, B., 2012. A new proprietary onion extract gel improves the appearance of new scars. *The Journal of Clinical and Aesthetic Dermatology* 5, 18–24.
- Dubey, G.P., Agarwal, A., Singh, A., 1993. Evaluation of D-400, an indigenous herbal preparation in diabetes mellitus. *Indian Journal of Internal Medicine* 3, 183–186.
- Duddukuri, G.R., Rao, D.E., Kaladhar, D.S.V.G.K., Sastry, Y.N., Rao, K.K., Chaitanya, K. K., Sireesha, Ch., 2011. Preliminary studies on *in vitro* antibacterial activity and phytochemical analysis of aqueous crude extract of *Shorea robusta* floral parts. *International Journal of Current Research* 3, 21–23.
- Duke, J.A., 2012. Ethnobotanical Uses. Germplasm Resources Information Network: Dr. Duke's Phytochemical and Ethnobotanical Databases. USDA.
- Edwin, S., Jarald, E.E., Deb, L., Jain, A., King, H., Dutt, K.R., Raj, A.A., 2008. Wound healing and antioxidant activity of *Achyranthes aspera*. *Pharmaceutical Biology* 46, 824–828.
- Eftekhari, F., Yousefzadi, M., Tafakor, V., 2005. Antimicrobial activity of *Datura innoxia* and *Datura stramonium*. *Fitoterapia* 76, 118–120.
- Engels, C., Schieber, A., Ganzle, M.G., 2012. Sinapic acid derivatives in defatted oriental mustard (*Brassica juncea* L.) seed meal extracts using UHPLC–DAD–ESI–MSⁿ and identification of compounds with antibacterial activity. *European Food Research and Technology* 234, 535–542.
- Etkin, N. (Ed.), 1994. Eating on the Wild Side. The Pharmacologic, Ecologic and Social Implications of Using Noncultigens. University of Arizona Press, Tucson.
- Evans, C.E., Banno, A., Samuel, O.A., 2002. Efficacy of some nupur medicinal plants against *Salmonella typhi*: an *in vitro* study. *Journal of Ethnopharmacology* 80, 21–24.
- Farahpour, M.R., Taghikhani, H., Habibi, M., Zandieh, M.A., 2011. Wound healing activity of flaxseed *Linum usitatissimum* L. in rats. *African Journal of Pharmacy and Pharmacology* 5, 2386–2389.
- Ferreira, A.A., Bezerra, G.A., Matos, F.J., Viana, G.S., 2000. Antinociceptive, anti-inflammatory and bronchodilator activities of Brazilian medicinal plants containing coumarin: a comparative study. *Journal of Ethnopharmacology* 70, 151–159.
- Ferreira, F.S., Brito, S.V., Ribeiro, S.C., Almeida, W.O., Alves, R.R.N., 2009. Zootherapeutics utilized by residents of the community Poco Dantas, Crato-CE, Brazil. *Journal of Ethnobiology and Ethnomedicine* 5, 21.
- Figueiredo, A.S., Modesto-Filho, J., 2008. Effect of defatted sesame (*Sesamum indicum* L.) flour on the blood glucose level in type 2 diabetic women. *Revista Brasileira de Farmacognosia* 18, 77–83.
- Fikru, A., Makonnen, E., Eguale, T., Debell, A., Mekonnen, G.A., 2012. Evaluation of *in vivo* wound healing activity of methanol extract of *Achyranthes aspera* L. *Journal of Ethnopharmacology* 143, 469–474.
- Friedman, J., Yaniv, Z., Dafni, A., Palewitch, D., 1986. A preliminary classification of the healing potential of medicinal plants, based on a rational analysis of an ethnopharmacological field survey among Bedouins in the Negev desert. *Israel Journal of Ethnopharmacology* 16, 275–287.
- FSI, 2011. India State of Forest Report 2011, Chapter 9.28 – Uttarakhand Forest Survey of India (FSI). Ministry of Environment & Forests, Government of India, Dehradun, Uttarakhand, India, p. 236.
- Gacche, R.N., Dhole, N.A., Jadhav, A.D., 2008. Antioxidant and anti-inflammatory related activities of certain botanicals frequently used in Ayurveda and other indigenous systems of medication. *Australian Journal of Medical Herbalism* 20, 1–27.
- Gaire, B.P., Subedi, L., 2013. A review on the pharmacological and toxicological aspects of *Datura stramonium* L. *Journal of Integrative Medicine* 11, 73–79.
- Gairola, S., Sharma, J., Bedi, Y.S., 2014. A cross-cultural analysis of Jammu, Kashmir and Ladakh (India) medicinal plant use. *Journal of Ethnopharmacology* 155, 925–986.
- Gairola, S., Sharma, J., Gaur, R.D., Siddiqi, T.O., Painuli, R.M., 2013. Plants used for treatment of dysentery and diarrhoea by the Bhoja community of district Dehradun, Uttarakhand, India. *Journal of Ethnopharmacology* 150, 989–1006.

- Ganesan, S., Kesaven, L., 2003. Ethnomedicinal plants used by the ethnic group of Valaiyans of Vellimalai Hills (Reserved forest) Tamil Nadu, India. *Journal of Economic and Taxonomic Botany* 27, 754–760.
- Gangwar, K.K., Deepali, Gangwar, R.S., 2010. Ethnomedicinal plant diversity in Kumaun Himalaya of Uttarakhand, India. *Nature and Science* 8, 66–78.
- Garg, V.K., Paliwal, S.K., 2011. Wound-healing activity of ethanolic and aqueous extracts of *Ficus benghalensis*. *Journal of Advanced Pharmaceutical Technology & Research* 2, 110–114.
- Garipelli, N., Chinnalalaiah, R., Potnuri, N., Pigili, R.K., 2012. Anti-inflammatory and anti-oxidant activities of ethanolic extract of *Euphorbia thymifolia* Linn whole plant. *International Journal of Pharmacy & Pharmaceutical Sciences* 4, 516–519.
- Gaur, R.D., 1999. Flora of the District Garhwal: North West Himalaya (with Ethnobotanical Notes). Transmedia, Srinagar (Garhwal).
- Gaur, R.D., Sharma, J., 2011. Indigenous knowledge on the utilization of medicinal plants diversity in Siwalik region of Garhwal Himalaya, Uttarakhand. *Journal of Forest Science* 27, 23–31.
- Gaur, R.D., Sharma, J., Painuli, R.M., 2010. Plants used in traditional healthcare of livestock by Gujjar community of Sub-Himalayan tracts, Uttarakhand, India. *Indian Journal of Natural Products and Resources* 1, 243–248.
- Gaur, R.D., Sharma, J., Painuli, R.M., 2011. Folk herbal medicines used by the Gujjar tribe of Sub-Himalayan tracts, Uttarakhand. *Journal of Economic and Taxonomic Botany* 35, 224–230.
- Gautam, S., Jain, A.K., Kumar, A., 2011. Potential antimicrobial activity of *Zizyphus nummularia* against medically important pathogenic microorganisms. *Asian Journal of Traditional Medicine* 6, 267–271.
- Gazzaneo, L.R.S., Lucena, R.F.P., Albuquerque, U.P., 2005. Knowledge and use of medicinal plants by local specialists in an region of Atlantic Forest in the state of Pernambuco (Northeastern Brazil). *Journal of Ethnobiology and Ethnomedicine* 1, 9. <http://dx.doi.org/10.1186/1746-4269-1-9>.
- Geetha, K., Davuluri, S.P., 2013. Purification of Trypsin inhibitor from *Achyranthes aspera* seeds. *International Journal of Advancement in Research & Technology* 2, 215–222.
- Ghimirey, A.K., Jin, C., Ghimire, B.K., Cho, D.H., 2009. Antioxidant activity and quantitative estimation of azadirachtin and nimbin in *Azadirachta indica* A. Juss grown in foothills of Nepal. *African Journal of Biotechnology* 8, 3084–3091.
- Ghimire, B.K., Seong, E.S., Kim, E.H., Ghimirey, A.K., Yu, C.Y., Ghimire, B.K., Chung, I. M., 2011. A comparative evaluation of the antioxidant activity of some medicinal plants popularly used in Nepal. *Journal of Medicinal Plants Research* 5, 1884–1891.
- Gholamhoseinian, A., Moradi, M.N., Sharifi-far, F., 2009. Screening the methanol extracts of some Iranian plants for acetylcholinesterase inhibitory activity. *Research in Pharmaceutical Sciences* 4, 105–112.
- Ghorband, D.P., Biradar, S.D., 2011. Traditional medicines knowledge in Dharmabad taluka of Nanded district, Maharashtra, India. *Indian Journal of Natural Products and Resources* 2, 498–503.
- Gidwani, B.K., Bhargava, S., Rao, S.P., Majoomdar, A., Pawar, D.P., Alaspure, R.N., 2009. Analgesic, anti-inflammatory and anti-hemorrhoidal activity of aqueous extract of *Lantana camara* Linn. *Research Journal of Pharmacy and Technology* 2, 378–381.
- Girach, R.D., Singh, S., Brahmam, M., Misra, M.K., 1999. Traditional treatment of skin diseases in Bhadrak District, Orissa. *Journal of Economic and Taxonomic Botany* 27, 754–760.
- Giri, S.N., Raja, K., Fata, J., Sun, C., Balambika, R., Bolton, D., 2014. *Clerodendrum viscosum* and Methods of Use. US Patent No. US 20140134731 A1, Original Assignee The City University of New York.
- Gnanaswari, K., Raju, V.R.R., 2012. Preliminary phytochemical screening and antimicrobial evaluation of *Leonotis nepetifolia* (L.) R.Br. *Journal of Natural Products and Plant Resources* 2, 689–692.
- Gokhale, A.B., Damre, A.S., Kulkarni, K.R., Saraf, M.N., 2002. Preliminary evaluation of anti-inflammatory and anti-arthritis activity of *S. lappa*, *A. speciosa* and *A. aspera*. *Phytomedicine* 9, 433–437.
- Golla, U., Bhimathathi, S.S.R., 2014. Evaluation of antioxidant and DNA damage protection activity of the hydroalcoholic extract of *Desmostachya bipinnata* L. Stapf (Article ID 215084). *The Scientific World Journal*, 8.
- Gommers, F.J., Voor in't Holt, D.J.M., 1976. Chemotaxonomy of compositae related to their host suitability for *Pratylenchus penetrans*. *Netherlands Journal of Plant Pathology* 82, 1–8.
- Gouthamchandra, K., Mahmood, R., Manjunatha, H., 2010. Free radical scavenging, antioxidant enzymes and wound healing activities of leaves extracts from *Clerodendrum infortunatum* L. *Environmental Toxicology and Pharmacology* 30, 11–18.
- Govindarajan, R., Rastogi, S., Vijayakumar, M., Shirwaikar, A., Rawat, A.K.S., Mehrotra, S., Pushpangadan, P., 2003. Studies on the antioxidant activities of *Desmodium gangeticum*. *Biological and Pharmaceutical Bulletin* 26, 1424–1427.
- Gowanloch, J.N., Brown, C.A., 1943. Poisonous Snakes, Plants and Black Widow Spider of Louisiana. Louisiana Department of Conservation, New Orleans, LA, USA.
- Goyal, M., Ghosh, M., Nagori, B.P., Sasmal, D., 2013. Analgesic and anti-inflammatory studies of cyclopeptide alkaloid fraction of leaves of *Zizyphus nummularia*. *Saudi Journal of Biological Sciences* 20, 365–371.
- Grassi, P., Urias Reyes, T.S., Sosa, S., Tubaro, A., Hofer, O., Zitterl-Eglseer, K., 2006. Anti-inflammatory activity of two diterpenes of *Hyptis suaveolens* from El Salvador. *Zeitschrift für Naturforschung C61*, 165–170.
- Greenberg, L.A., Lester, D., 1954. Handbook of Cosmetic Materials – The Properties, Uses, and Toxic and Dermatologic Actions. Interscience Publishers, Inc., New York p. 455.
- Grice, E.A., Kong, H.H., Conlan, S., Deming, C.B., Davis, J., Young, A.C., NISC Comparative Sequencing Program, Bouffard, G.G., Blakesley, R.W., Murray, P. R., Green, E.D., Turner, M.L., Segre, J.A., 2009. Topographical and temporal diversity of the human skin microbiome. *Science* 324, 1190–1192.
- Guha, G., Rajkumar, V., Mathew, L., Kumar, R.A., 2011. The antioxidant and DNA protection potential of Indian tribal medicinal plants. *Turkish Journal of Biology* 35, 233–242.
- Gunde, M.C., Meshram, S.S., Dangre, P.V., Amnerkar, N.D., 2013. Wound healing properties of methanolic extract of seeds of *Mucuna pruriens*. *International Journal of Pharmacognosy and Phytochemical Research* 5, 57–59.
- Gupta, C., Katewa, S.S., 2012. Antimicrobial testing of *Vanda tessellate* leaf extracts. *Journal of Medicinal and Aromatic Plant Science* 34, 158–162.
- Gupta, P.C., Sharma, N., Rao, C.V., 2011. Comparison of the antioxidant activity and total phenolic, flavonoid content of aerial part of *Cleome viscosa* L. *International Journal of Phytomedicine* 3, 386–391.
- Gupta, S., Raghuvanshi, M., Jain, D., 2010. Comparative studies on anti-inflammatory activity of *Coriandrum sativum*, *Datura stramonium* and *Azadirachta indica*. *Asian Journal of Experimental Biological Sciences* 1, 151–154.
- Gupta, V., Kumar, P., Bansal, P., Singh, R., 2009. Anti-inflammatory and antinociceptive activity of *Mitragyna parvifolia*. *Asian Journal of Medical Sciences* 1, 97–99.
- Hahm, T.S., Park, S.J., Martin, L., 2009. Effects of germination on chemical composition and functional properties of sesame (*Sesamum indicum* L.) seeds. *Bior-resource Technology* 100, 1643–1647.
- Hajare, S.W., Chandra, S., Sharma, J., Tandan, S.K., Lal, J., Telang, A.G., 2001. Anti-inflammatory activity of *Dalbergia sissoo* leaves. *Fitoterapia* 72, 131–139.
- Han, Y.K., Park, Y.J., Ha, Y.M., Park, D., Lee, J.Y., Lee, N., Yoon, J.H., Moon, H.R., Chung, H.Y., 2012. Characterization of a novel tyrosinase inhibitor, (2R,5R)-2-(2,4-dihydroxyphenyl)thiazolidine-4-carboxylic acid (MHY384). *Biochimica et Biophysica Acta* 1820, 542–549.
- Harish, B.G., Krishna, V., Kumar, H.S.S., Ahamed, B.M.K., Sharath, R., Swamy, H.M.K., 2008. Wound healing activity and docking of glycogen-synthase-kinase-3-beta-protein with isolated triterpenoid lupeol in rats. *Phytomedicine* 15, 763–767.
- Hasan, M.D., Ahmed, M.D., Sukla, M., Uddin, S.J., Masud, M.M., Sadhu, S.K., Ishibashi, M., 2006. Antioxidant, antinociceptive activity and general toxicity study of *Dendrophthoe falcata* and isolation of quercitrin as the major component. *Oriental Pharmacy and Experimental Medicine* 6, 355–360.
- Hasan, S.M.R., Hossain, M.M., Akter, R., Jamila, M., Mazumder, M.E.H., Rahman, S., 2009. DPPH free radical scavenging activity of some Bangladeshi medicinal plants. *Journal of Medicinal Plants Research* 3, 875–879.
- Hausen, B.M., 1973. Holzarten mit Gesundheitsschadigenden Inhaltstoffen. DRW-Verlag, Stuttgart.
- Health Canada, 2014. Summary of Health Canada's Assessment of a Health Claim About Ground Whole Flaxseed and Blood Cholesterol Lowering. Bureau of Nutritional Sciences, Food Directorate, Health Products and Food Branch, Health Canada. (<http://www.hc-sc.gc.ca/fn-an/label-etiquet/claims-reclam/assess-evalu/flaxseed-graines-de-lin-eng.php>) (accessed 24.09.14).
- Hegde, M.M., Lakshman, K., Girija, K., Kumar, B.S.A., Lakshmiprasanna, V., 2010. Assessment of antiarthralgic activity of *Desmostachya bipinnata* L. (Poaceae) root extracts. *Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas* 9, 312–318.
- Heinrich, M., Ankil, A., Frei, B., Weimann, C., Sticher, O., 1998. Medicinal plants in Mexico, Healer's consensus and cultural importance. *Social Science and Medicine* 47, 1859–1871.
- Heinrich, M., Rimpler, H., Antonio Barrera, N., 1992. Indigenous phytotherapy of gastrointestinal disorders in a Mixe Lowland community. *Journal of Ethnopharmacology* 36, 63–80.
- Hema, T.A., Arya, A.S., Suseelan, S., Celestial, R.K.J., Divya, P.V., 2013. Antimicrobial activity of five south Indian medicinal plants against clinical pathogens. *International Journal of Pharma and Bio Sciences* 4, 70–80.
- Hiremath, V.T., Vijaykumar, M.M.J., Taranath, T.C., 2010. Survey on ethnomedicinal plants of Jogimatti Forest Chitradurga District, Karnataka, India. *Environment and We An International Journal of Science and Technology* 5, 223–233.
- Ho, W.S., Ying, S.Y., Chan, P.C., Chan, H.H., 2006. Use of onion extract, heparin, allantoin gel in preventing of scarring in Chinese patients having laser removal of tattoos: a prospective randomized controlled trial. *Dermatologic Surgery* 32, 891–896.
- Hosamani, P.A., Lakshman, H.C., Sandeep, K.K., Hosamani, R.C., 2011. Antimicrobial activity of leaf extract of *Andrographis paniculata* Wall. *Science Research Reporter* 1, 92–95.
- Hosamath, P.V., 2011. Evaluation of antimicrobial activity of *Litsea glutinosa*. *International Journal of Pharmaceutical Applications* 2, 105–114.
- Hou, Y.Y., Wu, M.L., Hwang, Y.C., Chang, F.R., Wu, Y.C., Wu, C.C., 2009. The natural diterpenoid ovatodiolide induces cell cycle arrest and apoptosis in human oral squamous cell carcinoma Ca9-22 cells. *Life Sciences* 85, 26–32.
- Hsieh, S.C., Fang, S.H., Rao, Y.K., Tzeng, Y.M., 2008. Inhibition of pro-inflammatory mediators and tumor cell proliferation by *Anisomeles indica* extracts. *Journal of Ethnopharmacology* 118, 65–70.
- Hsu, H.F., Chang, S.F., Chen, Z.H., Huang, K.H., Fang, L.W., Chiou, S.J., Yen, J.H., Tsai, Y. D., Wang, C.P., Yuan, S.S.F., Hwang, J.Y., 2012a. Cytotoxic effect of *Anisomeles indica* extract on human pharynx squamous cancer cells. *Journal of Medicinal Plants Research* 6, 5002–5012.
- Hsu, H., Chen, Z., Chang, S., Wang, C., Chiou, S., Yen, J., Chang, C., Tsai, Y., Fang, L., Hwang, J., 2012b. Evaluating the anti-metastatic potential of *Anisomeles indica* extract by using human oral squamous carcinoma FaDu cells. *African Journal of Pharmacy and Pharmacology* 6, 1782–1791.

- Huang, H., Lien, H., Ke, H., Chang, L., Chen, C., Chang, T., 2012. Antioxidative characteristics of *Anisomeles indica* extract and inhibitory effect of ovotodioid on melanogenesis. *International Journal of Molecular Sciences* 13, 6220–6235.
- Huang, J., Li, H., Yang, J., Chen, Y., Liu, Y., Li, N., Nie, C., 2004. Chemical components of *Vetiveria zizanioides* volatiles. *Ying Yong Sheng Tai Xue Bao* 15, 170–172.
- Hussain, M., Farooq, U., Rashid, M., Bakhsh, H., Majeed, A., Khan, I.A., Rana, S.L., Shafeeq-ur-Rahman, Aziz, A., 2014. Antimicrobial activity of fresh latex, juice and extract of *Euphorbia hirta* and *Euphorbia thymifolia* – an *in vitro* comparative study. *International Journal of Pharma Sciences* 4, 546–553.
- Ibrahim, H.A., Ibrahim, H., 2009. Phytochemical screening and toxicity evaluation on the leaves of *Argemone mexicana* Linn. (Papaveraceae). *International Journal of Applied Sciences* 3, 39–43.
- Ilango, K., Maharajan, G., Narasimhan, S., 2003. Analgesic and anti-inflammatory activities of *Momordica dioica* fruit pulp. *Natural Product Sciences* 9, 210–212.
- Ilango, K., Maharajan, G., Narasimhan, S., 2012. Preliminary phytochemical screening and antibacterial activity of fruit pulp of *Momordica dioica* Roxb. (Cucurbitaceae). *African Journal of Basic and Applied Sciences* 4, 12–15.
- Ilavarasan, R., Mallika, M., Subramanian, V., 2006. Anti-inflammatory and free radical scavenging activity of *Ricinus communis* root extract. *Journal of Ethnopharmacology* 103, 478–480.
- Ilavarasan, R., Mallika, M., Venkataraman, S., 2011. Toxicological assessment of *Ricinus communis* Linn. root extracts. *Toxicology Mechanisms and Methods* 21, 246–250.
- Ilavarasan, R., Mallika, M., Venkataraman, S., 2005. Anti-inflammatory and antioxidant activities of *Cassia fistula* Linn. bark extracts. *African Journal of Traditional, Complementary and Alternative Medicines* 2, 70–85.
- Imamdas, F.B., Oswal, R.J., Chorage, T.V., Garje, K., 2011. *In Vitro* antimicrobial activity of *Cuscuta reflexa* Roxb. *International Research Journal of Pharmacy* 2, 214–216.
- Indhumathi, T., Mohandass, S., Shibi, A., 2014. Acute toxicity study of ethanolic extract of *Solanum incanum* L. fruit. *Asian Journal of Pharmaceutical and Clinical Research* 7, 98–100.
- Indra Kumar, S.V., 2002. Clinical trial of a herbal antiseptic cream in the treatment of minor injuries. *Bombay Hospital Journal* 44, 52–55.
- Jachak, S.M., Gautam, R., Selvam, C., Madhan, H., Srivastava, A., Khan, T., 2011. Anti-inflammatory, cyclooxygenase inhibitory and antioxidant activities of standardized extracts of *Tridax procumbens* L. *Fitoterapia* 82, 173–177.
- Jain, A., Katewa, S.S., Galav, P.K., Sharma, P., 2005. Medicinal plant diversity of Sitamata wildlife sanctuary, Rajasthan, India. *Journal of Ethnopharmacology* 102, 143–157.
- Jain, A., Soni, M., Deb, L., Jain, A., Rout, S.P., Gupta, V.B., Krishna, K.L., 2008. Antioxidant and hepatoprotective activity of ethanolic and aqueous extracts of *Momordica dioica* Roxb. leaves. *Journal of Ethnopharmacology* 115, 61–66.
- Jain, N., Jain, R., Jain, A., Jain, D.K., Chandel, H.S., 2010. Evaluation of wound-healing activity of *Acorus calamus* Linn. *Natural Product Research* 24, 534–541.
- Jain, R., Rawat, S., Jain, S.C., 2013. Phytochemicals and antioxidant evaluation of *Ficus racemosa* root bark. *Journal of Pharmacy Research* 6, 615–619.
- Jayasutha, J., Nithila, S.M.J., 2011. Evaluation of wound healing activity of ethanolic extract of *Aristolochia bracteata* and *Cassia tora* on wistar albino rats. *International Journal of PharmTech Research* 3, 1547–1550.
- Jaythirtha, M.G., Mishra, S.H., 2004. Preliminary immunomodulatory activities of methanol extracts of *Eclipta alba* and *Centella asiatica*. *Phytomedicine* 11, 361–365.
- JECFA (Joint FAO/WHO Expert Committee on Food Additives), 1981. Monograph on β -asarone. WHO Food Additive Series No. 16.
- Jeong, S.H., Ryu, Y.B., Curtis-Long, M.J., Ryu, H.W., Baek, Y.S., Kang, J.E., Lee, W.S., Park, K.H., 2009. Tyrosinase inhibitory polyphenols from roots of *Morus ihou*. *Journal of Agricultural and Food Chemistry* 57, 1195–1203.
- John, J., Mehta, A., Mehta, P., Mehta, P., 2011. Antimicrobial and phytochemical analysis of *Cassia tora* leaf extracts. *Journal of Basic and Applied Biology* 5, 125–131.
- Johnsy, G., Davidson, S.S., Kaviyaran, V., 2012. Indigenous knowledge of medicinal plants used for the treatment of skin diseases by the Kaani tribe of Kanyakumari district. *International Journal of Pharmacy and Pharmaceutical Sciences* 4, 309–313.
- Joshi, B., Tyagi, V., 2011. Traditional knowledge and utilization of medicinal plants of Himalayan Region. *Nature and Science* 9, 1–6.
- Joshi, P., Bhanu, P., Gahlot, M., 2014. Anti-inflammatory and analgesic activities of methanol extract of *Vallisneria spiralis* leaves. *International Journal of Pharmacy and Pharmaceutical Sciences* 6, 622–624.
- Jothy, S.L., Zakaria, Z., Chen, Y., Lau, Y.L., Latha, L.Y., Sasidharan, S., 2011. Acute oral toxicity of methanolic seed extract of *Cassia fistula* in mice. *Molecules* 16, 5268–5282.
- Joy, B., Mathew, M., 2010. Anti-oxidant studies and chemical investigation of ethanol extract of *Acalypha indica* Linn. In: Awaad, A.S., Govil, J.N., Singh, V.K. (Eds.), *Recent Progress in Medicinal Plants*, vol. 27, pp. 261–279.
- Jyothi, B., Sudarsanam, G., Sitaram, B., Prasada, B.G., Yasodamma, N., 2010. Ethnobotanical survey of medicinal plants used in the treatment of dermatogenic diseases in Chittoor District, Andhra Pradesh, India. *Ethnobotanical Leaflets* 14, 511–517.
- Kaithwas, G., Mukherjee, A., Chaurasia, A.K., Majumdar, D.K., 2011. Anti-inflammatory, analgesic and antipyretic activities of *Linum usitatissimum* L. (flaxseed/linseed) fixed oil. *Indian Journal of Experimental Biology* 49, 932–938.
- Kala, C.P., 2009. Aboriginal uses and management of ethnobotanical species in deciduous forests of Chhattisgarh state in India. *Journal of Ethnobiology and Ethnomedicine* 5, 20. <http://dx.doi.org/10.1186/1746-4269-5-20>.
- Kalita, S., Kumar, G., Karthik, L., Rao, K.V.B., 2012. *In vitro* antioxidant and DNA damage inhibition activity of aqueous extract of *Lantana camara* L. (Verbenaceae) leaves. *Asian Pacific Journal of Tropical Biomedicine* 2, S1675–S1679.
- Kalpna, Upadhyay, A., 2010. Anti-inflammatory evaluation of ethanolic extract of leaves of *Holoptelea integrifolia*, Planch. *Annals of Biological Research* 1, 185–195.
- Kamble, R.D., Rode, K.N., Maske, K.S., Shaikh, H.M., 2013. Anti-inflammatory and analgesic activity of various fractions of *Vetiveria zizanioides* (V.Z.) in rodents. *Asian Journal of Plant Science & Research* 3, 61–69.
- Kamble, S.Y., Patil, S.R., Sawant, P.S., Sawant, S., Pawar, S.G., Singh, E.A., 2010. Studies on plants used in traditional medicine by Bhilla tribe of Maharashtra. *Indian Journal of Traditional Knowledge* 9, 591–598.
- Kant, S., Agnihotri, M.S., Dixit, K.S., 2001. Clinical evaluation of *Boerhaavia diffusa* as an adjuvant in the treatment of pulmonary tuberculosis. *Phytomedica* 2, 89–94.
- Karabow, A.S., 1927. An occupational affection of the skin in workers preparing castor oil. *Moskovskii Meditsinskii Zhurnal* 7, 47.
- Karmakar, U.K., Ghosh, D., Sadhu, S.K., 2011. Assessment of analgesic, cytotoxic and antioxidant activities of *Vallisneria spiralis* (Roth) Kuntze. *Stamford Journal of Pharmaceutical Sciences* 4, 64–68.
- Karthikeyan, K., Gandhi, S.S., Srinivasan, R., Chidambaram, C., Subramanian, K., 2012. *In vitro* antibacterial activity of *Desmodium gangeticum* (L.) DC. *Asian Pacific Journal of Tropical Disease*, S421–S424.
- Kassuya, C.A., Silvestre, A., Menezes-de-Lima, O.J., Marotta, D.M., Rehder, V.L., Calixto, J.B., 2006. Antiinflammatory and antiallodynic actions of the lignan niranthin isolated from *Phyllanthus amarus*. Evidence for interaction with platelet activating factor receptor. *European Journal of Pharmacology* 546, 182–188.
- Katewa, S.S., Chaudhary, B.L., Jain, A., 2004. Folk herbal medicines from tribal area of Rajasthan, India. *Journal of Ethnopharmacology* 92, 41–46.
- Katewa, S.S., Galav, P.K., 2005. Traditional herbal medicines from Shekhawati region of Rajasthan. *Indian Journal of Traditional Knowledge* 4, 237–245.
- Katiyar, N.S., Rao, N.V., Gangwar, A.K., 2012. Evaluation of anti-inflammatory activity of stem extracts of *Cuscuta reflexa* (Roxb) in rats. *International Journal of Research in Pharmaceutical and Biomedical Sciences* 3, 1805–1808.
- Katzenschlanger, R., Evans, A., Manson, A., Patsalos, P.N., Ratnaraj, N., Watt, H., Timmermann, L., Van der Giessen, R., Lees, A.J., 2004. *Mucuna pruriens* in Parkinson's disease: a double blind clinical and pharmacological study. *Journal of Neurology, Neurosurgery and Psychiatry* 75, 1672–1677.
- Kaur, M., Dang, C.B.S., Singhai, A., Singh, M., Kosta, S., Singh, H., Peter, J., Jain, S., 2014. Toxicity profile of ethanolic extract of *Lawsonia inermis* leaves in albino wistar rats. *World Journal of Pharmacy and Pharmaceutical Sciences* 3, 835–848.
- Kaur, R., Jaiswal, M.L., Jain, V., 2013. Protective effect of *Lannea coromandelica* Houtt. Merrill. against three common pathogens. *Journal of Ayurveda and Integrative Medicine* 4, 224–228.
- Kaushik, D., Khokra, S.L., Kaushik, P., Saneja, A., Arora, D., 2009a. Anticonvulsant activity of *Mitragyna parvifolia* leaves extract. *Pharmacology Online* 3, 101–106.
- Kaushik, D., Khokra, S.L., Kaushik, P., Saneja, A., Sharma, C., Aneja, K.R., Chaudhary, B., Koshy, S., 2009b. A study of analgesic and antimicrobial potential of *Mitragyna parvifolia*. *International Journal of Pharmaceutical Sciences and Drug Research* 1, 6–8.
- Kaushik, D., Saneja, A., Kaushik, P., Lal, S., Yadav, V., 2009c. Antioxidant and anti-inflammatory activities of *Mitragyna parvifolia* leaves extract. *Der Pharmacia Lettre* 1, 75–82.
- Keller, K., Stahl, E., 1983. Composition of the essential oil from beta-asarone free calamus. *Planta Medica* 47, 71–74.
- Khalid, M.D.S., Singh, R.K., Reddy, I.V.N., Kumar, S.J., Kumar, B.S., Kumar, G.N.S., Rao, K.S., 2011. Anti-inflammatory activity of aqueous extract of *Ipomoea carnea* Jacq. *Pharmacology Online* 1, 326–331.
- Khalil, N.M., Sperotto, J.S., Manfron, M.P., 2006. Anti-inflammatory activity of the hydroalcoholic extract of leaves of *Sida rhombifolia* L. (Malvaceae). *Acta Farmaceutica Bonaerense* 25, 260–261.
- Khan, B.A., Akhtar, N., Hussain, I., Abbas, K.A., Rasul, A., 2013. Whitening efficacy of plant extracts including *Hippophae rhamnoides* and *Cassia fistula* extracts on the skin of Asian patients with melasma. *Postepy Dermatologii i Alergologii* XXX, 226–232.
- Khare, C.P., 2007. *Indian Medicinal Plants: An Illustrated Dictionary*. Springer-Verlag, Berlin/Heidelberg.
- Kharmawphlang, S., Sarma, B.P., 2003. Psoriasis and its management by indigenous drug. Bioprospecting of commercially important plants. In: *Proceedings of the National Symposium on "Biochemical approaches for utilization and exploitation of commercially important plants"*. Jorhat, India, 12–14 November 2003, pp. 286–289.
- Khilfi, D., Hayouni, E.A., Valentin, A., Cazaux, S., Moukarzel, B., Hamdi, M., Bouajila, J., 2013. LC–MS analysis, anticancer, antioxidant and antimalarial activities of *Cynodon dactylon* L. extracts. *Industrial Crops and Products* 45, 240–247.
- Khuda, F., Iqbal, Z., Khan, A., Zakiullah, Nasir, F., Shah, Y., 2013. Anti-inflammatory activity of the topical preparation of *Valeriana wallichii* and *Achyranthes aspera* leaves. *Pakistan Journal of Pharmaceutical Sciences* 26, 451–454.
- Khyade, M.S., Vaikos, N., 2011. Comparative phytochemical and antibacterial studies on the bark of *Wrightia tinctoria* and *Wrightia arborea*. *International Journal of Pharma and Bio Sciences* 2, 176–181.
- Kim, H., Chen, F., Wang, X., Chung, H.Y., Jin, Z., 2005. Evaluation of antioxidant activity of Vetiver (*Vetiveria zizanioides* L.) oil and identification of its antioxidant constituents. *Journal of Agricultural and Food Chemistry* 53, 7691–7695.

- Kim, H., Han, T.H., Lee, S.G., 2009. Anti-inflammatory activity of a water extract of *Acorus calamus* L. leaves on keratinocyte HaCaT cells. *Journal of Ethnopharmacology* 122, 149–156.
- Kim, H.Y., Yokozawa, T., Cho, E.J., Cheigh, H.S., Choi, J.S., Chung, H.Y., 2003. *In vitro* and *in vivo* antioxidant effects of mustard leaf (*Brassica juncea*). *Phytotherapy Research* 17 (5), 465–471.
- Kingston, C., Jeeva, S., Jeeva, G.M., Kiruba, S., Mishra, B.P., Kannan, D., 2009. Indigenous knowledge of using medicinal plants in treating skin diseases in Kanyakumari district, Southern India. *Indian Journal of Traditional Knowledge* 8, 196–200.
- Kishore, N., Mishra, A.K., Chansouria, J.P.N., 1993. Fungitoxicity of essential oils against dermatophytes. *Mycoses* 36, 211–215.
- Kowalska, J., Pszczola, K., Wilimowska-Pelc, A., Lorenc-Kubis, I., Zuziak, E., Lugowski, M., Legowska, A., Kwiatkowska, A., Sleszynska, M., Lesner, A., Walewska, A., Zablotna, E., Rolka, K., Wilusz, T., 2007. Trypsin inhibitors from the garden four o'clock (*Mirabilis jalapa*) and spinach (*Spinacia oleracea*) seeds: isolation, characterization and chemical synthesis. *Phytochemistry* 68, 1487–1496.
- Krishanti, M.P., Rathinam, X., Kasi, M., Ayyalu, D., Surash, R., Sadasivam, K., Subramaniam, S., 2010. A comparative study on the antioxidant activity of methanolic leaf extracts of *Ficus religiosa* L., *Chromolaena odorata* (L.), King & Robinson, *Cynodon dactylon* (L.) Pers. and *Tridax procumbens* L. *Asian Pacific Journal of Tropical Medicine*, 348–350.
- Kshirsagar, R., Upadhyay, S., 2009. Free radical scavenging activity screening of medicinal plants from Tripura, Northeast India. *Natural Product Radiance* 8, 117–122.
- Kulkarni, Y.A., Warriar, S., 2012. Acute and repeated dose toxicity studies of methanol extract of *Bauhinia variegata* leaves. *The FASEB Journal* 26, 1124.5.
- Kumar, A., Kashyap, P., 2013. Wound healing activity of *Cynodon dactylon* (L.) Pers. in albino wistar rats. *International Journal of Phytopharmacy* 3, 63–67.
- Kumar, S.G., Malipatil, M., Patil, M.B., 2009. Evaluation of leaves of *Amaranthus spinosus* for antimicrobial and wound healing activity in rats. *Advances in Pharmacology and Toxicology* 10, 71–74.
- Kumar, S.V., Sankar, P., Varatharajan, R., 2009. Anti-inflammatory activity of roots of *Achyranthes aspera*. *Pharmaceutical Biology* 47, 973–975.
- Kumar, B.S.A., Lakshman, K., Jayaveera, K.N., Shekar, D.S., Kumar, A.A., Manoj, B., 2010. Antioxidant and antipyretic properties of methanolic extract of *Amaranthus spinosus* leaves. *Asian Pacific Journal of Tropical Medicine* 3, 702–706.
- Kumar, V., Kumar, R., Yadav, S., Singh, S., Pandeya, S.N., 2010. Evaluation of analgesic & anti-inflammatory activity of hydro-alcoholic extract of *Desmostachya bipinnata* (L.) Stapf root on experimental animals. *International Journal of Pharmaceutical Science and Drug Research* 2, 213–215.
- Kumar, K.A., Sharvanee, S., Patel, J., Choudhary, R.K., 2010. Chemical composition and antimicrobial activity of the essential oil of *Desmostachya bipinnata* Linn. *International Journal of Phytomedicine* 2, 436–439.
- Kumar, H.G., Priyadarshini, V.R., Vinothini, G., Letchoumy, V.P., Nagini, S., 2010. The neem limonoids azadirachtin and nimbolide inhibit cell proliferation and induce apoptosis in an animal model of oral oncogenesis. *Investigational New Drugs* 28, 392–401.
- Kumar, C.M., Sathisha, U.V., Dharmesh, S., Rao, A.G.A., Singh, S.A., 2011. Interaction of sesamol (3,4-methylenedioxyphenol) with tyrosinase and its effect on melanin synthesis. *Biochimie* 93, 562–569.
- Kumar, D.L., Rao, K.N., Madhavi, B., Kumar, D.S., Banji, D., 2011. Anti oxidation activity of *Wrightia tinctoria* Roxb bark and *Schrebera swietenoides* Roxb bark extract. *Journal of Pharmacy Research* 4, 396–397.
- Kumar, V., Tripathi, M.K., Kohli, S., 2011. Antibacterial activity of *Cannabis sativa* against some pathogens isolated from burns of patient. *International Journal of Phytomedicine and Related Industries* 3, 224–243.
- Kumar, P., Vidyasagar, G.M., 2008. Traditional knowledge on medicinal plants used for the treatment of skin diseases in Bidar district, Karnataka. *Indian Journal of Traditional Knowledge* 7, 273–276.
- Kumar, P.K.S., Subramoniam, A., Pushpangadan, P., 2000. Aphrodisiac activity of *Vanda tessellata* (Roxb.) Hook. ex Don extract in male mice. *Indian Journal of Pharmacology* 32, 300–304.
- Kumar, P.S., Sucheta, S., Deepa, V.S., Selvamani, P., Latha, S., 2008. Antioxidant activity in some selected Indian medicinal plants. *African Journal of Biotechnology* 7, 1826–1828.
- Kumar, R., Sankar, R.V., 2003. Ethnobotany of Malyali tribes in Malpattu Village, Javvadu hills of Eastern Ghats, Thiruvannamalai Hills, Tamil Nadu. *Journal of Economic and Taxonomic Botany* 27, 715–726.
- Kumar, S., Jena, P.K., Sabnam, S., Kumari, M., Tripathy, P.K., 2012. Study of plants used against the skin diseases with special reference to *Cassia fistula* L. among the king (Dongaria Kandha) of Niyamgiri: a primitive tribe of Odisha, India. *International Journal of Drug Development and Research* 4, 256–264.
- Kumar, V.L., Basu, N., 1994. Anti-inflammatory activity of the latex of *Calotropis procera*. *Journal of Ethnopharmacology* 44, 123–125.
- Kumari, P., Singh, G.S., 2009. Ethnobotanical study of medicinal plants used by the Taungya community in Terai Arc Landscape, India. *Journal of Ethnopharmacology* 123, 167–176.
- Kundu, M., Das, S., Kumari, S., Basu, S.P., 2011. Studies on the aqueous matured leaf extract of *Bambusa bambos* “Druce” as a potent hypothermic agent. *IJPI'S Journal of Pharmacology and Toxicology* 1, 54–60.
- Kuvar, S.D., Bapat, U.C., 2010. Medicinal plants used by Kokani tribals of Nasik district Maharashtra to cure cut and wounds. *Indian Journal of Traditional Knowledge* 9, 114–115.
- Lal, B., Singh, K.N., 2008. Indigenous herbal remedies to cure skin disorders by the natives of Lahaul-Spiti in Himachal Pradesh. *Indian Journal of Traditional Knowledge* 7, 237–241.
- Lal, B., Vats, S.K., Singh, R.D., Gupta, A.K., 1996. Plants used as ethnomedicine and supplementary food by Gaddis of Himachal Pradesh, India. In: Jain, S.K. (Ed.), *Ethnobiology in Human Welfare*. Deep Publications, New Delhi, pp. 384–387.
- Lalla, J.K., Nandedkar, S.Y., Paranjape, M.H., Talreja, N.B., 2001. Clinical trials of ayurvedic formulations in the treatment of acne vulgaris. *Journal of Ethnopharmacology* 78, 99–102.
- Lam, F.Y., Ferrell, W.R., 1991. Neurogenic component of different models of acute inflammation in the rat knee joint. *Annals of the Rheumatic Diseases* 50, 747–751.
- Lardos, A., Heinrich, M., 2013. Continuity and change in medicinal plant use: the example of monasteries on Cyprus and historical *iatriosofia* texts. *Journal of Ethnopharmacology* 150, 202–214.
- Laupattarakasem, P., Wangsrimongkol, T., Surarit, R., Hahnvanjanawong, C., 2006. *In vitro* and *in vivo* anti-inflammatory potential of *Cryptolepis buchanani*. *Journal of Ethnopharmacology* 108, 349–354.
- Lavhale, M.S., Mishra, S.H., 2007. Evaluation of free radical scavenging activity of *Butea monosperma* Lam. *Indian Journal of Experimental Biology* 45, 376–384.
- Leary, W.P., Reyes, A.J., van der Byl, K., 1986. Effects of single doses of two antihypertensive rauwolfia-diuretic combinations on urinary excretion. *South African Medical Journal* 70, 95–98.
- Lee, K.T., Kim, B.J., Kim, J.H., Heo, M.Y., Kim, H.P., 1997. Biological screening of 100 plant extracts for cosmetic use (I): inhibitory activities of tyrosinase and DOPA auto-oxidation. *International Journal of Cosmetic Science* 19, 291–298.
- Leonti, M., Cabras, S., Weckerle, C.S., Solinas, M.N., Casu, L., 2010. The causal dependence of present plant knowledge on herbals – contemporary medicinal plant use in Campania (Italy) compared to Matthioli (1568). *Journal of Ethnopharmacology* 130, 379–391.
- Leonti, M., Casu, L., Sanna, F., Bonsignore, L., 2009. A comparison of medicinal plant use in Sardinia and Sicily – De Materia Medica revisited? *Journal of Ethnopharmacology* 121, 255–267.
- Leonti, M., Nebel, S., Rivera, D., Heinrich, M., 2006. Wild gathered food plants in the European Mediterranean: a comparative analysis. *Economic Botany* 60, 130–142.
- Li, R.W., Leach, D.N., Myers, S.P., Lin, G.D., Leach, G.J., Waterman, P.G., 2004. A new anti-inflammatory glucoside from *Ficus racemosa* L. *Planta Medica* 70, 421–426.
- Lin, C.C., Cheng, H.Y., Yang, C.M., Lin, T.C., 2002. Antioxidant and antiviral activities of *Euphorbia thymifolia* L. *Journal of Biomedical Science* 9, 656–664.
- Lodhi, S., Singhi, A.K., 2011. Preliminary pharmacological evaluation of *Martynia annua* Linn. Leaves for wound healing. *Asian Pacific Journal of Tropical Biomedicine* 1, 421–427.
- Luqman, S., Kumar, R., Kaushik, S., Srivastava, S., Darokar, M.P., Khanuja, S.P.S., 2009. Antioxidant potential of the root of *Vetiveria zizanioides* (L.) Nash. *Indian Journal of Biochemistry & Biophysics* 46, 122–125.
- Macwan, C., Patel, H.V., Kalia, K., 2010. A comparative evaluation of *in vitro* antioxidant properties of Bamboo *Bambusa arundinacea* leaves extracts. *Journal of Cell and Tissue Research* 10, 2413–2418.
- Mahat, M.A., Patil, B.M., 2007. Evaluation of anti-inflammatory activity of methanol extract of *Phyllanthus amarus* in experimental animal models. *Indian Journal of Pharmaceutical Sciences* 69, 33–36.
- Mahesh, A.R., Kumar, H., Ranganath, M.K., Devkar, R.A., 2012. Detail study on *Boerhavia diffusa* plant for its medicinal importance – a review. *Research Journal of Pharmaceutical Sciences* 1, 28–36.
- Mahida, Y., Mohan, J.S.S., 2007. Screening of plants for their potential antibacterial activity against *Staphylococcus* and *Salmonella* spp. *Natural Product Radiance* 6, 301–305.
- Mahire, N.B., Tote, M.V., Jain, A.P., Undale, V.R., Bhosle, A.V., 2009. Anti-inflammatory effect of *Premna latifolia* leaves. *Pharmacology Online* 3, 929–937.
- Mainasara, M., Aliero, B., Yakubu, A., 2012. Phytochemical and antibacterial properties of root and leaf extracts of *Calotropis procera*. *Nigerian Journal of Basic and Applied Sciences* 20, 1–6.
- Maioli, E., Torricelli, C., Valacchi, G., 2012. Rottlerin and cancer: novel evidence and mechanisms. *The Scientific World Journal* 12, 350826.
- Mairh, A.K., Mishra, P.K., Kumar, J., Mairh, A., 2010. Traditional botanical wisdom of Birhore tribes of Jharkhand. *Indian Journal of Traditional Knowledge* 9, 467–470.
- Maity, T.K., Mandal, S.C., Mukherjee, P.K., Saha, K., Das, J., Pal, M., Saha, B.P., 1998. Studies on anti-inflammatory effect of *Cassia tora* leaf extract (fam. Leguminosae). *Phytotherapy Research* 12, 21–23.
- Maiyo, Z.C., Ngure, R.M., Matasyoh, J.C., Chepkorir, R., 2010. Phytochemical constituents and antimicrobial activity of leaf extracts of three *Amaranthus* plant species. *African Journal of Biotechnology* 9, 3178–3182.
- Majumdar, K., Datta, B.K., 2007. A study on ethnomedicinal usage of plants among the folklore herbalists and Tripuri medical practitioners, Part II. *Natural Product Radiance* 6, 66–73.
- Majumder, R., Jami, M.S.I., Alam, M.E.K., Alam, M.B., 2013. Antidiarrheal activity of *Lannea coromandelica* Linn. Bark extract. *American–Eurasian Journal of Scientific Research* 8, 128–134.
- Malan, R., Walia, A., Saini, V., Gupta, S., 2011. Comparison of different extracts leaf of *Brassica juncea* Linn. on wound healing activity. *European Journal of Experimental Biology* 1, 33–40.
- Malarvili, T., Gomathi, N., 2009. Effect of *Achyranthes aspera* (Linn.) seeds on redox and oxidative status in plasma and selected tissues of rats fed with high doses of fructose. *Biosciences, Biotechnology Research Asia* 6, 659–664.

- Mali, P.Y., Panchal, S.S., 2013. A review on phyto-pharmacological potentials of *Euphorbia thymifolia* L. *Ancient Science of Life* 32, 165–172.
- Malik, N., Dhawan, V., Bahl, A., Kaul, D., 2009. Inhibitory effects of *Terminalia arjuna* on platelet activation *in vitro* in healthy subjects and patients with coronary artery disease. *Platelets* 20, 183–190.
- Mamatha, G.C., Prabhakar, T., Madhuri, V., Neelima, K., Venkatanagaraju, E., Chandrasekar, S.B., 2014. Anti-arthritis activity of *Euphorbia thymifolia* Linn. *World Journal of Pharmacy and Pharmaceutical Sciences* 3, 1323–1331.
- Mamatha, N., Pavan, Keshava Murthy, K.R., Venkatesh, D.A., 2006. Data on 100 medicinal plants used by Soligas of Biligirirangana hills of Mysore District, Karnataka. *My Forest* 42, 121–139.
- Mamgain, P., Singh, R.H., 1994. Controlled clinical trial of the lekhania drugvaca (*Acorus calamus*) in cases of Ischaemic heart diseases. *Journal of Research in Ayurveda and Siddha* 15, 35–51.
- Manalisha, D., Chandra, K.J., 2012. Preliminary phytochemical analysis and acute oral toxicity study of *Mucuna pruriens* Linn. in albino mice. *International Research Journal of Pharmacy* 3, 181–183.
- Mandal, S., Kundu, P., Roy, B., Mandal, R.K., 2002. Precursor of the inactive 2S seed storage protein from the Indian mustard *Brassica juncea* is a novel trypsin inhibitor. Characterization, post-translational processing studies, and transgenic expression to develop insect-resistant plants. *The Journal of Biological Chemistry* 277, 37161–37168.
- Mandal, S.C., Maity, T.K., Das, J., Saba, B.P., Pal, M., 2000. Anti-inflammatory evaluation of *Ficus racemosa* Linn. leaf extract. *Journal of Ethnopharmacology* 72, 87–92.
- Mangal, A., Sharma, M.C., 2009. Evaluation of certain medicinal plants for antiobesity properties. *Indian Journal of Traditional Knowledge* 8, 602–605.
- Manikandan, S., Devi, R.S., 2005. Antioxidant property of α -asarone against noise stress-induced changes in different regions of rat brain. *Pharmacological Research* 52, 467–474.
- Mantle, D., Gok, M.A., Lennard, T.W., 2001. Adverse and beneficial effects of plant extracts on skin and skin disorders. *Adverse Drug Reactions and Toxicological Reviews* 20, 89–103.
- Manu, K.A., Kuttan, G., 2007. Effect of Punarnavine, an alkaloid from *Boerhaavia diffusa*, on cell-mediated immune responses and TIMP-1 in B16F-10 metastatic melanoma-bearing mice. *Immunopharmacology and Immunotoxicology* 29, 569–586.
- Marques-Neto, J.F., Lapa, A., Kubota, M., 1988. Efeitos de "Ageratum conyzoides, Linee" no tratamento da artrose. *Revista Brasileira de Reumatologia* 28, 109–114.
- Mary, K.V., 2011. Studies on phytochemical screening and antibacterial Activities of *Lantana camara* linn. *Plant Science Feed* 1, 74–79.
- Mas-Chamberlin, C., Mondon, P., Lamy, F., Peschard, O., Lintner, K., 2005. Reduction of hair-loss: matrilines and plant molecules to the rescue. In: *Proceedings of the 7th Scientific Conference of the Asian Society of Cosmetic Chemists*, Bangkok, Thailand.
- MatAli, R., Houghton, P.J., Raman, A., Hout, J.R., 1998. Antimicrobial and anti-inflammatory activities of extracts and constituents of *Oroxylum indicum* (L.) Vent. *Phytomedicine* 5, 375–381.
- Matpal, M., Agarwal, K., Saini, P., 2013. *In vitro* anti-inflammatory activity of *Ficus benghalensis* bark. *International Research Journal of Pharmacy* 4, 107–108.
- Mazid, M.A., Datta, B.K., Nahar, L., Bashir, S.A.M.K., Bachar, S.C., Sarker, S.D., 2009. Antinociceptive, anti-inflammatory and diuretic properties of *Polygonum barbatum* (L.) Hara var. *barbata*. *Revista Brasileira de Farmacognosia* 19, 749–754.
- Meechai, I., Puripattavanong, J., Dej-adisai, S., 2010. Screening of anti-tyrosinase activity from Thai medicinal plants. In: *The 9th NRCT-JSPS Joint Seminar. Natural Medicine Research for the Next Decade: New Challenges and Future Collaboration*. December 8–9 2010, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok, Thailand. pp. 119–121.
- Meena, K.L., Yadav, B.L., 2011. Some ethnomedicinal plants used by the Garasia tribe of district Sirohi, Rajasthan. *Indian Journal of Traditional Knowledge* 10, 354–357.
- Mehdinezhad, B., Rezaei, A., Mohajeri, D., Safarmashaei, S., 2012. Comparison of *in vivo* wound healing activity of *Verbascum thapsus* flower extract with zinc oxide on experimental wound model in Rabbits. *American-Eurasian Journal of Toxicological Sciences* 4, 24–30.
- Mehta, F.A., Patel, B.G., Pandya, S.S., Ahir, K.B., Patel, S.B., 2009. Antinociceptive and antiinflammatory activity of *Achyranthes aspera* L. extracts. *Pharmacology Online* 3, 978–985.
- Minh, H.C., Tuan, N.A., 2013. Study on toxicity of the plant *Cryptolepis buchananii* on experimental animals. *Journal of Food and Nutrition Sciences* 9, 1.
- Mirmiran, P., Bahadoran, Z., Golzarand, M., Rajab, A., Azizi, F., 2013. Ardeh (*Sesamum indicum*) could improve serum triglycerides and atherogenic lipid parameters in Type 2 Diabetic patients: a randomized clinical trial. *Archives of Iranian Medicine* 16, 651–656.
- Mishra, A., Sharma, A.K., Kumar, S., Saxena, A.K., Pandey, A.K., 2013. *Bauhinia variegata* leaf extracts exhibit considerable antibacterial, antioxidant and anticancer activities (Article ID 915436). *BioMed Research International*, 10.
- Mishra, S.B., Verma, A., Mukerjee, A., Vijayakumar, M., 2012. *Amaranthus spinosus* L. (Amaranthaceae) leaf extract attenuates streptozotocin-nicotinamide induced diabetes and oxidative stress in albino rats: A histopathological analysis. *Asian Pacific Journal of Tropical Biomedicine*, S1647–S1652.
- Mishra, P.K., Shukla, R., Singh, P., Prakash, B., Dubey, N.K., 2012. Antifungal and antiflatulogenic efficacy of *Caesulia axillaris* Roxb. essential oil against fungi deteriorating some herbal raw materials, and its antioxidant activity. *Industrial Crops and Products* 36, 74–80.
- Mishra, S.L., Sinhamahapatra, P.K., Nayak, A., Das, R., Sannigrahi, S., 2010. *In vitro* antioxidant potential of different parts of *Oroxylum indicum*: a comparative study. *Indian Journal of Pharmaceutical Sciences* 72, 267–269.
- Mistry, S., Dutt, K.R., Jena, J., 2013. Protective effect of *Sida cordata* leaf extract against CCl₄ induced acute liver toxicity in rats. *Asian Pacific Journal of Tropical Medicine* 3, 280–284.
- Mohammad, F., Mazumder, A., 2013. Wound healing potentiality of methanolic extract of aerial parts of *Cleome viscosa*. *International Journal of PharmTech Research* 5, 978–982.
- Mojahid-ul-Islam, Elhddad, S., 2012. Phytochemical investigation and evaluation of analgesic activity of ethanolic extract of *Dalbergia sissoo* (Roxb.) bark. *Journal of Natural Product and Plant Resources* 2, 701–704.
- Mokhtarian, F., Griffin, D.E., 1984. The role of mast cells in virus-induced inflammation in the murine central nervous system. *Cell Immunology* 86, 491–500.
- Monteiro, E.M.H., Chibli, L.A., Yamamoto, C.H., Pereira, M.C.S., Vilela, F.M.P., Rodarte, M.P., Pinto, M.A.D.O.P., Amaral, M.D.P.H.D., Silverio, M.S., Araujo, A.L.S.D.M., Araujo, A.D.L.A.D., Del-Vechio-Vieira, G., Sousa, O.V.D., 2014. Antinociceptive and anti-inflammatory activities of the Sesame oil and Sesamin. *Nutrients* 6, 1931–1944.
- Moreman, D.E., 1996. An analysis of the food plants and drug plants of Native North America. *Journal of Ethnopharmacology* 52, 1–22.
- Morton, J.F., 1962. Ornamental plants with toxic and/or irritant properties. II. *Proceedings of Florida State Horticulture Society* 75, 484–491.
- Mozaffarpur, S.A., Naseri, M., Esmailidooki, M.R., Kamalinejad, M., Bijani, A., 2012. The effect of cassia fistula emulsion on pediatric functional constipation in comparison with mineral oil: a randomized, clinical trial. *Daru* 30, 83.
- Mukherjee, P.K., Badami, S., Wahile, A.M., Rajan, S., Suresh, B., 2001. Evaluation of tyrosinase inhibitory activity of some Indian spices. *Journal of Natural Remedies* 1/2, 125–129.
- Muniappan, M., Sundararaj, T., 2003. Anti-inflammatory and antiulcer activities of *Bambusa arundinacea*. *Journal of Ethnopharmacology* 88, 161–167.
- Muniappan, M.J., Senthilkumar, J., Inbaraj, S.D., 2014. Evaluation of *Bambusa arundinacea* leaves for wound healing in albino rats. *Global Journal of Pharmacology* 8, 84–86.
- Muralidhar, A., Babu, K.S., Sankar, T.R., Reddanna, P., Latha, J., 2013. Wound healing activity of flavonoid fraction isolated from the stem bark of *Butea monosperma* (Lam) in albino Wistar rats. *European Journal of Experimental Biology* 3, 1–6.
- Murthy, K.S.R., Lakshmi, N., Ramulu, D.R., 2011. Biological activity and phytochemical screening of the oleoresin of *Shorea robusta* Gaertn.f. *Tropical and Subtropical Agroecosystems* 14, 787–791.
- Murti, K., Kumar, U., 2011. Antimicrobial activity of *Ficus benghalensis* and *Ficus racemosa* roots L. *American Journal of Microbiology* 2, 21–24.
- Murti, K., Kumar, U., 2012. Enhancement of wound healing with roots of *Ficus racemosa* L. in albino rats. *Asian Pacific Journal of Tropical Biomedicine* 2, 276–280.
- Murti, K., Lambale, V., Panchal, M., Shah, M., 2011. Healing potential of roots of *Scoparia dulcis* in Albino rats. *Pharmacologia* 2, 369–373.
- Murugan, M., Mohan, V.R., 2011. Antibacterial activity of *Mucuna pruriens* (L.) Dc. var. *pruriens*—an ethnomedicinal plant. *Science Research Reporter* 1, 69–72.
- Musa, E.M., El Badwi, S.M., Jah Elnabi, M.A., Osman, E.A., Dahab, M.M., 2011. Anti-inflammatory activity of the plant *Cannabis sativa* (L) petroleum ether extract in albino rats. *Research in Pharmacy* 1, 18–25.
- Muthu, C., Ayyanar, M., Nagappan, R., Ignacimuthu, S., 2006. Medicinal plants used by traditional healers in Kancheepuram District of Tamil Nadu, India. *Journal of Ethnobiology and Ethnomedicine* 2, 43. <http://dx.doi.org/10.1186/1746-4269-2-43>.
- Muthuraman, A., Singh, N., 2012. Acute and sub-acute oral toxicity profile of *Acorus calamus* (Sweet flag) in rodents. *Asian Pacific Journal of Tropical Biomedicine*, S1017–S1023.
- Nadella, D., Paarakh, P.M., 2011. Antibacterial activity of different extracts of stem bark of *Holoptelea integrifolia* Roxb (Planch). *International Research Journal of Pharmacy* 2, 111–113.
- Nadkarni, A.K., 1976. Dr. K.M. Nadkarni's Indian Materia Medica. With Ayurvedic, Unani-Tibbi, Siddha, Allopathic, Homeopathic, Naturopathic & Homo Remedies, Appendices & Indexes, Revised enlarged and reprinted 3rd edition, vols. 1 & 2. Popular Prakashan, Bombay, p. 636.
- Nagarani, G., Abirami, A., Siddhuraju, P., 2014. A comparative study on antioxidant potentials, inhibitory activities against key enzymes related to metabolic syndrome, and anti-inflammatory activity of leaf extract from different *Momordica* species. *Food Science and Human Wellness* 3, 36–46.
- Nagda, D., Saluja, A., Nagda, C., 2009. Antioxidant activities of methanolic and aqueous extract from leaves of *Martynia annua* Linn. *Pharmacognosy Journal* 1, 288–297.
- Nahar, L., Nasrin, F., Zahan, R., Mosaddik, Md. A., 2013. Anti-nociceptive and anti-inflammatory activities of *Wrightia arborea*. *Pakistan Journal of Biological Sciences* 16, 485–490.
- Nair, P.R., Vijayan, N.P., Madhavikutty, P., Nair, C.N.B., Chandrika, A., 1988. Clinical evaluation of sahacharadi and nirgundi tailas in saisaveeyavata (Poliomyelitis). *Ancient Science of Life* VIII, 25–29.
- Nandagopalan, V., Anand, S.P., Laxmi prabha, A., Selvakumar, U., Doss, A., 2011. An ethnobotanical study in the Pudukkottai District, South India. *Asian Journal of Experimental Biological Sciences* 2, 412–421.
- Nantitanon, W., Chowwanapoonpohn, S., Okonogi, S., 2007. Antioxidant and antimicrobial activities of *Hyptis suaveolens* essential oil. *Scientia Pharmaceutica* 75, 35–46.

- Nayak, B.S., Raju, S.S., Eversley, M., Ramsabhadra, A., 2009. Evaluation of wound healing activity of *Lantana camara* L. – a preclinical study. *Phytotherapy Research* 23, 241–245.
- Nayak, B.S., Suresh, R., Rao, A.V., Pillai, G.K., Davis, E.M., Ramkissoon, V., McRae, A., 2005. Evaluation of wound healing activity of *Vanda roxburghii* R.Br (Orchidaceae): a preclinical study in a rat model. *The International Journal of Lower Extremity Wounds* 4, 200–204.
- Nayak, S., Behera, S.K., Nisha, M., 2004. Ethnomedicinal botanical survey of Kalahandi District of Orissa. *Indian Journal of Traditional Knowledge* 3, 72–79.
- Naz, R., Bano, A., 2012. Antimicrobial potential of *Ricinus communis* leaf extract in different solvents against pathogenic bacterial and fungal strains. *Asian Pacific Journal of Tropical Biomedicine* 2, 944–947.
- Naz, R., Bano, A., 2013. Phytochemical screening, antioxidants and antimicrobial potential of *Lantana camara* in different solvents. *Asian Pacific Journal of Tropical Disease* 3, 480–486.
- Negi, C.S., Nautiyal, S., Dasila, L., Rao, K.S., Maikhuri, R.K., 2002. Ethnomedicinal plant uses in a small tribal community in a part of Central Himalaya, India. *Journal of Human Ecology* 14, 23–31.
- Negi, V.S., Maikhuri, R.K., Vashishtha, D.P., 2011. Traditional healthcare practices among the villages of Rawain valley, Uttarakash, Uttarakhand, India. *Indian Journal of Traditional Knowledge* 10, 533–537.
- Neto, L.G.D.N., Pinto, L.D.S., Bastos, R.M., Evaristo, F.F.V., Vasconcelos, M.A.D., Carneiro, V.A., Arruda, F.V.S., Porto, A.L.F., Leal, R.B., Junior, V.A.D.S., Cavada, B. S., Teixeira, E.H., 2011. Effect of the Lectin of *Bauhinia variegata* and its recombinant Isoform on surgically induced skin wounds in a Murine model. *Molecules* 16, 9298–9315.
- Nikam, P.S., Nikam, S.V., Sontakke, A.V., Khanwelkar, C.C., 2011. Role of *Phyllanthus amarus* treatment in Hepatitis-C. *Biomedical Research* 22, 319–322.
- Nishino, H., Hayashi, T., Arisawa, M., Satomi, Y., Iwashima, A., 1993. Antitumor-promoting activity of scopadulcic acid B, isolated from the medicinal plant *Scoparia dulcis* L. *Oncology* 50, 100–103.
- Nithya, V., Brinda, P., Anand, K.V., 2011. Wound healing activity of *Leonotis nepetaefolia* R.Br., in Wistar Albino rats. *Asian Journal of Pharmaceutical and Clinical Research* 4, 23–26.
- Notka, F., Meier, G.R., Wagner, R., 2004. Concentrated inhibitory activities of *Phyllanthus amarus* on HIV replication *in vitro* and *ex vivo*. *Antiviral Research* 64, 93–102.
- Nurul, I.M., Mizuguchi, H., Shahriar, M., Venkatesh, P., Maeyama, K., Mukherjee, P.K., Hattori, M., Choudhuri, M.S.K., Takeda, N., Fukui, H., 2011. *Albizia lebbek* suppresses histamine signaling by the inhibition of histamine H1 receptor and histidine decarboxylase gene transcriptions. *International Immunopharmacology* 11, 1766–1772.
- Oberdorfer, S., Grisold, W., Hinterholzer, G., Rosner, M., 2002. Coma with focal neurological signs caused by *Datura stramonium* intoxication in a young man. *Journal of Neurology, Neurosurgery & Psychiatry* 73, 458–459.
- Obogwu, M.B., Akindele, A.J., Adeyemi, O.O., 2014. Hepatoprotective and *in vivo* antioxidant activities of the hydroethanolic leaf extract of *Mucuna pruriens* (Fabaceae) in antitubercular drugs and alcohol models. *Chinese Journal of Natural Medicines* 12, 273–283.
- Ono, M., Nishida, Y., Masuoka, C., Li, J.C., Okawa, M., Ikeda, T., 2004. Lignan derivatives and a norditerpene from the seeds of *Vitex negundo*. *Journal of Natural Products* 67, 2073–2075.
- Onuoha, O.G., Ayo, J.A., Osuagwu, V., Iruolaje, F.O., 2013. Investigation of the antibacterial activity of *Ageratum conyzoides* extract on microorganisms isolated from septic wound. *Topclass Journal of Herbal Medicine* 2, 182–188.
- Orisakwe, O.E., Afonne, O.J., Chude, M.A., Obi, E., Dioka, C.E., 2003. Sub-chronic toxicity studies of the aqueous extract of *Boerhavia diffusa* leaves. *Journal of Health Science* 49, 444–447.
- Ouedraogo, G.G., Ouedraogo, M., Lamien-Sanou, A., Lompo, M., Gombri-Lompo, O. M., Guissou, P.I., 2013. Acute and subchronic toxicity studies of roots barks extracts of *Calotropis procera* (Ait.) R. Br used in the treatment of sickle cell disease in Burkina Faso. *British Journal of Pharmacology and Toxicology* 4, 194–200.
- Paarakh, P.M., 2010. *Terminalia arjuna* (Roxb.) Wt. and Arn.: a review. *International Journal of Pharmacology* 6, 515–534.
- Pachkore, G.L., Dhale, D.A., Dharasurkar, A.N., 2011. Antimicrobial and phytochemical screening of *Hyptis suaveolens* (L.Poit) Lamiaceae. *International Multidisciplinary Research Journal* 1, 01–03.
- Pahwa, R., Chatterjee, V.C., 1988. The toxicity of Indian *Calotropis procera* R.Br latex in the black rat, *Rattus rattus* Linn. *Veterinary and Human Toxicology* 30, 305–308.
- Pal, A., Sharma, P.P., Mukherjee, P.K., 2009. A clinical study of Kutaja (*Holarrhena antidysenterica* Wall.) on Shonitarsha. *AYU* 30, 369–372.
- Pammel, L.H., 1911. A Manual of Poisonous Plants. Chiefly of North America, with Brief Notes on Economic and Medicinal Plants, and Numerous Illustrations. Torch Press, Cedar Rapids, IA.
- Pandey, P.K., Jamal, F., 2010. Identification and evaluation of Trypsin inhibitor in the seed extracts of *Butea monosperma* (Flame of Forest). *International Journal of Biotechnology and Biochemistry* 6, 513–520.
- Panduraju, T., Parvathi, B., Rammoan, M., Reddy, C.S., 2011. Wound healing properties of *Cleome viscosa* Linn. Hygeia – Journal for Drugs and Medicines 3, 41–45.
- Parhar, K., Biswas, S., 1998. Medicinal plant resources of Doon valley and adjacent Siwaliks – I. *Journal of Non-Timber Forest Products* 5, 29–33.
- Parimala, S., Shashidhar, Gh., Sridevi, Ch., Jyothi, V., Suthakaran, R., 2009. Anti-inflammatory activity of *Celastrus paniculatus* seeds. *International Journal of PharmTech Research* 1, 1326–1329.
- Parkash, V., Sandhu, P., 2012. *In vitro* antimycobiotic and antibacterial action of seed extract of *Celastrus paniculatus* Willd. (Jyotismati). *Journal of Antimicrobials* 127, 123–132.
- Parmar, M.Y., Shah, P.A., Thakkar, V.T., Gandhi, T.R., 2008. Hepatoprotective activity of *Vetiveria zizanioides* Linn. against ethanol-induced liver damage in rats. *Pharmacognosy Magazine* 4, S216–S221.
- Parra-Delgado, H., Ruiz, G.G., Camacho, A.N., Martínez-Vázquez, M., 2004. Anti-inflammatory activity of some extracts and isolates from *Leonotis nepetaefolia* on TPA-induced Edema Model. *Revista de la Sociedad Química de México* 48, 293–295.
- Parthasarthy, J.S., Ravichandran, G., Gladstone, M.S.B., Pande, P.S., 2009. Safety and efficacy study of nourishing baby oil in baby massage. *Indian Medical Journal* 103, 209–213.
- Patel, J.R., Tripathi, P., Sharma, V., Chauhan, N.S., Dixit, V.K., 2011. *Phyllanthus amarus*: ethnomedicinal uses, phytochemistry and pharmacology: a review. *Journal of Ethnopharmacology* 138, 286–313.
- Patel, R.P., Trivedi, B.M., 1962. The *in vitro* antibacterial activity of some medicinal oils. *Indian Journal of Medical Research* 50, 218–222.
- Patel, V.R., Patel, P.R., Kajal, S.S., 2010. Antioxidant activity of some selected medicinal plants in western region of India. *Advances in Biological Research* 4, 23–26.
- Patil, B.S., Naikwade, N.S., Kondawar, M.S., Magdum, C.S., Awale, V.B., 2009. Traditional uses of plants for wound healing in the Sangli district, Maharashtra. *International Journal of PharmaTech Research* 3, 876–878.
- Patil, P.R., Patil, V.M., Verma, S., Vijayanath, V., Surpur, R.R., 2012. A study of anti-inflammatory activity of *Azadirachta indica* (Neem) root in Wistar albino rats. *Medico-Legal Update* 12, 104–106.
- Patil, R.P., Nimbalkar, M.S., Jadhav, U.U., Dawkar, V.V., Govindwar, S.P., 2010. Antiaflatoxicogenic and antioxidant activity of an essential oil from *Ageratum conyzoides* L. *Journal of the Science of Food and Agriculture* 90, 608–614.
- Patil, S., Anarthe, S., Jadhav, R., Surana, S., 2011. Evaluation of anti-inflammatory activity and *in-vitro* antioxidant activity of Indian Mistletree, the Hemiparasite *Dendrophthoe falcata* L. F. (Loranthaceae). *Iran Journal of Pharmacy Research* 10, 253–259.
- Patil, U.H., Gaikwad, D.K., 2011. Phytochemical evaluation and bactericidal potential of *Terminalia arjuna* stem bark. *International Journal of Pharmaceutical Sciences and Research* 2, 614–619.
- Pattanayak, S.P., Sunita, P., 2008. Wound healing, anti-microbial and antioxidant potential of *Dendrophthoe falcata* (L.f) Ettingsh. *Journal of Ethnopharmacology* 120, 241–247.
- Pattnaik, A., Sarkar, R., Sharma, A., Yadav, K.K., Kumar, A., Roy, P., Mazumder, A., Karmakar, S., Sen, T., 2013. Pharmacological studies on *Buchanania lanzan* Spreng. – a focus on wound healing with particular reference to anti-biofilm properties. *Asian Pacific Journal of Tropical Biomedicine* 3, 967–974.
- Pawar, C.R., Kolhe, V.N., Khedhar, P.A., 2011. Anti-inflammatory activity of *Typha angustata* (Typhaceae). *International Journal of Research in Ayurveda and Pharmacy* 2, 1598–1600.
- Pawlikowska-Pawlega, B., Gawron, A., 1995. Effect of quercetin on the growth of mouse fibroblast cells *in vitro*. *Polish Journal of Pharmacology* 47, 531–535.
- Peng, H., Lai, C., Lin, C., Chou, S., 2014. Effect of *Vetiveria zizanioides* essential oil on melanogenesis in melanoma cells: downregulation of tyrosinase expression and suppression of oxidative stress (Article ID 213013). *The Scientific World Journal*, 9.
- Peng, L., Liu, S., Xu, S., Chen, L., Shan, Y., Wei, W., Liang, W., Gao, J., 2013. Inhibitory effects of salidroside and paeonol on tyrosinase activity and melanin synthesis in mouse B16F10 melanoma cells and ultraviolet B-induced pigmentation in guinea pig skin. *Phytotherapy* 20, 1082–1087.
- Perry, L.M., Metzger, J., 1980. Medicinal plants of East and Southeast Asia. MIT Press, Cambridge, MA p. 620.
- Perumal, P., Sekar, V., Rajesh, V., Gandhimathi, S., Kumar, R.S., Nazimudin, K.H.S., 2010. *In vitro* antioxidant activity of *Argemone mexicana* roots. *International Journal of PharmTech Research* 2, 1477–1482.
- Phillips, O., Gentry, A.H., Reynel, C., Wilkin, P., Galvez-Durand, C.B., 2002. Quantitative ethnobotany and Amazonian conservation. *Conservation Biology* 8, 225–248.
- Pingale, S.S., 2013. Toxicity study for *Argemone mexicana* L. *World Journal of Pharmaceutical Sciences* 1, 151–155.
- Policegoudra, R.S., Chattopadhyay, P., Aradya, S.M., Shivaswamy, R., Singh, L., Veer, V., 2014. Inhibitory effect of *Tridax procumbens* against human skin pathogens. *Journal of Herbal Medicine* 4, 83–88.
- Policepatel, S.S., Manikrao, V.G., 2013. Ethnomedicinal plants used in the treatment of skin diseases in Hyderabad Karnataka region, Karnataka, India. *Asian Pacific Journal of Tropical Biomedicine* 3, 882–886.
- Ponrasu, T., Suguna, L., 2012. Efficacy of *Annona squamosa* on wound healing in streptozotocin-induced diabetic rats. *International Wound Journal* 9, 613–623.
- Poornima, R., Prabakaran, G., 2012. Preliminary phytochemical screening and antibacterial activity of *Acalypha indica* and *Euphorbia hirta* of family Euphorbiaceae against some pathogenic organisms. *International Journal of Agricultural Sciences* 2, 34–38.
- Pour, B.M., Latha, L.Y., Sasidharan, S., 2011. Cytotoxicity and oral acute toxicity studies of *Lantana camara* leaf extract. *Molecules* 16, 3663–3674.

- Prabha, M.R., Vasantha, K., 2012. Phytochemical and antibacterial activity of *Calotropis procera* (ait.) R. Br. flowers. *International Journal of Pharma and Bio Sciences* 3, 1–6.
- Pradhan, B.K., Badola, H.K., 2008. Ethnomedicinal plant use by Lepcha tribe of Dzongu valley, bordering Khangchendzonga Biosphere Reserve, in North Sikkim, India. *Journal of Ethnobiology and Ethnomedicine* 4, 22. <http://dx.doi.org/10.1186/1746-4269-4-22>.
- Prakash, D., Niranjana, A., Tewari, S.K., 2001. Some nutritional properties of the seeds of three *Mucuna* species. *International Journal of Food Sciences and Nutrition* 52, 79–82.
- Prakash, D., Singh, B.N., Upadhyay, G., 2007. Antioxidant and free radical scavenging activities of phenols from onion (*Allium cepa*). *Food Chemistry* 102, 1389–1393.
- Prakasha, H.M., Krishnappa, M., 2006. People's knowledge on medicinal plants in Shringeri taluk, Karnataka. *Indian Journal of Traditional Knowledge* 5, 352–357.
- Pranjpe, P., Kulkarni, P.H., 1995. Comparative efficacy of four ayurvedic formulations in the treatment of acne vulgaris: a double-blind randomised placebo-controlled clinical evaluation. *Journal of Ethnopharmacology* 49, 127–132.
- Prasad, M.P., Sushant, S., Chikkaswamy, B.K., 2012. Phytochemical analysis, antioxidant potential, antibacterial activity and molecular characterization of *Clerodendrum* species. *International Journal of Molecular Biology* 3, 71–76.
- Prasad, M.S., Venkateshwarlu, G., Dhanalakshmi, C.H., Kumar, D.S., Alekha, K., Kumar, B.P., Rajkumar, J.V., 2011. Wound healing activity of *Pongamia pinnata* in albino wistar rats. *Research Journal of Pharmaceutical, Biological and Chemical Sciences* 2, 1096–1100.
- Prasad, N., Nandi, D., Arora, S., Pandey, A., 2014. *In vitro* evaluation of antibacterial properties of *Moringa oleifera*, *Dalbergia sissoo* and *Alstonia scholaris*. *Journal of Biology, Agriculture and Healthcare* 4, 54–62.
- Premalatha, B., 2000. *Semecarpus anacardium* Linn. Nuts – a boon in alternative medicine. *Indian Journal of Experimental Biology* 38, 1177–1182.
- Prietom, J.M., Recio, M.C., Giner, R.M., Mániz, S., Ríos, J.L., 2003. Pharmacological approach to the pro- and anti-inflammatory effects of *Ranunculus sceleratus* L. *Journal of Ethnopharmacology* 89, 131–137.
- Priya, C.L., Rao, K.V.B., 2012. Ethnobotanical and current ethnopharmacological aspects of *Argemone mexicana* Linn. *International Journal of Pharmaceutical Sciences and Research* 3, 2143–2148.
- Puapattanakul, O., 1980. Research on the use of *Annona squamosa* seeds and leaves to treat hair lice. *Journal of the Pharmacologists Association, Thailand* 34, 93.
- Punam, J., Bhanu, P., Manoj, G., 2012. Fatty acid composition and physico-chemical properties of the oil extracted from root bark of *Vallisneria spiralis* (Roth) Kuntze. *International Research Journal of Pharmacy* 3, 165–166.
- Punjani, B.L., 2002. Ethnobotanical aspects of some plants of Aravali hills in North Gujarat. *Ancient Science of Life* 21, 268–280.
- Purkayastha, J., Nath, S.C., Islam, M., 2005. Ethnobotany of medicinal plants from Dibru-Saikhowa Biosphere Reserve of North east India. *Fitoterapia* 76, 121–127.
- Purohit, S.K., Solanki, R., Soni, R., Mathur, V., 2013. Evaluation of wound healing activity of ethanolic extract of *Azadirachta indica* leaves in male albino rats. *Asian Journal of Pharmacy and Technology* 3, 73–75.
- Radhika, L.G., Meena, C.V., Peter, S., Rajesh, K.S., Rosamma, M.P., 2011. Phytochemical and antimicrobial study of *Oroxylum indicum*. *Ancient Science of Life* 30, 114–120.
- Radwan, M.M., Elshohly, M.A., Slade, D., Ahmed, S.A., Khan, I.A., Ross, S.A., 2009. Biologically active cannabinoids from high-potency *Cannabis sativa*. *Journal of Natural Products* 72, 906–911.
- Rahate, K.P., Rajasekaran, A., Arulkumar, K.S.G., 2012. Potential of *desmostachya bipinnata* stapf (poaceae) root extracts in inhibition of cell proliferation of cervical cancer cell lines. *International Journal of Research in Pharmaceutical Sciences* 3, 5–11.
- Rahman, M.A., Akter, N., Rashid, H., Ahmed, N.U., Uddin, N., Islam, M.S., 2012. Analgesic and anti-inflammatory effect of whole *Ageratum conyzoides* and *Emilia sonchifolia* alcoholic extracts in animal models. *African Journal of Pharmacy and Pharmacology* 6, 1469–1476.
- Rahman, M.A., Uddin, M.N., Ahmed, N.U., Islam, M.D., 2012. Phytochemical analysis and effect of whole *Eclipta alba* ethanolic extract on the central nociception and inflammation in rodent models. *Journal of Pharmaceutical and Biomedical Sciences* 25, 130–140.
- Rahman, M.A., Bachar, S.C., Rahmatullah, M., 2010. Analgesic and anti-inflammatory activity of methanolic extract of *Acalypha indica* Linn. *Pakistan Journal of Pharmaceutical Sciences* 23, 256–258.
- Rahman, M.S., Salehin, M.F., Jamal, M.A.H.M., Parvin, A., Alam, M.K., 2011. Antibacterial activity of *Argemone mexicana* L. against water borne microbes. *Research Journal of Medicinal Plant* 5, 621–626.
- Rai, S., Aggarwal, K.K., Babu, C.R., 2008. Isolation of a serine Kunitz trypsin inhibitor from leaves of *Terminalia arjuna*. *Current Science* 94, 1509–1512.
- Raja, R.V., Ramanathan, T., Savitha, S., 2009. Studies on wound healing property of coastal medicinal plants. *Journal of Bioscience and Technology* 1, 39–44.
- Raja, V., Ovais, M., Dubey, A., 2013. Phytochemical screening and antibacterial activity of *Lawsonia inermis* leaf extract. *International Journal of Microbiological Research* 4, 33–36.
- Rajakumar, N., Shivanna, M.B., 2009. Ethno-medicinal application of plants in the eastern region of Shimoga district, Karnataka, India. *Journal of Ethnopharmacology* 126, 64–73.
- Rajani, D.P., Ashok, P., 2009. *In vitro* antioxidant and antihyperlipidemic activities of *Bauhinia variegata* Linn. *Indian Journal of Pharmacology* 41, 227–232.
- Rajanna, S., Kolhapure, S.A., 2005. Evaluation of the efficacy and safety of “Muscle & Joint Rub” in the treatment of sprains, contusions and inflammatory musculoskeletal disorders. *The Antiseptic* 102, 389–393.
- Rajasekaran, S., Dinesh, M.G., Kansraj, C., Baig, F.H.A., 2014. *Amaranthus spinosus* leaf extracts and its anti-inflammatory effects on cancer. *Indian Journal of Research in Pharmacy and Biotechnology* 2, 1058–1064.
- Rajasekharan, S., Srivastava, T.N., 1977. Ethnobotanical study on vacha and a preliminary clinical trial on bronchial asthma. *Journal of Research in Indian Medicine, Yoga and Homoeopathy* 12, 92–96.
- Rajkapoor, B., Jayakar, B., Muruges, N., Sakthisekaran, D., 2006. Chemoprevention and cytotoxic effect of *Bauhinia variegata* against N-nitrosodiethylamine induced liver tumors and human cancer cell lines. *Journal of Ethnopharmacology* 104, 407–409.
- Ramadas, S.R., Ghotge, N.S., Ashalata, S., Mathur, N.P., Gour, V., Rao, S., 2000. Ethnoveterinary remedies used in common surgical conditions in some districts of Andhra Pradesh and Maharashtra, India. *Ethnobotany* 12, 100–112.
- Ramakrishna, S., Geetha, K.M., Gopal, P.V.V.S.B., Kumar, P.R., Madhav, P.C., Umachandar, L., 2011. Effect of *Mallotus philippensis* Muell.-Arg leaves against hepatotoxicity of carbon tetrachloride in rats. *International Journal of Pharma Sciences and Research* 2, 74–83.
- Ramakrishnaiah, G., Hariprasad, T., 2012. *In vitro* antimicrobial activity of leaves and bark extracts of *Ficus religiosa* (Linn.). *International Research Journal of Pharmacy* 3, 178–179.
- Ramprasad, V.R., Palanivelu, S., Panchanatham, S., 2005. Anti-inflammatory effect of *Semecarpus anacardium* Linn. Nut extract in acute and chronic inflammatory conditions. *Biological and Pharmaceutical Bulletin* 27, 2028–2031.
- Rani, S., Ranam, J.C., Jeelani, S.M., Gupta, R.C., Kumari, S., 2013. Ethnobotanical notes on 30 medicinal polypetalous plants of district Kangra of Himachal Pradesh. *Journal of Medicinal Plants Research* 7, 1362–1369.
- Rasik, A.M., Raghubir, R., Gupta, A., Shukla, A., Dubey, M.P., Srivastava, S., Jain, H.K., Kulshrestha, D.K., 1999. Healing potential of *Calotropis procera* on dermal wounds in Guinea pigs. *Journal of Ethnopharmacology* 68, 261–266.
- Rathi, A., Rao, Ch.V., Khatoon, S., Mehrotra, S., 2003. Ethnopharmacological evaluation of *Peristrophe bicalyculata* Nees. for anti-inflammatory and analgesic activity. *Natural Product Sciences* 9, 195–199.
- Rathi, A., Rao, Ch.V., Ravishankar, B., De, S., Mehrotra, S., 2004. Anti-inflammatory and anti-nociceptive activity of the water decoction *Desmodium gangeticum*. *Journal of Ethnopharmacology* 95, 259–263.
- Ratnasooriya, W.D., Jayakody, J.R.A.C., Premakumara, G.A.S., Ediriweera, E.R.H.S.S., 2005. Antioxidant activity of water extract of *Scoparia dulcis*. *Fitoterapia* 76, 220–222.
- Ravichandran, G., Bharadwaj, V.S., Kolhapure, S.A., 2004a. Evaluation of efficacy and safety of Acne-N-Pimple cream in acne vulgaris. *The Antiseptic* 101, 249.
- Ravichandran, G., Bharadwaj, V.S., Kolhapure, S.A., 2004b. Evaluation of the clinical efficacy of “Anti-Dandruff Shampoo” in the treatment of dandruff. *The Antiseptic* 101, 5–8.
- Ravichandran, G., Bharadwaj, V.S., Kolhapure, S.A., 2005. Evaluation of the efficacy and safety of “Anti-Wrinkle cream” in the treatment of facial skin wrinkles: a prospective, open, phase III clinical trial. *The Antiseptic* 102, 65–70.
- Ravichandran, G., Singh, A., Patki, P.S., Mitra, S.K., 2008. Clinical evaluation of herbal Hair Loss Cream in management of *Alopecia aerata*: an open study. *Medicine Update* 15, 31–34.
- Ray, S., Sheikh, M., Mishra, S., 2011. Ethnomedicinal plants used by tribals of East Nimar region, Madhya Pradesh. *Indian Journal of Traditional Knowledge* 10, 367–371.
- Reddy, C.S., Reddy, K.N., Rao, K.T., Pattananaik, C., 2007. Ethnomedicinal studies on medicinal plants used by the chenchus of Nallamalais in Kurnool district, Andhra Pradesh, India. *Research Journal of Medicinal Plant* 1, 128–133.
- Reddy, C.V., Kamble, A., 2014. Toxicity study of *Achyranthus aspera*. *International Letters of Natural Sciences* 9, 85–96.
- Reddy, J.S., Rao, P.R., Reddy, M.S., 2002. Wound healing effects of *Heliotropium indicum*, *Plumbago zeylanicum* and *Acalypha indica* in rats. *Journal of Ethnopharmacology* 79, 249–251.
- Reddy, K.N., Trimurthulu, G., Reddy, Ch.S., 2010. Medicinal plants used by ethnic people of Medak district, Andhra Pradesh. *Indian Journal of Traditional Knowledge* 9, 184–190.
- Reddy, S.B., Reddy, K.K.R., Naidu, V.G., Madhusudhana, K., Agwane, S.B., Ramakrishna, S., Diwan, P.V., 2008. Evaluation of antimicrobial, antioxidant and wound-healing potentials of *Holoptelea integrifolia*. *Journal of Ethnopharmacology* 115, 249–256.
- Resmi, C.R., Venukumar, M.R., Latha, M.S., 2006. Antioxidant activity of *Albizia lebbek* (Linn.) Benth. in alloxan diabetic rats. *Indian Journal of Physiology and Pharmacology* 50, 297–302.
- Revathi, P., Parimalazhagan, T., 2010. Traditional Knowledge on Medicinal Plants Used by the Irula Tribe of Hasanur Hills, Erode District, Tamil Nadu, India. *Ethnobotanical Leaflets* 14, 136–160.
- Riel, M.A., Kyle, D.E., Milhous, W.K., 2002. Efficacy of scopadulcic acid A against *Plasmodium falciparum* in vitro. *Journal of Natural Products* 65, 614–615.
- Riyaz, S., Manisha, R.L., Pal, R., Shamshuddin, M., Putta, R.K., 2012. Anti-inflammatory evaluation of alcoholic and aqueous extracts of *Allium cepa* Linn. (Liliaceae) in rats. *Novel Science International Journal of Medical Sciences* 1, 186–190.
- Ramanjaneyulu, K., Bhargavi, A., Buddaraj, P.R.V., Pavani, S.K., Mehavineela, P., 2011. Evaluation of phytochemical and antibacterial activity of *Butea monosperma* leaf extracts. *International Journal of Research in Ayurveda and Pharmacy* 2, 1563–1565.
- Roy, K., Shivakumar, H., Sarkar, S., 2009. Wound healing potential of leaf extracts of *Ficus religiosa* on wistar albino strain rats. *International Journal of PharmTech Research* 1, 506–508.

- Roy, M.K., Nakahara, K., Na, T.V., Trakoontivakorn, G., Takenaka, M., Isobe, S., Tsuchida, T., 2007. Baicalein, a flavonoid extracted from a methanolic extract of *Oroxylum indicum* inhibits proliferation of a cancer cell line *in vitro* via induction of apoptosis. *Pharmazie* 62, 149–153.
- Roy, N., Laskar, R.A., Sk, I., Kumari, D., Ghosh, T., Begum, N.A., 2011. A detailed study on the antioxidant activity of the stem bark of *Dalbergia sissoo* Roxb., an Indian medicinal plant. *Food Chemistry* 126, 1115–1121.
- Rumzhum, N.N., Rahman, M.M., Islam, M.S., Chowdhury, S.A., Sultana, R., Parvin, M. N., 2008. Cytotoxicity and antioxidant activity of extracts from *Mirabilis jalapa*. *Stamford Journal of Pharmaceutical Sciences* 1, 85–88.
- Sadashiv, P.S., Krishna, A.R., 2011. Acute toxicity study for *Achyranthes aspera* leaves. *Journal of Pharmacy Research* 4, 2221.
- Saha, A., Ahmed, M., 2009. The analgesic and anti-inflammatory activities of the extract of *Albizia lebbek* in animal model. *Pakistan Journal of Pharmaceutical Sciences* 22, 74–77.
- Saha, S., Subrahmanyam, E.V.S., 2013. Evaluation of anti-inflammatory activity of ethanolic extract of seeds of *Holarrhena pubescens* (Buch.-Ham.) Wall. *International Journal of Pharmacy & Pharmaceutical Sciences* 5, 915–919.
- Sahoo, A.K., Narayanan, N., Sahana, S., Rajan, S.S., Mukherjee, P.K., 2008. *In vitro* antioxidant potential of *Semecarpus anacardium* L. *Pharmacology Online* 3, 327–335.
- Sahu, G., Gupta, P.K., 2012. A review on *Bahunia variegata* Linn. *International Research Journal of Pharmacy* 3, 48–51.
- Sahu, M.C., Debata, N.K., Padhy, R.N., 2012. Antibacterial activity of *Argemone mexicana* L. against multidrug resistant *Pseudomonas aeruginosa*, isolated from clinical samples. *Asian Pacific Journal of Tropical Biomedicine*, S800–S807.
- Saikia, A.P., Ryakala, V.K., Sharma, P., Goswami, P., Bora, U., 2006. Ethnobotany of medicinal plants used by Assamese people for various skin ailments and cosmetics. *Journal of Ethnopharmacology* 106, 149–157.
- Sakarkar, D.M., Sakarkar, U.M., Shrikhande, V.N., Vyas, J.V., Mandavgade, S., Jaiswal, S.B., Purohit, R.N., 2004. Wound healing properties of Henna leaves. *Natural Product Radiance* 3, 406–412.
- Sakulpanich, A., Chewchinda, S., Sithisarn, P., Gritsanapan, W., 2012. Standardization and toxicity evaluation of *Cassia fistula* pod pulp extract for alternative source of herbal laxative drug. *Pharmacognosy Journal* 4, 6–12.
- Saleem, T.S.M., Basha, S.D., Mahesh, G., Rani, P.V.S., Kumar, N.S., Chetty, C.M., 2011. Analgesic, anti-pyretic and anti-inflammatory activity of dietary sesame oil in experimental animal models. *Pharmacologia* 2, 172–177.
- Sambrekar, S.N., Patil, P.A., Patil, S.A., 2011. Wound healing activity of root extracts of *Commelina benghalensis* Linn. *Research Journal of Pharmacy and Technology* 4, 1772–1776.
- Sandhiya, S., Subhasree, N., Shivapriya, S., Agrawal, A., Dubey, G.P., 2013. *In vitro* antioxidant and antimicrobial potential of *Bambusa arundinacea* (Retz.) Willd. *International Journal of Pharmacy and Pharmaceutical Sciences* 5, 359–362.
- Sannigrahi, S., Mazumder, U.K., Pal, D., Mishra, S.L., 2012. Terpenoids of methanol extract of *Clerodendrum infortunatum* exhibit anticancer activity against Ehrlich's ascites carcinoma (EAC) in mice. *Pharmaceutical Biology* 50, 304–309.
- Sapkale, A.P., Thorat, M.S., Dighe, D.A., Munot, N.M., Singh, M.C., 2013. Investigation of anti-inflammatory activity of whole flower extract of *Butea monosperma* by ex-vivo and in-vitro techniques and their correlation. *International Journal of Pharmacy & Pharmaceutical Sciences* 5, 84–86.
- Saravanan, R., Dhachinamoorthi, D., Senthilkumar, K., Sriakshmi, M., DivyaSri, T., 2012. Antibacterial activity of *Euphorbia hirta* extracts. *International Journal of Research in Ayurveda and Pharmacy* 3, 439–441.
- Sarg, T., Ateya, A., Abdel-Ghani, A., Badr, W., Shams, G., 1999. Phytochemical and pharmacological studies of *Dalbergia sissoo* growing in Egypt. *Pharmaceutical Biology* 37, 54–62.
- Sarrell, E.M., Cohen, H.A., Kahan, E., 2003. Naturopathic treatment for ear pain in children. *Pediatrics* 111, e574–579.
- Sathish, R., Ahmed, M.H., Natarajan, K., Lalitha, K.G., 2010. Evaluation of wound healing and antimicrobial activity of *Lannea coromandelica* (Houtt) Merr. *Journal of Pharmacy Research* 3, 1225–1228.
- Sathya, M., Kokilavani, R., Ananta, T.K.S., 2012. Acute and subacute toxicity studies of ethanolic extract of *Acalypha indica* Linn in male wistar albino rats. *Asian Journal of Pharmaceutical and Clinical Research* 5, 97–100.
- Satish, A., Kumar, R.P., Rakshith, D., Satish, S., Ahmed, F., 2013. Antimutagenic and antioxidant activity of *Ficus benghalensis* stem bark and *Moringa oleifera* root extract. *International Journal of Chemical and Analytical Science* 4, 45–48.
- Sawant, M., Isaac, J.C., Narayanan, S., 2004. Analgesic studies on total alkaloids and alcohol extracts of *Eclipta alba* (Linn.) Hassk. *Phytotherapy Research* 18, 111–113.
- Schrader, F.C., Barho, M., Steiner, I., Ortmann, R., Schlitzer, M., 2012. The antimalarial pipeline – an update. *International Journal of Medical Microbiology* 302, 165–171.
- Schwartz, L., Tulipan, L., Birmingham, D.J., 1957. *Occupational Diseases of the Skin*, 3rd edn. Henry Kimpton, London.
- Seetharam, Y.N., Chalgeri, G., Haleshi, C., Vijay, 1999. Folk medicine and ethnomedicine of North-Eastern Karnataka. *Ethnobotany* 11, 32–37.
- Sen, S.K., Behara, L.M., 2003. Ethnomedicinal plants used against skin diseases in Bargar district of Orissa. *Ethnobotany* 15, 90–96.
- Senthamilselvi, M.M., Kesavan, D., Sulochana, N., 2012. An anti-inflammatory and anti-microbial flavone glycoside from flowers of *Cleome viscosa*. *Organic and Medicinal Chemistry Letters* 2, 19. <http://dx.doi.org/10.1186/2191-2858-2-19>.
- Senthilkumar, M., Urumoorthi, P.G., Janardhanan, K., 2006a. Some medicinal plants used by the Irular, the tribal people of Marudhamalai hills, Coimbatore, Tamilnadu. *Natural Product Radiance* 5, 382–388.
- Senthilkumar, M., Sripriya, R., Vijaya Raghavan, H., Sehgal, P.K., 2006b. Wound healing potential of *Cassia fistula* on infected albino rat model. *Journal of Surgical Research* 131, 283–289.
- Senthilkumar, S., Akram, A.S., Fareed Ahmed, T.S., Mohamed Jaabir, M.S., 2010. Phytochemical analysis and antimicrobial activity of the ethanolic extract of *Acorus calamus* rhizome. *Orient Journal of Chemistry* 26, 223–227.
- Sermakkani, M., Thangapandian, V., 2010. Phytochemical and antibacterial activity of *Martynia annua* L. against the different pathogenic bacteria. *Journal of Herbal Medicine and Toxicology* 4, 221–224.
- Shagal, M.H., Modibbo, U.U., Liman, A.B., 2012. Pharmacological justification for the ethnomedicinal use of *Datura stramonium* stem-bark extract in treatment of diseases caused by some pathogenic bacteria. *International Research Journal of Pharmacy and Pharmacology* 2, 16–19.
- Shah, P.D., Ghag, M., Deshmukh, P.B., Kulkarni, Y., Joshi, S.V., Vyas, B.A., Shah, D.R., 2012. Toxicity study of ethanolic extract of *Acorus calamus* rhizome. *International Journal of Green Pharmacy* 6, 29–35.
- Shah, N.T., Pandya, T.N., Sharma, P.P., Patel, B.R., Acharya, R., 2012. Mootral Karma of Kusha [*Imperata cylindrical* Beauv.] and Darbha [*Desmostachya bipinnata* Stapf.] – a comparative study. *AYU* 33, 387–390.
- Shahabuddin, K.U., Sarwar, M.S., Mohiuddin, E., 2006. Clinical evaluation of some herbal medicine for amoebiasis. *Pakistan Journal of Pharmacology* 23, 9–12.
- Shahavi, V.M., Desai, S.K., 2008. Antiinflammatory activity of *Butea monosperma*. *Fitoterapia* 79, 82–85.
- Shankar, A., Parsai, M.R., Naqvi, S.M.A., Jain, J.P., 1980. A clinical trial of apamarga (*Achyranthes aspera*) in cases of shoth (general anasarca). *Journal of Research in Ayurveda and Siddha* 1, 514–528.
- Sharma, J., Gairola, S., Gaur, R.D., Painuli, R.M., 2011. Medicinal plants used for primary healthcare by Tharu tribe of Udham Singh Nagar, Uttarakhand, India. *International Journal of Medicinal and Aromatic Plants* 1, 228–233.
- Sharma, R.K., Rajani, G.P., Sharma, V., Komala, N., 2011. Effect of ethanolic and aqueous extracts of *Bauhinia variegata* Linn. On gentamicin-induced nephrotoxicity in rats. *International Journal of Pharmaceutical Sciences* 45, 192–198.
- Sharma, J., Gairola, S., Gaur, R.D., Painuli, R.M., 2012. The treatment of jaundice with medicinal plants in indigenous communities of the Sub-Himalayan region of Uttarakhand, India. *Journal of Ethnopharmacology* 143, 262–291.
- Sharma, K.K., Kotoky, J., Kalita, J.C., Barthakur, R., 2012b. Evaluation of antidermatophytic activity of *Ranunculus sceleratus* and *Pongamia pinnata* available in North Eastern Region of India. *Asian Pacific Journal of Tropical Biomedicine*, S806–S811.
- Sharma, P., Nath, A.K., Kumari, R., Bhardwaj, S.V., 2012. Purification, characterization and evaluation of insecticidal activity of trypsin inhibitor from *Albizia lebbek* seeds. *Journal of Forestry Research* 23, 131–137.
- Sharma, K.K., Kotoky, J., Kalita, J.C., Barthakur, R., 2012a. Evaluation of antidermatophytic activity of *Ranunculus sceleratus* and *Pongamia pinnata* available in North Eastern Region of India. *Asian Pacific Journal of Tropical Biomedicine*, S808–S811.
- Sharma, J., Gaur, R.D., Painuli, R.M., 2013a. Traditional herbal medicines used for the treatment of skin disorders by the Gujjar tribe of Sub-Himalayan tract, Uttarakhand. *Indian Journal of Traditional Knowledge* 12, 736–746.
- Sharma, J., Gairola, S., Gaur, R.D., Painuli, R.M., Siddiqi, T.O., 2013b. Ethnomedicinal plants used for treating epilepsy by indigenous communities of sub-Himalayan region of Uttarakhand, India. *Journal of Ethnopharmacology* 150, 353–370.
- Sharma, A.K., Sharma, A.K., Sharma, R.B., 2013. Survey of medicinal plants of Hatras district used in Skin diseases. *Current Research in Medicine and Medical Sciences* 3, 16–18.
- Sharma, J., Painuli, R.M., 2011. Plants used for the treatment of Rheumatism by the Bhoja tribe of District Dehradun, Uttarakhand. *International Journal of Medicinal and Aromatic Plants* 1, 28–32.
- Sharma, J., Painuli, R.M., Gaur, R.D., 2010. Plants used by the rural communities of district Shahjahanpur, Uttar Pradesh. *Indian Journal of Traditional Knowledge* 9, 798–803.
- Sharma, K., Shukla, S.D., Mehta, P., Bhatnagar, M., 2002. Fungistatic activity of *Semecarpus anacardium* Linn. F. Nut extract. *Indian Journal of Experimental Biology* 40, 314–318.
- Sharma, M.C., Sharma, S., 2010. Phytochemical screening of methanolic extract and antibacterial activity of *Eclipta alba* and *Morina citrifolia* L. *Middle-East Journal of Scientific Research* 6, 445–449.
- Sharma, P.K., Chauhan, N.S., Lal, B., 2004. Observations on the traditional phytotherapy among the inhabitants of Parvati valley in western Himalaya, India. *Journal of Ethnopharmacology* 92, 167–176.
- Sharma, R.K., 2010. Pharmacological Evaluation of *Bauhinia Variegata* Linn. for Wound Healing and Nephroprotective Activity (M.Sc. thesis). Rajiv Gandhi University of Health Sciences, Karnataka.
- Sharma, S.K., Gupta, V.K., 2007. *In vitro* antioxidant study of *Ficus religiosa* Linn. root. *International Journal of Chemical Sciences* 5, 2365–2371.
- Sheela, Q.R.X., Arockiasamy, P., Singh, J.D., Ilavarasan, R., Ramani, V.A., 2011. Anti-ulcer evaluation of *Polygonum barbatum* linn leaf extract. *International Journal of Pharmacy and Pharmaceutical Sciences* 3, 81–84.
- Shelly, W.B., Arthur, R.P., 1955. Mucinain, the active pruritogenic proteinase of cowhage. *Science* 122, 469–470.
- Shendkar, C.D., Chandrachud, P.S., Lavate, S.M., Jagtap, S.D., Deshpande, N.R., 2012. Antibacterial screening of *Achyranthes aspera* Linn. root extracts. *Annals of Biological Research* 3, 2202–2204.
- Shenoy, C., Patil, M.B., Kumar, R., Patil, S., 2009a. Preliminary phytochemical investigation and wound healing activity of *Allium cepa* Linn. (Liliaceae). *International Journal of Pharmacy and Pharmaceutical Sciences* 2, 167–179.

- Shenoy, C., Patil, M.B., Kumar, R., 2009b. Wound healing activity of *Hyptis suaveolens* (L.) Poit (Lamiaceae). *International Journal of PharmTech Research* 1, 737–744.
- Shiddamallayya, N., Yasmeen, A., Gopakumar, K., 2010. Hundred common forest medicinal plants of Karnataka in primary healthcare. *Indian Journal of Traditional Knowledge* 9, 90–95.
- Shim, Y.Y., Gui, B., Arnison, P.G., Wang, Y., Reaney, M.J.T., 2014. Flaxseed (*Linum usitatissimum* L.) bioactive compounds and peptide nomenclature: A review. *Trends in Food Science & Technology* 38, 5–20.
- Shirwaikar, A., Rajendran, K., Kumar, C.D., 2004. *In vitro* antioxidant studies of *Annona squamosa* Linn. leaves. *Indian Journal of Experimental Biology* 42, 803–807.
- Shivanna, M.B., Rajakumar, N., 2010. Ethno-medico-botanical knowledge of rural folk in Bhadravathi taluk of Shimoga district, Karnataka. *Indian Journal of Traditional Knowledge* 9, 158–162.
- Shreedhara, C.S., Vaidya, V.P., 2006. Screening of *Momordica dioica* for hepatoprotective, antioxidant, and anti-inflammatory activities. *Natural Product Sciences* 12, 157–161.
- Shukla, R., Singh, S., Bhandari, C.R., 1973. Preliminary clinical trials on antidiabetic action of *Azadirachta indica*. *Medicine and Surgery* 13, 11–12.
- Siddhuraju, P., Mohan, P.S., Becker, K., 2002. Studies on the antioxidant activity of Indian Laburnum (*Cassia fistula* L.): a preliminary assessment of crude extracts from stem bark, leaves, flowers and fruit pulp. *Food Chemistry* 79, 61–67.
- Siddiqui, B.S., Ali, S.T., Rizwani, G.H., Begum, S., Tauseef, S., Ahmad, A., 2012. Antimicrobial activity of the methanolic bark extract of *Holarrhena pubescens* (Buch. Ham), its fractions and the pure compound conessine. *Natural Product Research* 26, 987–992.
- Silja, V.P., Varma, S.K., Mohanan, K.V., 2008. Ethnomedicinal plant knowledge of the Mullu kuruma tribe of Wayanad district, Kerala. *Indian Journal of Traditional Knowledge* 7, 604–612.
- Sindhoor, K.L., Gurram, S.K., Nagarjuna, S., Reddy, Y.P., 2012. Comparative study of anti-inflammatory activity of petroleum ether and ethanolic extracts of *Brassica juncea*. *International Journal of PharmTech Research* 4, 1172–1176.
- Singh, A., Singh, P.K., 2009. An ethnobotanical study of medicinal plants in Chandauli District of Uttar Pradesh, India. *Journal of Ethnopharmacology* 121, 324–329.
- Singh, A.K., Raghubanshi, A.S., Singh, J.S., 2002. Medical ethnobotany of the tribals of Sonaghati of Sonbhadra district, Uttar Pradesh, India. *Journal of Ethnopharmacology* 81, 31–41.
- Singh, K.K., Maheswari, J.K., 1994. Traditional phytotherapy of some Medicinal plants used by the Tharus of the Nainital district, Uttar Pradesh, India. *Pharmaceutical Biology* 32, 51–58.
- Singh, B.R., Singh, B.R., Singh, R.L., Prakash, D., Singh, D.P., Sarma, B.K., Upadhyay, G., Singh, H.B., 2009. Polyphenolics from various extracts/fractions of red onion (*Allium cepa*) peel with potent antioxidant and antimutagenic activities. *Food and Chemical Toxicology* 47, 1161–1167.
- Singh, P.P., Ambika, Chauhan, S.M.S., 2009. Activity guided isolation of antioxidants from the leaves of *Ricinus communis* L. *Food Chemistry* 114, 1069–1072.
- Singh, M., Kumar, V., Singh, I., Gauttam, V., Kalia, A.N., 2010. Anti-inflammatory activity of aqueous extract of *Mirabilis jalapa* Linn. leaves. *Pharmacognosy Research* 2, 364–367.
- Singh, K.P., Daboriya, V., Kumar, S., Singh, S., 2010. Antibacterial activity and phytochemical investigations on *Nicotiana glauca* (wild tobacco). *Romanian Journal of Biology – Plant Biology* 55, 135–142.
- Singh, R., Seherawat, A., Sharma, P., 2011. Hypoglycemic, antidiabetic and toxicological evaluation of *Momordica dioica* fruit extracts in Alloxan induced diabetic rats. *Journal of Pharmacology and Toxicology* 6, 454–467.
- Singh, H., Singh, V., Kumar, D., Chaudhary, A.K., 2011. Wound healing and antimicrobial potential of *Oroxylum indicum* Vent. in albino mice. *The Natural Products Journal* 1, 128–134.
- Singh, M., Kale, R.K., 2010. Abstract 64: Chemopreventive potential of *Bauhinia variegata* against DMBA-induced papillomagenesis in Swiss albino mice. In: Abstracts: AACR International Conference on Frontiers in Cancer Prevention Research. December 6–9, 2009, Houston, TX; Cancer Prevention Research 3, A64.
- Singh, R., Gupta, A.K., 2008. Antimicrobial and antitumor activity of the fractionated extracts of *Curculigo orchioides*. *International Journal of Green Pharmacy* 2, 34–36.
- Singh, R., Siddiqui, M.A., Baurah, M.C., 1978. Plant dermatitis in Delhi. *Indian Journal of Medical Research* 68, 650–655.
- Singh, S., Singh, G.B., 2006. Anti-inflammatory activity of *Lannea coromandelica* bark extract in rats. *Phytotherapy Research* 8, 311–313.
- Singh, S., Bothara, S.B., 2012. Acute toxicity studies of natural materials extracted from indigenous edible fruits available in Chhattisgarh. *Inventi Rapid: Planta Activa* 2012, 1–3.
- Sinha, M.K., Patel, D.K., Kanungo, V.K., 2013. Medicinal plants used in the treatment of skin diseases in Central Bastar of Chhattisgarh, India. *Global Advanced Research Journal of Medicinal Plants* 2, 1–3.
- Sinha, S.N., 2012. Antibacterial potential of crude methanolic extract of *Leonotis nepetifolia* (L.) R. B.R. *International Research Journal of Pharmacy* 3, 277–278.
- Sireeratawong, S., Lertprasertsuke, N., Srisawat, U., Thupia, A., Ngamjariyawat, A., Suwanlikhid, N., Jaijoy, K., 2008. Acute and subchronic toxicity study of the water extract from root of *Sida rhombifolia* Linn. in rats. *Songklanakarin Journal of Science and Technology* 30, 729–737.
- Sivakumar, B., Velmurugan, C., Kumar, P.R.L., 2013. Diuretic activity of methanolic extract of '*Albizia lebbek*'. *International Journal of PharmTech Research* 5, 404–406.
- Sivaranjani, R., Ramakrishnan, K., 2012. Traditional uses of medicinal plants in treating skin diseases in Nagapattinam district of Tamilnadu, India. *International Research Journal of Pharmacy* 3, 201–204.
- Slaviero, G., 1916. Disease among workers in hemp cultivation. *The Lancet* 187, 580.
- Somchit, M.N., Rashid, R.A., Abdullah, A., Zuraini, A., Zakaria, Z.A., Sulaiman, M.R., Arifah, A.K., Mutalib, A.R., 2010. *In vitro* antimicrobial activity of leaves of *Acalypha indica* Linn. (Euphorbiaceae). *African Journal of Microbiological Research* 4, 2133–2136.
- Souder, P., 1963. Poisonous plants of Guam. In: Keegan, H.L., Macfarlane, W.V. (Eds.), *Venomous and Poisonous Animals and Noxious Plants of the Pacific Region*. Pergamon Press, New York, pp. 15–29.
- Sourabie, T.S., Ouedraogo, N., Sawadogo, W.R., Nikema, J.B., Guissou, I.P., Nacoulma, O.G., 2012. Biological evaluation of anti-inflammatory and analgesic activities of *Argemone mexicana* Linn. (Papaveraceae) aqueous leaf extract. *International Journal of Pharma Sciences and Research* 3, 451–458.
- Speranza, L., Franceschelli, S., Pesce, M., Menghini, L., Patrino, A., Vinciguerra, I., De Lutiis, M.A., Felaco, M., Felaco, P., Grilli, L., 2009. Anti-inflammatory properties of the plant *Verbascum mallophorum*. *Journal of Biological Regulators & Homeostatic Agents* 23, 189–195.
- Spiewak, R., 2000. Occupational skin diseases among farmers. In: Zagorski, J. (Ed.), *Occupational and ParaOccupational Diseases in Agriculture*. Institute of Agriculture Medicine, Lublin, Poland, pp. 142–152.
- Sreelekshmi, R., Latha, P.G., Arafat, M.M., Shyamal, S., Shine, V.J., Anuja, G.I., Suja, S. R., Rajasekharan, S., 2007. Anti-inflammatory, analgesic and anti-lipid peroxidation studies on stem bark of *Ficus religiosa* Linn. *Natural Product Radiance* 6, 377–381.
- Srinivasan, K., Muruganandan, S., Lal, J., Chandra, S., Tandan, S.K., Prakash, V.R., 2001. Evaluation of anti-inflammatory activity of *Pongamia pinnata* leaves in rats. *Journal of Ethnopharmacology* 78, 151–157.
- Subramanian, P.M., Misra, G.S., 1978. Chemical constituents of *Ficus bengalensis* (part II). *Polish Journal of Pharmacology and Pharmacy* 30, 559–562.
- Subramanian, R., Krishnaswamy, G., Devaraj, A., Sethuraman, P., Jayakumararaj, R., 2011. Wound healing ethnopharmacological potentials of selected medicinal plants used by Malayali Tribes. *International Research Journal of Pharmacy* 2, 132–137.
- Sultana, B., Anwar, F., Przybylski, R., 2007. Antioxidant activity of phenolic components present in barks of *Azadirachta indica*, *Terminalia arjuna*, *Acacia nilotica*, and *Eugenia jambolana* Lam. trees. *Food Chemistry* 104, 1106–1114.
- Sumitra, M., Manikandan, P., Suguna, L., 2005. Efficacy of *Butea monosperma* on dermal wound healing in rats. *The International Journal of Biochemistry & Cell Biology* 37, 566–573.
- Sundriyal, A., Bijem, K.R.V., Kalia, A.N., 2013. Antiepileptic potential of *Anisomeles indica* (Linn.) Kuntze aerial parts in pentylenetetrazole-induced experimental convulsions in Wistar rats. *Indian Journal of Experimental Biology* 51, 715–720.
- Suresh, M., Ayyanar, M., Amalraj, L., Mehalingam, P., 2012. Ethnomedicinal Plants used to treat Skin Diseases in Pothigai hills of Western Ghats, Tirunelveli District, Tamil Nadu, India. *Journal of Biosciences Research* 3, 112–121.
- Susmitha, S., Vidyamol, K.K., Ranganayaki, P., Vijayaragavan, R., 2013. Phytochemical extraction and antimicrobial properties of *Azadirachta indica* (Neem). *Global Journal of Pharmacology* 7, 316–320.
- Sutar, R.C., Kasture, S.B., Kalaichelvan, V.K., 2014. Evaluation of anticonvulsant activity of leaf extracts of *Holoptelea integrifolia* (roxb.) Planch in experimental animals. *International Journal of Pharmacy & Pharmaceutical Sciences* 6, 308–311.
- Tahiliani, P., Kar, A., 2000. *Achyranthes aspera* elevates thyroid hormone levels and decreases hepatic lipid peroxidation in male rats. *Journal of Ethnopharmacology* 71, 527–532.
- Takada, Y., Aggarwal, B.B., 2003. Betulinic acid suppresses carcinogen-induced NF- κ B activation through inhibition of I κ B kinase and p65 phosphorylation: abrogation of cyclooxygenase-2 and matrix metalloproteinase-9. *The Journal of Immunology* 171, 3278–3286.
- Talekar, Y.P., Das, B., Paul, T., Talekar, D.Y., Apte, K.G., Parab, P.B., 2012. Wound healing potential of *Vitex negundo* Linn in experimental animals. *International Journal of Pharmacy and Pharmaceutical Sciences* 4, 543–546.
- Tamboli, A.M., Karpe, S.T., Shaikh, S.A., Manikrao, A.M., Kature, D.V., 2011. Hypoglycemic activity of extracts of *Oroxylum indicum* (L.) Vent roots in animal models. *Pharmacologyonline* 2, 890–899.
- Taylor, J.M., Jones, W.I., Hagan, E.C., Gross, M.A., Davis, D.A., Cook, E.K., 1967. Toxicity of oil of calamus (Jammu variety). *Toxicology and Applied Pharmacology* 10, 405–412.
- Telang, R.S., Chatterjee, S., Varshneya, C., 1999. Studies on analgesic and anti-inflammatory activities of *Vitex negundo* Linn. *Indian Journal of Pharmacology* 31, 363–366.
- Thoh, M., Kumar, P., Nagarajaram, H.A., Manna, S.K., 2010. Azadirachtin interacts with the tumor necrosis factor (TNF) binding domain of its receptors and inhibits TNF-induced biological responses. *The Journal of Biological Chemistry* 285, 5888–5895.
- Thompson, L.U., Seidl, M.M., Rickard, S.E., Orcheson, L.J., Fong, H.H., 1996. Antitumor effect of a mammalian lignan precursor from flaxseed. *Nutrition and Cancer* 26, 159–165.
- Tiangda, C.H., Gritsanapan, W., Sookvanichsilp, N., Limchalcarn, A., 2000. Anti-head lice activity of a preparation of *Annona squamosa* seed extract. *The Southeast Asian Journal of Tropical Medicine and Public Health* 31, 174–177.

- Tiwari, S.K., Lahkar, M., Dash, S., Samudrala, P.K., Thomas, J.M., Augustine, B.B., 2013. Preliminary phytochemical, toxicity and anti-inflammatory evaluation of *Commelina benghalensis*. International Journal of Green Pharmacy 7, 201–205.
- Tomar, A., 2009. Folk medicinal uses of plant roots from Meerut district, Uttar Pradesh. Indian Journal of Traditional Knowledge 8, 298–301.
- Torane, R.C., Kamble, G.S., Khatiwora, E., Ghayal, N.A., Deshpande, N.R., 2011. Antioxidant capacity of leaves and stem of *Ehretia laevis*. International Journal of Pharmacy and Pharmaceutical Sciences 3, 149–151.
- TPL, 2013. The Plant List, A Working List of All Known Plant Species. Version 1.1, released in September 2013. (www.theplantlist.org) (accessed in January 2014).
- Tripathi, A.K., Singh, R.H., 1995. Clinical study on an indigenous drug vaca (*Acorus calamus*) in the treatment of depressive illness. Journal of Research in Ayurveda and Siddha 16, 24–34.
- Tripathi, R., Kaushik, D., Tripathi, A., Rasal, V.P., Khan, S.A., 2006. Acute and subacute toxicity studies on vetiver oil in rats. FABAD Journal of Pharmaceutical Sciences 31, 71–77.
- Tripathi, V.R., Kumar, S., Garg, S.K., 2011. A study on trypsin, *Aspergillus flavus* and *Bacillus* sp. protease inhibitory activity in *Cassia tora* (L.) syn *Senna tora* (L.) Roxb. seed extract. BMC Complementary and Alternative Medicine 11, 56.
- Tripathi, Y.B., Upadhyay, A.K., 2001. Antioxidant property of *Mucuna pruriens* Linn. Current Science 80, 1377–1378.
- Tsai, J.C., Peng, W.H., Chiu, T.H., Lai, S.C., Lee, C.Y., 2011. Anti-inflammatory effects of *Scoparia dulcis* L. and betulinic acid. The American Journal of Chinese Medicine 39, 943–956.
- Turker, A.U., Camper, N.D., 2002. Biological activity of common mullein, a medicinal plant. Journal of Ethnopharmacology 82, 117–125.
- Udupa, A.L., Kulkarni, D.R., Udupa, S.L., 1995. Effect of *Tridax procumbens* extracts on wound healing. Pharmaceutical Biology 33, 37–40.
- Umamaheswari, A., Nuni, A., Shreevidya, R., 2010. Evaluation of antibacterial activity of *Boerhaavia diffusa* L. leaves. International Journal of Green Pharmacy 4, 75–78.
- Uniyal, B., Shiva, V., 2005. Traditional knowledge on medicinal plants among rural women of the Garhwal Himalaya, Uttarakhand. Indian Journal of Traditional Knowledge 4, 259–266.
- Upadhyay, B., Parveen, Dhaker, A.K., Kumar, A., 2010. Ethnomedicinal and ethnopharmaco-statistical studies of Eastern Rajasthan, India. Journal of Ethnopharmacology 129, 64–86.
- Upadhyay, R.K., Balekar, N., Jain, D.K., Jain, N.K., 2013. Evaluation of wound healing activity of stem bark of *Albizia lebbek*. International Journal of Pharmaceutical & Biological Archive 4, 672–677.
- Upadhyaya, O.P., Kumar, K., Tiwari, R.K., 1998. Skin treatments of Bihar plants. Pharmaceutical Biology 36, 20–24.
- Urooj, A., Ahmed, F., 2013. *Ficus racemosa* and *Morus indica*: Emerging alternative antihyperglycemic agents. Open Conference Proceedings Journal 4, 59–65.
- Vagdevi, H.M., Hanumanthappa, B.C., Vaidya, V.P., Raghavendra, R., Parashurama, T. R., 2011. *In vitro* antibacterial and antifungal evaluation of *Vallisneria spiralis* (roth) kuntze. International Journal of Pharmaceutical Research and Development 3, 29–33.
- Vanderjagt, T.J., Ghattas, R., Vanderjagt, D.J., Crossey, M., Glew, R.H., 2002. Comparison of the total antioxidant content of 30 widely used medicinal plants of New Mexico. Life Sciences 70, 1035–1040.
- Varde, A.B., Ainapure, S.S., Naik, S.R., Amladi, S.R., 1988. Anti-inflammatory activity of coconut oil extract of *Acorus calamus*, *Ocimum sanctum* and *Ocimum basilicum* in rats. Indian Drugs 25, 226–228.
- Varma, J., Tripathi, M., Ram, V.J., Pandey, V.B., Dubey, N.K., 2002. γ -Asarone – the fungitoxic principle of the essential oil of *Caesulia axillaris*. World Journal of Microbiology and Biotechnology 18, 277–279.
- Vasantharaj, S., Sathiyavimal, S., Hemashenpagam, N., 2013. Phytochemical analysis and antibacterial activity of *Azadirachta indica*. International Journal of Pharmaceutical Sciences Review and Research 22, 59–61.
- Veerabadrar, U., Venkatraman, A., Souprayan, A., Narayanasamy, M., Perumal, D., Elumalai, S., Sivalingam, S., Devaraj, V., Perumal, A., 2013. Evaluation of antioxidant potential of leaves of *Leonotis nepetifolia* and its inhibitory effect on MCF7 and Hep2 cancer cell lines. Asian Pacific Journal of Tropical Disease 3, 103–110.
- Venkata, S.P., Murali, M.Ch., Kandra, P., Sravani, R., Akondi, R.B., 2012. Screening of antimicrobial and antioxidant potentials of *Acacia caesia*, *Dillenia pentagyna* and *Buchanania lanzan* from Mareduilli Forest of India. Journal of Pharmacy Research 5, 1734–1738.
- Venkatachalam, U., Muthukrishnan, S., 2012. Free radical scavenging activity of ethanolic extract of *Desmodium gangeticum*. Journal of Acute Medicine 2, 36–42.
- Venkataswamy, R., Mubarak, H.M., Doss, A., Ravi, T.K., Sukumar, M., 2010. Ethnobotanical study of medicinal plants used by Malasar tribals in Coimbatore District of Tamil Nadu (South India). Asian Journal of Experimental Biological Sciences 1, 387–392.
- Verma, S., Chauhan, N.S., 2007. Indigenous medicinal plants knowledge of Kunihat forest division, district Solan. Indian Journal of Traditional Knowledge 6, 494–497.
- Vieira, C., Fetzter, S., Sauer, S.K., Evangelista, S., Averbeck, B., Kress, M., Reeh, P.W., Cirillo, R., Lippi, A., Maggi, C.A., Manzini, S., 2001. Pro- and anti-inflammatory actions of ricinoleic acid: similarities and differences with capsaicin. Naunyn-Schmiedeberg's Archives of Pharmacology 364, 87–95.
- Vijayalakshmi, A., Geetha, M., 2014. Anti-psoriatic activity of flavonoids from *Cassia tora* leaves using the rat ultraviolet B ray photodermatitis model. Revista Brasileira de Farmacognosia 24, 322–329.
- Vijayalakshmi, T., Muthulakshmi, V., Sachdanandam, P., 2000. Toxic studies on biochemical parameters carried out in rats with *Serankottai nei*, a siddha drug-milk extract of *Semecarpus anacardium* nut. Journal of Ethnopharmacology 69, 9–15.
- Vijayan, A., Liju, V.B., John, R., Parthipan, B., Renuka, C., 2007. Traditional remedies of Kani tribes of Kottor reserve forest, Agasthyavanam, Thiruvananthapuram, Kerala. Indian Journal of Traditional Knowledge 6, 589–594.
- Vijikumar, S., 2011. *Cuscuta reflexa* Roxb. – wonderful miracle plant in ethnomedicine. Indian Journal of Natural Sciences 11, 676–683.
- Virani, N.V., Chandola, H.M., Vyas, S.N., Jadeja, D.B., 2010. Clinical study on erectile dysfunction in diabetic and non-diabetic subjects and its management with *Ficus religiosa* Linn. AYU 31, 272–279.
- Vohra, S.B., Shah, S.A., Sharma, K., Naqvi, S.A.H., Dandiya, P.C., 1989. Antibacterial, antipyretic, analgesic and anti-inflammatory studies on *Acorus calamus* Linn. Annals of National Academy of Medical Sciences 25, 13–20.
- Vyjayanthi, G., Kulkarni, C., Abraham, A., Kolhapure, S.A., 2004. Evaluation of anti-dandruff activity and safety of polyherbal hair oil: an open pilot clinical trial. The Antiseptic 101, 368–372.
- Wadankar, G.D., Malode, S.N., Sarambekar, S.L., 2011. Traditionally used medicinal plants for wound healing in the Washim District, Maharashtra (India). International Journal of PharmTech Research 3, 2080–2084.
- Wafa, G., Amadou, D., Larbi, K.M., Hela, E.F.O., 2014. Larvicidal activity, phytochemical composition, and antioxidant properties of different parts of five populations of *Ricinus communis* L. Industrial Crops and Products 56, 43–51.
- Wake, R.R., Patil, A.N., Khadabadi, S.S., 2011. In-vitro antimicrobial activity of extracts of seeds of *Cleome viscosa* Linn. International Journal of Pharmaceutical Sciences and Research 2, 2232–2236.
- Wang, Y.K., Hong, Y.J., Wei, M., Wu, Y., Huang, Z.Q., Chen, R.Z., Chen, H.Z., 2010. Curculigolide attenuates human umbilical vein endothelial cell injury induced by H₂O₂. Journal of Ethnopharmacology 132, 233–239.
- Wani, T.A., Chandrashekar, H.H., Kumar, D., Prasad, R., Gopal, A., Sardar, K.K., Tandan, S.K., Kumar, D., 2012a. Wound healing activity of ethanolic extract of *Shorea robusta* Gaertn. F. Resin. Indian Journal of Experimental Biology 50, 277–281.
- Wani, T.A., Chandrashekar, H.H., Kumar, D., Prasad, R., Sardar, K.K., Kumar, D., Tandan, S.K., 2012b. Anti-inflammatory and antipyretic activities of the ethanolic extract of *Shorea robusta* Gaertn. f. resin. Indian Journal of Biochemistry & Biophysics 49, 463–467.
- Warokar, A.S., Ghante, M.H., Duragkar, N.J., Bhusari, K.P., 2010. Anti-inflammatory and antioxidant activities of methanolic extract of *Buchanania lanzan* Kernel. Indian Journal of Pharmaceutical Education and Research 44, 363–368.
- Weber, L.F., 1937. External causes of dermatitis. A list of irritants. Archives of Dermatology and Syphilology 35, 129–179.
- White, J.C., 1887. *Dermatitis venenata*: An Account of the Action of External Irritants Upon the Skin. Cupples and Hurd, Boston.
- Wijaya, R., Neumann, G.M., Condron, R., Hughes, A.B., Polya, G.M., 2000. Defense proteins from seed of *Cassia fistula* include a lipid transfer protein homologue and a protease inhibitory plant defensin. Plant Science 6, 243–255.
- Wojdylo, A., Oszmianski, J., Czemerys, R., 2007. Antioxidant activity and phenolic compounds in 32 selected herbs. Food Chemistry 105, 940–949.
- Wu, C., Liang, C., Shiu, L., Chang, L., Lin, T., Lan, C.E., Tsai, J., Wong, T., Wei, K., Lin, T., Chang, N., Sheu, H., 2011. *Solanum incanum* extract (SR-T100) induces human cutaneous squamous cell carcinoma apoptosis through modulating tumor necrosis factor receptor signalling pathway. Journal of Dermatological Science 63, 83–92.
- Wu, Q., Fu, D., Hou, A., Lei, G., Liu, Z., Chen, J., Zhou, T., 2005. Antioxidative phenols and phenolic glycosides from *Curculigo orchoides*. Chemical and Pharmaceutical Bulletin 53, 1065–1067.
- Yadav, V., Jayalakshmi, S., Singla, R.K., Patra, A., 2011. Preliminary assessment of anti-inflammatory activity of *Callicarpa macrophylla* Vahl. leaves extracts. Indo Global Journal of Pharmaceutical Sciences 1, 219–222.
- Yadav, M., Khan, K.K., Beg, M.Z., 2012. Ethnobotanical plants used for curing skin diseases by tribals of Rewa district (Madhya Pradesh). Indian Journal of Life Sciences 2, 123–126.
- Yadav, V., Jayalakshmi, S., Singla, R.K., Patra, A., 2012. Evaluation of antibacterial activity of *Callicarpa macrophylla* Vahl. stem extracts. WebmedCentral Ayurvedic Medicine 3, WMC003651.
- Yadava, R.N., Tiwari, L., 2007. New antifungal flavone from *Butea monosperma*. Journal of Enzyme Inhibition and Medicinal Chemistry 22, 497–500.
- Yesmin, M.N., Uddin, S.N., Mubassara, S., Akond, M.A., 2008. Antioxidant and antibacterial activities of *Calotropis procera* Linn. American-Eurasian Journal of Agricultural & Environmental Sciences 4, 550–553.
- Yisa, J., 2009. Phytochemical analysis and antimicrobial activity of *Scoparia dulcis* and *Nymphaea lotus*. Australian Journal of Basic and Applied Sciences 3, 3975–3979.
- Yogesh, H.S., Kichadi, S.C., Muchandi, I.S., Gopalakrishna, B., 2013. Evaluation of anti-inflammatory activity of *Cynodon dactylon* Pers. on carrageenan induced paw edema in rats. Indian Journal of Natural Products and Resources 4, 151–154.
- Yogisha, S., Samiulla, D.S., Prashanth, D., Padmaja, R., Amit, A., 2002. Trypsin inhibitory activity of *Lawsonia inermis*. Fitoterapia 73, 690–691.
- Yusufoglu, H.S., 2011. Topical anti-inflammatory and wound healing activities of herbal gel of *Ziziphium nummularia* L. (F. Rhamnaceae) leaf extract. International Journal of Pharmacology 7, 862–867.

- Zachariah, S.M., Aleykutty, N.A., Viswanad, V., Jacob, S., Prabhakar, V., 2011. In-vitro antioxidant potential of methanolic extracts of *Mirabilis jalapa* Linn. Free Radicals and Antioxidants 1, 42–86.
- Zade, V., Wikhe, M., Dabhadkr, D., Dawada, S., Patil, U., 2013. Antifertility efficacy of *Cannabis sativa* leaves on female albino rats. The International Journal of Science Inventions Today 2, 107–117.
- Zaveri, M., Jain, S., 2010. Anti-inflammatory and analgesic activity of root bark of *Oroxylum indicum*, Vent. Journal of Global Pharma Technology 2, 79–87.
- Zeashan, H., Amresh, G., Singh, S., Rao, C.V., 2009. Hepatoprotective and antioxidant activity of *Amaranthus spinosus* against CCl₄ induced toxicity. Journal of Ethnopharmacology 125, 364–366.
- Zhang, H., Jiang, L., Ye, S., Ye, Y., Ren, F., 2010. Systematic evaluation of antioxidant capacities of the ethanolic extract of different tissues of jujube (*Ziziphus jujube* Mill.) from China. Food and Chemical Toxicology 48, 1461–1465.
- Zhang, M., Chen, Y., 1996. Chemical constituents of *Eclipta alba* (L.) Hassk. Zhongguo Zhong Yao Za Zhi 21, 480–481.
- Zhenbao, J., Fei, T., Ling, G., Guanjin, T., Xiaolin, D., 2007. Antioxidant properties of extracts from juemingzi (*Cassia tora* L.) evaluated *in vitro*. LWT-Food Science and Technology 40, 1072–1077.
- Zhou, Y., Liu, Y.E., Cao, J., Zeng, G., Shen, C., Li, Y., Zhou, M., Chen, Y., Pu, W., Potters, L., Shi, Y.E., 2009. Vitexins, nature-derived lignan compounds, induce apoptosis and suppress tumor growth. Clinical Cancer Research 15, 5161–5169.
- Zohri, A.N., Abdel-Gawad, K., Saber, S., 1995. Antibacterial, antidermatophytic and antitoxigenic activities of onion (*Allium cepa* L.) oil. Microbiological Research 150, 167–172.